arXiv:hep-th/9308146v1 30 Aug 1993

Finite and infinite W algebras

and their applications

Tjark Tjin

Supervisor: Prof. dr. ir. F. A. Bais

The various chapters of this thesis are based on the following papers:

- Chapter 3: T. Tjin, Finite W algebras, *Phys. Lett* **B292** (1992)60; J. de Boer and T. Tjin, Quantization and representation theory of finite W algebras, *Comm. Math. Phys.* to appear.

- Chapter 4: F. A. Bais, T. Tjin, P. van Driel, Covariantly coupled chiral algebras, *Nucl. Phys.* **B357** (1991) 632; J. de Boer and T. Tjin, The relation between quantum W algebras and Lie algebras, *Comm. Math. Phys.* to appear; F. A. Bais, T. Tjin, P. van Driel, J. de Boer, J. Goeree, W algebras, W gravities and their moduli spaces, To be published in the proceedings of the 'workshop on Superstrings and related Topics', Trieste, Juli 1992.

- Chapter 5: T. Tjin, P. van Driel, Coupled WZNW-Toda models and Covariant KdV hierarchies, Preprint ITFA-91-04; T. Tjin, Finite W symmetry in finite dimensional integrable systems, To be published in the proceedings of the workshop on 'Low dimensional Topology and Physics', Isaac Newton Institute, Cambridge, 1992.

Contents

1 General Introduction **3**

1.1 Conformal Field Theory and W algebras 4

1.2 String theories of elementary particle physics 17

1.3 The Quantum Hall effect 25

1.4 Phase transitions in 2 dimensions 28

1.5 Solitary wave phenomena 31

1.6 Outline of the thesis 37

2 Classical and quantum reduction of WZW models **39**

2.1 Constrained WZW models 40

2.2 Reduction of Poisson manifolds 43

2.3 Gauged WZW models 47

2.4 Quantum framework for constrained WZW models 48

3 Finite W algebras **51**

3.1 The Classical Theory of Finite W algebras 53

3.1.1 Kirillov Poisson structures 53

3.1.2 sl_2-embeddings 55

3.1.3 Reductions associated to sl_2 embeddings 59

3.1.4 Generalized finite Miura transformations 69

3.1.5 Examples 72

3.1.6 An algebraic description of finite W algebras 76

3.1.7 The symplectic geometry of finite W algebras 81

3.2 Quantum finite W Algebras 85

3.2.1 BRST quantization of finite W algebras 86

3.2.2 Spectral Sequences for Double Complexes 88
3.2.3 The BRST Cohomology 90
3.2.4 Example 93
3.3 The Representation Theory of Finite W algebras 94
3.3.1 The representation theory of $\bar{W}_3^{(2)}$ 95
3.3.2 Induced represention s 100
3.3.3 Fock realizations 104
3.4 Relation with Lie algebra cohomology 108

4 W algebras from affine Lie algebras 117
4.1 Classical W algebras 119
4.1.1 Generalized Drinfeld-Sokolov reduction 119
4.1.2 An algebraic description of W algebras 130
4.2 Quantum W algebras 134
4.2.1 Quantization of generalized DS reductions 134
4.2.2 Examples 140

5 W symmetry in integrable systems 148
5.1 Generalized finite Toda systems 149
5.2 Generalized Toda field theory 152
5.3 Covariant KdV type hierarchies 154
5.4 Discussion 161

Chapter 1

General Introduction

In this introductory chapter we give a brief review of conformal field theory (CFT), define the concept of a W algebra and discuss some of their potential applications to high energy physics, condensed matter physics and solitary wave phenomena. In 2 dimensions the conformal group is infinite dimensional and its Lie algebra is the Virasoro algebra. The particular structure of the representation theory of this algebra has lead to a partial classification of CFT's but it has turned out that for a complete classification it is necessary to consider *extensions* of conformal symmetry. Such extensions are called W algebras. In the first section of this chapter we review some basic aspects of CFT and W algebras in order to set the stage. Having developed the necessary background we consider string theory which is the most promising candidate for a unified theory of the four known forces of nature and its particle spectrum. In this context CFT's arize as perturbative groundstates around which one expands the theory. As another promising application we consider some recent work on the role played by CFT in the quantum Hall effect. The crucial insight here is that (as first proposed by Laughlin) in the Hall state the system behaves like an incompressible quantum fluid such that the only nontrivial dynamics is concentrated around the boundary of the sample. The boundary theory can then be described by a 2 dimensional CFT. Next we discuss the result that 2 dimensional lattice models, such as the Ising model, at criticality correspond to 2 dimensional CFT's. This relation is based on the fact that at criticality fluctuations in the system are correlated over long distances (i.e. the correlation length goes to infinity) which allows one to define a continuum limit in which the lattice system corresponds to a 2 dimensional quantum field theory. Due to the scale covariance in critical lattice systems this field theory will by conformally invariant and the critical exponents will be determined by the representation theory of the W algebra underlying the CFT. In the last section of this chapter we discuss the role played by W algebras in the theory of integrable hierarchies with solitary wave solutions. Here W algebras appear in two distinct ways: first of all through the second Hamiltonian structures and also through

the so called weak action angle structure which is intimately related to the integrability of the hierarchy.

We hope that with this introductory chapter we have provided the reader with sufficient motivation to proceed to the more technical chapters in which we study W algebras in detail.

1.1 Conformal Field Theory and W algebras

The distinctive feature of the conformal group in 2 dimensions is that it is, in contrast with other dimensions, infinite dimensional. It turns out that this infinite symmetry has dramatic consequences as was first shown in [8]. In this section we shall give a brief account of this vast subject. For more detailed reviews we refer the reader to [12, 37].

In what follows we shall always take spacetime to be $M = \mathbf{R} \times S^1$ with coordinates $(x^0, x^1) \equiv (t, x)$ and metric $ds^2 = dt^2 - dx^2$. Here we think of space to be compactified, i.e $x = x + 2\pi$. A conformal transformation is a diffeomorphism (i.e. a smooth and invertible map) from spacetime to itself that preserves the causal structure (that is the lightlike lines are mapped to lighlike lines and the orientation of time is preserved). The set of lightlike lines on M is isomorphic to $S^1 \times S^1$ and from this it is easy to see that the set of conformal diffeomorphisms is given by $Diff_c(M) = Diff(S^1) \times Diff(S^1)$ where in light cone coordinates the first factor corresponds to transformations $x^+ \mapsto \hat{x}^+(x^+)$ and the second factor corresponds to $x^- \mapsto \hat{x}^-(x-)$. Note that in these coordinates the metric has the form $ds^2 = dx^+ dx^-$ and is form invariant (up to a scalar factor) under $Diff_c(M)$. Let $\epsilon = \epsilon(x^\pm)$ be a (small) function and let A be a smooth function of $x^\pm$, then one has

$$A(x^\pm + \epsilon(x^\pm)) = \exp(\sum_n \epsilon_n e^{inx^\pm} \frac{d}{dx^\pm}) A(x^\pm) \tag{1.1.1}$$

where ϵ_n are the Fourier coefficients of $\epsilon(x^\pm)$. From this follows that the space of conformal Killing vectors is spanned by

$$X_n^\pm = e^{inx_\pm} \frac{d}{dx_\pm} \tag{1.1.2}$$

which have the following commutation relations

$$[X_n^\pm, X_m^\pm] = -i(n-m) X_{n+m}^\pm \tag{1.1.3}$$

Now, let $\mathcal{F}$ be the 'algebra of fields' of a certain relativistic quantum field theory on M with Hilbert space $\mathcal{H}$ (i.e. $\mathcal{F}$ consists of operator valued distributions $\phi(x)$ over

M such that $[\phi_1(x), \phi_2(y)] = 0$ if x and y are spacelike separated. Furthermore the spectrum of the energy momentum operators P^μ is concentrated in the closed forward light cone $\bar{V}^+$ ($p^2 \geq 0;\ p^0 = E \geq 0$) and the Hilbert space $\mathcal{H}$, which is an irreducible representation of $\mathcal{F}$, contains precisely one state, called the physical vacuum, which is unvariant under the unitary representation of the Poincare group that $\mathcal{H}$ carries). Physical operators like P^μ are usually unbounded and can therefore not be defined on the whole Hilbert space. As usual we therefore assume the existence of a common domain of definition $\mathcal{D} \subset \mathcal{H}$ for all fields which is dense in $\mathcal{H}$.

In a CFT the Hilbert space will carry a unitary projective representation of the conformal group. This means that any element $f \in Diff_c(M)$ will be represented by a unitary operator $U(f)$ which is determined up to a phase. If the theory is conformally invariant the representation U must give rise to a representation R of the conformal group on the field algebra $\mathcal{F}$ such that

$$U(f)\phi(x)U(f)^{-1} = R_f(\phi(f(x))) \tag{1.1.4}$$

for any $f \in Diff_c(M)$ and $\phi \in \mathcal{F}$. Equation (1.1.4) expresses the covariance of the theory under conformal transformations.

Let us consider an infinitesimal conformal transformation generated by a conformal Killing vector X on M. of course it will be represented on $\mathcal{H}$ by a Hermitean operator $T(X)$ which, due to the projectivity of the representation, is determined up to a constant. The fact that T is a (projective) representation of the Lie algebra of conformal Killing vectors is expressed by the equation

$$[T(X), T(X')] = T([X, X']) + c(X, X')\mathbf{1} \tag{1.1.5}$$

where $c(X, X')$ is a 'central extension'. For an 'ordinary' representation this term would have been absent. Defining $L_n = T(X_n^+)$ and $\bar{L}_n = T(X_n^-)$ and using equations (1.1.3) and (1.1.5) one finds

$$\begin{aligned} [L_n, L_m] &= (n-m)L_{n+m} + \frac{c}{12}m(m^2-1)\delta_{n+m,0} \\ [\bar{L}_n, \bar{L}_m] &= (n-m)\bar{L}_{n+m} + \frac{\bar{c}}{12}m(m^2-1)\delta_{n+m,0} \\ [\bar{L}_n, L_m] &= 0 \end{aligned} \tag{1.1.6}$$

where the central terms are fixed up to scalars $c, \bar{c}$ by the Jacobi identities. In fact Lorentz invariance requires c and $\bar{c}$ to be equal. We conclude the Hilbert space of any CFT carries a representation of $Vir \oplus Vir$.

From eq. (1.1.2) follows that $\frac{d}{dt} = X_0^+ + X_0^-$ while $\frac{d}{dx} = X_0^+ - X_0^-$ which means that $L_0 \pm \bar{L}_0$ are the generators of time and spatial translations respectively in $\mathcal{H}$.

Therefore $L_0 + \bar{L}_0$ must be equal to the Hamiltonian $H \equiv P^0$ of the system while $L_0 - \bar{L}_0$ corresponds to the momentum operator $P \equiv P^1$. Furthermore since space is compact we have $\phi(x + 2\pi) = \phi(x)$ which means that the operator implementing spatial translations by 2π given by $e^{2\pi iP}$ commutes with all fields in $\mathcal{F}$. Since $\mathcal{H}$ is an irreducible representation of $\mathcal{F}$ we find by Schur's lemma that $e^{2\pi iP}$ must be a multiple of the identity operator, i.e. $e^{2\pi iP} = \alpha\mathbf{1}$. Let $|p\rangle$ be a momentum eigenstate $P|p\rangle = p|p\rangle$, then this implies that $e^{2\pi ip}|p\rangle = \alpha|p\rangle$ and furthermore by taking $|p\rangle = |0\rangle$ we see that $\alpha = 1$. We conclude that the spectrum of $P = L_0 - \bar{L}_0$ is a subset of $\mathbf{Z}$ (i.e. it is quantized). In CFT the eigenvalues of $L_0 - \bar{L}_0$ are called the 'conformal spins'.

It can be shown [37] that $T(X)$ can be written as

$$T(X) = \frac{1}{2\pi} \int_\Sigma X^\mu T_{\mu\nu} d\sigma^\nu \tag{1.1.7}$$

where $X = X^\mu \partial_\mu$ is a conformal Killing vector and Σ is some spacelike curve. In (1.1.7) we have implicitly fixed the additive freedom in $T(X)$ by fixing the trace of $T_{\mu\nu}$ to be zero, $T^\mu_\mu = 0$. Furthermore varying Σ we find the conservation law $\partial^\mu T_{\mu\nu} = 0$. In light cone coordinates, and using the tracelessness of $T_{\mu\nu}$ this equation reads $\partial_- T_{++} = 0$ and $\partial_+ T_{--} = 0$, i.e. the only two nonzero components of $T_{\mu\nu}$ depend on one light cone variable only. This is an extremely strong statement for it implies that for any $f = f(x^+)$ one has the conservation law $\partial_-(fT_{++}) = 0$ (and similar equations for T_{--}). The charge

$$Q_f = \int_{S^1} dx\ f(x^+) T_{++}(x^+) \tag{1.1.8}$$

is then likewise conserved. Since f is arbitrary we conclude that in 2 dimensions conformal invariance leads to an *infinite set of independent conserved quantities*. Note that from eq.(1.1.7) immediately follows that $T_{++}(x^+) = \sum_n L_n e^{-inx_+}$ and $T_{--}(x^-) = \sum_n \bar{L}_n e^{-inx^-}$. It is well known that $T_{\mu\nu}$ can be identified with the stress energy tensor of the theory.

Let $|E, p\rangle \in \mathcal{H}$ be an eigenstate of H and P with energy E and momentum p. It is easy to check, using the commutation relations (1.1.6), that $L_n|E,p\rangle$ and $\bar{L}_n|E,p\rangle$ are also eigenstates of H (and P), but with energy $E - n$. Therefore the operators L_n and $\bar{L}_n$ lower the energy for positive n. Since the energy is bounded from below (remember that $p^0 \geq 0$) there must exist a state $|E_0, p_0\rangle$ such that $L_n|E_0,p_0\rangle = \bar{L}_n|E_0,p_0\rangle = 0$ for all $n > 0$. Define $h = \frac{1}{2}(E_0 + p_0)$ and $\bar{h} = \frac{1}{2}(E_0 - p_0)$, called 'conformal dimensions', then from $H = L_0 + \bar{L}_0$ and $P = L_0 - \bar{L}_0$ follows that $L_0|E_0,p_0\rangle = h|E_0,p_0\rangle$ and $\bar{L}_0|E_0,p_0\rangle = \bar{h}|E_0,p_0\rangle$. Therefore denote $|E_0,p_0\rangle \equiv |h,\bar{h}\rangle$. The linear span of all states that one can obtain by successively acting with the generators $\{L_{-n}\}_{n>0}$ on $|h,\bar{h}\rangle$ is an irreducible Virasoro representation denoted by $L(h,c)$. The same is true when we act with the generators $\{\bar{L}_{-n}\}_{n>0}$ on $|h,\bar{h}\rangle$ but now we obtain the representation $L(\bar{h},c)$.

In this way the Hilbert space $\mathcal{H}$ decomposes into irreducible $Vir \oplus Vir$ representations

$$\mathcal{H} = \bigoplus_{i,\bar{i}} N_{i\bar{i}}\, \mathcal{H}_i \otimes \mathcal{H}_{\bar{i}} \tag{1.1.9}$$

where $\mathcal{H}_i = L(h_i, c)$, $\mathcal{H}_{\bar{i}} = L(\bar{h}_{\bar{i}}, c)$ and $N_{i\bar{i}}$ are multiplicities (sometimes the set of these multiplicities is called the 'operator content' of the CFT). The states $|h, \bar{h}\rangle$ which satisfy $L_n|h, \bar{h}\rangle = \bar{L}_n|h, \bar{h}\rangle = 0$ for $n > 0$ are called 'highest weight states'. They are the states in the Virasoro irreducible representations with lowest energy. of course the (unique) vacuum state $|0>$ of the theory is such a highest weight state with $h = \bar{h} = 0$ and the Virasoro representation $L(0, c)$ is called the 'vacuum representation'.

Note that one only needs to know the set of highest weight states to *completely diagonalize* the Hamiltonian, for applying the Virasoro generators to them one can find a basis of the Hilbert space consisting only of eigenvectors of the Hamiltonian. Also note that since the energy momentum eigenvalues are restricted to $\bar{V}^+$ (i.e. $E^2 - p^2 \geq 0$) it follows that $E \geq |p|$. From this it is easy to show that $h_i, \bar{h}_{\bar{i}} \geq 0$ for all $i, \bar{i}$. Therefore the Virasoro representations appearing in CFT are the so called unitary irreducible 'highest weight' modules. In order to classify conformal field theories one would like to find a list of all Virasoro representations of this type for given c.

It turns out [58] that the Virasoro algebra only has unitary representations for $c > 0$. For $c > 0$ we have to distinguish between two possibilities: $0 < c < 1$ and $c \geq 1$. For $c \geq 1$ there exists a unique unitary irreducible Virasoro representation $L(h, c)$ for any positive and real h. For $0 < c < 1$ the situation is drastically different however. To begin with there only exist unitary irreducible Virasoro representations for

$$c = c(m) = 1 - \frac{6}{m(m+1)} \tag{1.1.10}$$

where $m \in \{3, 4, \ldots\}$. Furthermore the irreducible representation $L(c(m), h)$ is only unitary for

$$h = h^{(m)}(r, s) = \frac{((m+1)r - ms)^2 - 1}{4m(m+1)} \tag{1.1.11}$$

where $1 \leq r \leq m-1$ and $1 \leq s \leq m$. Note the remarkable fact that for $c = c(m)$ there are only a *finite* number of values for h that lead to unitary Virasoro representations. From this follows that if we are given a conformally invariant quantum field theory and by explicit calculation we find the central value c to be one of the values $c(m)$ then the direct sum decomposition (1.1.9) is finite if the multiplicities $N_{i\bar{i}}$ are finite (in fact for many conformal field theories that have been studied they are either 0 or 1). For conformal field theories with $c \geq 1$ the decomposition (1.1.9) can be infinite even if the multiplicities are all finite.

We are now in a position to roughly describe the idea of a W algebra. Suppose the Virasoro algebra is just a subalgebra of the total symmetry algebra of the quantum field theory. With respect to this larger algebra the Hilbert space may have a finite decomposition even if the decomposition w.r.t. Virasoro algebra is infinite. A conformal field theory with such a larger symmetry is called a 'rational conformal field theory'. Before we introduce W algebras more formally we still have to introduce the concept of a 'primary field'.

Consider the (infinitesimal) action of the conformal Killing vectorfields $X^{\pm}_{\pm 1}, X^{\pm}_0$ on $\mathcal{F}$. Define the operators $\mathcal{L}_{\pm}, \mathcal{L}_0$ (and their barred counterparts) by

$$\begin{aligned} [L_{-1}, \phi(x^+, x^-)] &= e^{-ix^+}(\mathcal{L}_{-1}\phi)(x^+, x^-) \\ [L_0, \phi(x^+, x^-)] &= (\mathcal{L}_{-1} + \mathcal{L}_0)\phi(x^+, x^-) \\ [L_1, \phi(x^+, x^-)] &= e^{ix^+}(\mathcal{L}_{-1} + 2\mathcal{L}_0 + \mathcal{L}_1)\phi(x^+, x^-) \end{aligned} \tag{1.1.12}$$

These equations are nothing but infinitesimal versions of eq. (1.1.4). It can now be shown [37] that the field algebra $\mathcal{F}$ decomposes into a direct sum of $\mathcal{L}_0$ and $\bar{\mathcal{L}}_0$ eigenspaces $\mathcal{F}^{(h,\bar{h})} = \{\phi \in \mathcal{F} \mid \mathcal{L}_0\phi = h\phi\,;\ \bar{\mathcal{L}}_0\phi = \bar{h}\phi\}$. An element of $\mathcal{F}^{(h,\bar{h})}$ is called a 'field of left conformal dimension h and right conformal dimension $\bar{h}$'. Furthermore one can show that if $\phi \in \mathcal{F}^{(h,\bar{h})}$ and

$$\phi(x^+, x^-) = \sum_{k,\bar{k}} e^{-ikx^+ - i\bar{k}x^-}\phi_{k,\bar{k}} \tag{1.1.13}$$

is its Fourier expansion, then $\phi_{k,\bar{k}}|0>= 0$ for $k > -h$ or $\bar{k} > -\bar{h}$ and also that the map $F : \mathcal{F} \to \mathcal{H}$ defined by $F(\phi) = \phi_{-h,-\bar{h}}|0>$ is an isomorphism (to be completely precise the image of F is the common domain $\mathcal{D}$ of definition of all fields in $\mathcal{F}$).

As we have seen the Hilbertspace contains a set of states that through the Virasoro algebra generate all other states. We called these the highest weight states $|h, \bar{h}\rangle$. The inverse image under the isomorphism F of such a highest weight state is called a *primary field*. Using the fact that $\mathcal{L}_0 = F^{-1}L_0F$ one can easily show that $\mathcal{L}_0\phi = h\phi$ which means that the primary field $\phi \equiv F^{-1}(|h, \bar{h}\rangle)$ is an element of $\mathcal{F}^{(h,\bar{h})}$ as was to be expected. Also, expanding a primary field into Fourier modes as in (1.1.13) one finds that

$$[L_n, \phi_{k,\bar{k}}] = (n(h-1) - k)\phi_{k+n,\bar{k}} \tag{1.1.14}$$

and a similar equation for the commutations relation of $\phi_{k,\bar{k}}$ with $\bar{L}_n$. Equation (1.1.14) expresses the transformation properties of a primary field under infinitesimal conformal transformations. Integrating this equation we obtain

$$U(f)\phi(x)U(f)^{-1} = \left(\frac{df^+}{dx^+}\right)^h \left(\frac{df^-}{dx^-}\right)^{\bar{h}} \phi(f(x)) \tag{1.1.15}$$

where $f \equiv (f^+, f^-) \in Diff_c(M)$. Fields of conformal dimensions $(h, 0)$ are called 'left chiral' and do not depend on x^- while fields of conformal dimensions $(0, \bar{h})$ are called 'right chiral' and do not depend on x^+. Obviously a field of conformal dimensions $(0, 0)$ is constant and conversely a constant has conformal dimensions $(0, 0)$.

Consider the operators $\mathcal{L}_n = F^{-1} L_n F$ and their barred counterparts $\bar{\mathcal{L}}_n = F^{-1} \bar{L}_n F$ which map the field algebra $\mathcal{F}$ into itself. Obviously we have $\mathcal{L}_n \phi = 0$ if ϕ is a primary field and $n > 0$. Define now the conformal family $[\phi]$ associated to the primary field ϕ to be the subset of $\mathcal{F}$ spanned by fields of the form

$$\phi^{\{p\}\{\bar{p}\}} \equiv \mathcal{L}_{-p_1} \dots \mathcal{L}_{-p_k} \bar{\mathcal{L}}_{-\bar{p}_1} \dots \bar{\mathcal{L}}_{-\bar{p}_l} \, \phi \tag{1.1.16}$$

where $p_1 \geq \dots \geq p_k \geq 0$; $\bar{p}_1 \geq \dots \geq \bar{p}_l \geq 0$; $k, l \geq 0$ and $\{p\} = (p_1, \dots, p_k)$. The fields (1.1.16) are called 'descendent fields'. Note that $[\phi] \simeq [\phi]_L \otimes [\phi]_R$ where $[\phi]_L$ is spanned by $\phi^{\{p\}\{0\}}$ and $[\phi]_R$ by $\phi^{\{0\}\{\bar{p}\}}$. Furthermore from eq.(1.1.9) follows that

$$\mathcal{F} = \bigoplus_{i,\bar{i}} N_{i,\bar{i}} \, [\phi^{(h_i, \bar{h}_{\bar{i}})}]_L \otimes [\phi^{(h_i, \bar{h}_{\bar{i}})}]_R \tag{1.1.17}$$

where $\phi^{(h,\bar{h})}$ is a primary field with conformal dimensions $(h, \bar{h})$.

The subalgebras $\mathcal{A}$ and $\bar{\mathcal{A}}$ of $\mathcal{F}$ defined by

$$\mathcal{A} = \bigoplus_h \mathcal{F}^{(h,0)} \quad \text{and} \quad \bar{\mathcal{A}} = \bigoplus_{\bar{h}} \mathcal{F}^{(0,\bar{h})} \tag{1.1.18}$$

are called the 'left and right *chiral algebras*' of the conformal field theory respectively. $\mathcal{A}$ and $\bar{\mathcal{A}}$ commute for if $\phi_L = \phi_L(x^+) \in \mathcal{A}$ and $\phi_R = \phi_R(y^-) \in \bar{\mathcal{A}}$ then by using invariance of $\phi_L(\phi_R)$ under translations in x^- (respectively x^+) one can always spatially separate (x^+, x^-) and (y^+, y^-) which means that the fields must commute. From this follows that the Hilbert space decomposes into a direct sum of irreducible $\mathcal{A} \otimes \bar{\mathcal{A}}$ representations

$$\mathcal{H} = \bigoplus_{(a,\bar{a})} N^{\mathcal{A}}_{a,\bar{a}} \, \mathcal{H}^{\mathcal{A}}_a \otimes \bar{\mathcal{H}}^{\bar{\mathcal{A}}}_{\bar{a}} \tag{1.1.19}$$

Note that for a CFT the chiral algebras always contain the Virasoro algebra due to the fact that, as we have seen, $T_{++} \in \mathcal{A}$ and $T_{--} \in \bar{\mathcal{A}}$. It can therefore happen that the decomposition (1.1.19) is finite even if the decomposition (1.1.9) is infinite. In that case the CFT is called a rational conformal field theory (RCFT). A W algebra [62] is a chiral algebra that is generated (in a way that we shall explain later) by a set of chiral primary fields.

In the literature CFT is usually formulated within the framework of Euclidean field theory (this is actually quite natural from the point of view of the applications). First one can extend the fields defined on Minkowski space M to fields in two complex variables by putting

$$\phi(\xi,\bar{\xi}) = e^{iP.\Delta x}\phi(x^0,x^1)e^{-iP.\Delta x} \tag{1.1.20}$$

where $\Delta x = (iy^0, iy^1)$, $\xi = x^0 - iy^0$ and $\bar{\xi} = x^1 + iy^1$. The Euclidean domain is defined as usual by $x^0 = y^1 = 0$ and $y^0 = \tau$ which effectively implies a Wick rotation $t \to -i\tau$. It is more or less customary in CFT to work in the so called 'radial picture' since it allows one to use the powerful theory of complex functions and contour integrals. For this define $z = e^{i(\xi+\bar{\xi})}$ and $\bar{z} = e^{i(\xi-\bar{\xi})}$. In the Euclidean domain we have $z = e^{\tau+ix^1}$ and $\bar{z} = e^{\tau-ix^1}$ which means that $z^* = \bar{z}$. Also the infinite past and future are mapped to $z = 0$ and $z = \infty$ respectively on the plane and equal time surfaces become circles of constant radius. The Wick rotated spacetime $M = \mathbf{R} \times S^1$ has become the complex plane with the origin removed. Furthermore left and right chiral fields depend holomorphically and antiholomorphically on z respectively.

Usually however one leaves the restriction to the Euclidean domain until after one has developed the theory. We are therefore dealing with a theory on $\mathbb{C}^2$ in which z and $\bar{z}$ are considered to be independent complex variables. It is then often convenient to introduce slightly different fields when working in the radial picture. Let $\phi(x^0,x^1)$ be a field and extend it to $\mathbb{C}^2$, $\phi = \phi(z,\bar{z})$. Then define the field $\tilde{\phi}(z,\bar{z})$ by $\phi(z,\bar{z}) = z^h\bar{z}^{\bar{h}}\tilde{\phi}(z,\bar{z})$. It can now easily be seen that the 'radial field' $\tilde{\phi}$ has the expansion

$$\tilde{\phi}(z,\bar{z}) = \sum_{k,\bar{k}} z^{-k-h}\bar{z}^{-\bar{k}-\bar{h}}\tilde{\phi}_{k,\bar{k}} \tag{1.1.21}$$

and furthermore that $\tilde{\phi}_{k,\bar{k}}|0>=0$ for $k > -h$ or $\bar{k} > -\bar{h}$. From now on we will omit the explicit $\bar{z}$ dependence of fields where possible.

In Euclidean field theory products of operators $A(z)B(w)$ are only defined for $|z| > |w|$ (Euclidean correlation functions are only defined if the operators in the correlator are time ordened. Remember that in the radial picture $|z| > |w|$ means that z is later than w. of course in the Euclidean functional integral definition of correlation functions time ordening is automatic). Therefore define the radial ordening operator R as follows: $R(A(z)B(w)) = A(z)B(w)$ if $|z| > |w|$, and $R(A(z)B(w)) = B(w)A(z)$ if $|z| < |w|$.

Now consider the commutator between two fields $A(z)$ and $B(z)$. Without loss of generality we can take $A(z)$ and $B(z)$ to be fields of left conformal dimensions h_A and h_B respectively. Again using translational invariance of these fields in $\bar{z}$ it is easy to show that $[A(z),B(w)]$ is zero unless $z = w$. Therefore the equal time commutator can

be expressed as a linear combination of derivatives of $\delta(z-w)$, i.e.

$$[A(z),B(w)] = \sum_{n=1}^{h_A+h_B} \{AB\}_{(n)}(w)\frac{(-\partial)^{n-1}}{(n-1)!}\delta(z-w) \tag{1.1.22}$$

where $\{AB\}_{(n)}(z)$ are elements of $\mathcal{F}$. The reason that the summation runs only up to h_A+h_B is that by performing a scale transformation on this equation one finds that the field $\{AB\}_{(n)}$ has conformal dimension h_A+h_B-n. Since there exist no fields with negative conformal dimension the summation terminates at h_A+h_B. The left hand side of the equal time commutator (1.1.22) can be written as

$$[A(z),B(w)] = \left(\lim_{|z|>|w|} - \lim_{|z|<|w|}\right) R(A(z)B(w)) \tag{1.1.23}$$

There exists a similar expression for the delta function.

$$\delta(z-w) = \left(\lim_{|z|>|w|} - \lim_{|z|<|w|}\right) \frac{1}{z-w} \tag{1.1.24}$$

From the above follows that the functions $A(z)B(z) - \sum_n \{AB\}_{(n)}(w)(z-w)^{-n}$ and $B(z)A(z) - \sum_n \{AB\}_{(n)}(w)(z-w)^{-n}$ are holomorphic for $|z|>|w|$ and $|z|<|w|$ respectively. Since they have the same (boundary) value at $|z|=|w|$ we find that they are holomorphic for all z and are the same function. One can now expand this function in a power series. The result is

$$R(A(z)B(w)) = \sum_{k=h_A+h_B}^{-\infty} \frac{\{AB\}_{(k)}(w)}{(z-w)^k} \tag{1.1.25}$$

i.e. the product of two fields can be expressed as a power series in terms of $z-w$. This is called the *operator product expansion* (OPE).

Let $\{\phi_n(z,\bar{z})\}_{n\in I}$ be the set of primary fields in a certain CFT. From eq.(1.1.17) follows that we can expand the OPE between ϕ_n and ϕ_m as follows

$$\phi_n(z,\bar{z})\phi_m(w,\bar{w}) = \sum_k \sum_{\{p\}\{\bar{p}\}} C^k_{nm;\{p\}\{\bar{p}\}}(z-w,\bar{z}-\bar{w})\,\phi_k^{\{p\}\{\bar{p}\}}(w,\bar{w}) \tag{1.1.26}$$

Define the number N^k_{nm} to be 1 if $C^k_{nm;\{p\}\{\bar{p}\}} \neq 0$ for some $\{p\},\{\bar{p}\}$ and 0 otherwise. Obviously these numbers encode whether in the OPE of ϕ_n and ϕ_m there appear elements

of the conformal family of ϕ_k. It can be shown that a three point function $\langle\phi_n\phi_m\phi_k\rangle$ can only be nonvanishing if $N_{nm}^k \neq 0$ which means that N_{nm}^k encodes which interactions are present in the theory. It is usual to denote this information in terms of the so called 'fusion rules'

$$[\phi_n] \cdot [\phi_m] = \sum_k N_{nm}^k [\phi_k] \tag{1.1.27}$$

which obviously is a shorthand notation for which conformal families interact with which. Given the unitary representations of a certain chiral algebra $\mathcal{A}$ the fusion rules provide a useful characterization of the associated CFT. The calculation of the fusion rules of conformal field theories with extended symmetries is still an open problem at this time.

Using the OPE between A and B we can calculate the commutator between their modes which are defined by

$$A_n = \oint_0 \frac{dz}{2\pi i} z^{n+h_A-1} A(z) \tag{1.1.28}$$

and a similar equation for B_n. The commutator $[A_n, B_m]$ is the difference of two terms both containing two contour integrals around the origin. The difference of two of those contours can be replaced by a contour surrounding w [12]. One then obtains

$$[A_n, B_m] = \oint_0 \frac{dw}{2\pi i} w^{n+h_B-1} \oint_w \frac{dz}{2\pi i} z^{n+h_A-1} R(A(z)B(w)) \tag{1.1.29}$$

Inserting eq.(1.1.25) one can calculate from this the commutator between A_n and B_m.

As usual the vacuum expectation value $< 0|A(z)B(w)|0 >$ of a product of two fields diverges when $z \to w$. We therefore define a normal ordened product as follows: Let the left conformal dimensions of $A(z)$ and $B(z)$ be h_A and h_B respectively, then the normal ordened product, denoted by $(AB)(z)$, is defined by $(AB)(z) = A_-(z)B(z) + B(z)A_+(z)$ where $A_-(z) = \sum_{n \leq -h_a} A_n z^{-n-h_A}$ and $A_+ = A - A_-$. If

$$A(z)B(w) = \sum_{k=h_a+h_B}^{-\infty} \frac{\{AB\}_{(k)}(w)}{(z-w)^k} \tag{1.1.30}$$

is the OPE between A and B then it can easily be shown that $(AB)(z) = \{AB\}_{(0)}(z)$.

Let $\{A_i(z)\}$ be a certain set of chiral fields. The algebra they generate is the subset of $\mathcal{F}$ that can be obtained by taking arbitrary linear combinations, derivatives and normal ordened products of these fields. A W algebra is then a chiral algebra which is generated by $T(z)$ and a set of chiral primary fields. These algebras play an important role in CFT for reasons that we explained above.

of course the simplest example of a W algebra is the Virasoro algebra, i.e. the chiral algebra generated by $T(z)$ alone. Let us now look at some extensions of this algebra. A very important example, is the algebra generated by the fields $T(z)$ and $\{J^a(z)\}$, where the index a runs over a basis of a simple Lie algebra g, and the fields $J^a(z)$ are all chiral primary fields of conformal dimension 1. The OPEs of the fields $J^a(z)$ are given by

$$J^a(z)J^b(w) = \frac{kg^{ab}}{(z-w)^2} + \frac{f_c^{ab}J^c(w)}{z-w} + \dots \tag{1.1.31}$$

where k is a central value called the level and g^{ab} is the Cartan Killing form on the Lie algebra g. The field algebra generated by the fields $J^a(z)$ itself contains a stress energy tensor w.r.t. which the fields $J^a(z)$ are primary of conformal dimension 1. It reads

$$T(z) = \frac{1}{2(k+h)} g_{ab}(J^aJ^b)(z) \tag{1.1.32}$$

where h is the dual coxeter number of g. The stress energy tensor (1.1.32) is called the affine Sugawara stress energy tensor and one can show by explicit calculation that the central value associated to it is

$$c(g;k) = \frac{k\dim(g)}{k+h} \tag{1.1.33}$$

The modes of the fields J^a form an affine Lie algebra. The structure and representation theory of this algebra is well known [42, 27] and it will be shown in this thesis that many (perhaps all) W algebras are reductions of it.

The W algebras considered above all have in common that the modes of the fields that generate the W algebra span a Lie algebra, i.e. the commutator of two basis elements is again a linear combination of basis elements. This needn't always be true as was first realized by Zamolodchikov [62]. He consider the example of an algebra generated by $T(z)$ and a chiral primary field $W(z)$ of conformal dimension 3 with OPE

$$\begin{aligned} W(z)W(w) &= \frac{c/3}{(z-w)^6} + \frac{2T(w)}{(z-w)^4} + \frac{\partial T(w)}{(z-w)^3} + \\ &+ \frac{1}{(z-w)^2}\left(2\beta\Lambda(w) + \frac{3}{10}\partial^2 T(w)\right) + \\ &+ \frac{1}{z-w}\left(\beta\partial\Lambda(w) + \frac{1}{15}\partial^3 T(w)\right) + \dots \end{aligned} \tag{1.1.34}$$

where $\Lambda = (TT) - \frac{3}{10}\partial T$ and

$$\beta = \frac{16}{22+5c} \tag{1.1.35}$$

Since the right hand side of this OPE contains terms quadratic in T the commutator of the modes of $W(z)$ will involve quadratic terms. The W algebra generated by $T(z)$ and $W(z)$ is called W_3. Like the Virasoro algebra it has a discrete series of values for c for which there exist only a finite number of conformal dimensions that give rise to unitary representations (note that there is an additional label (or quantum number) corresponding to the W_0 eigenvalue). These representations have central value

$$c = c(m) = 2\left(1 - \frac{12}{m(m+1)}\right) \tag{1.1.36}$$

with $m = 3, 4, \ldots$. Note that the critical value $c = 1$ of the Virasoro algebra has been shifted to a higher value by the fact that more symmetry is present. The W_3 algebra has generalizations to arbitrary N, i.e. algebras whose generators have conformal weights $(2, 3, \ldots, N)$ called W_N. Like the Virasoro algebra and W_3 all these algebras admit a (principal) series of (W) minimal models. The central charges associated to these minimal models are (for W_N)

$$c(m) = (N-1)\left(1 - \frac{N(N+1)}{m(m+1)}\right) \tag{1.1.37}$$

Again there exists only finitely many allowed conformal dimensions for each c-value such that the corresponding representation is unitary.

In the applications of CFT to string theory and statistical mechanics it is necessary to consider CFT on higher genus Riemann surfaces. CFT on the cilinder can be transfered to the torus as follows: Define the torus by identifying the spatial S^1 at $t = 0$ with the spatial S^1 at $t = 2\pi\tau_0$. In doing so one can introduce a shift in x by $2\pi\tau_1$. Define then the partition function $Z = \mathrm{Tr}_{\mathcal{H}}(e^{2\pi i\tau_1 P_1} e^{2\pi i\tau_0 P_0})$ where as before P_0 and P_1 are the energy and momentum operators respectively. In the Euclidean theory we have to identify z with qz where $q = e^{2\pi i\tau}$ and $\tau = \tau_1 + i\tau_0$. The Torus partition function can then be shown to be

$$Z(q) = \mathrm{Tr}_{\mathcal{H}}(q^{L_0 - \frac{c}{24}} \bar{q}^{\bar{L}_0 - \frac{c}{24}}) \tag{1.1.38}$$

Inserting the decompostion (1.1.19) into Z one finds

$$Z(q) = \sum_{(a,\bar{a})} N_{a\bar{a}}\, \chi_a(q) \chi_{\bar{a}}(\bar{q}) \tag{1.1.39}$$

where $\chi_a(q)$ is the 'character' of the representation $\mathcal{H}_a$ of the chiral algebra $\mathcal{A}$

$$\chi_a(q) = \mathrm{Tr}_{\mathcal{H}_a}(q^{L_0 - \frac{c}{24}}) \tag{1.1.40}$$

In order for the CFT to make sense on the torus it should not depend on the way in which the torus is obtained from the cilinder, which means that Z should be invariant under the modular group $SL(2;\mathbf{Z})$ of the torus. This group is generated by the transformations

$$T : \tau \mapsto \tau + 1 \quad \text{and} \quad S : \tau \mapsto -\frac{1}{\tau} \tag{1.1.41}$$

and the fact that Z must be invariant under these transformations places strong constraints on the coefficients $N_{a\bar{a}}$. The study of these constraints is called the 'modular bootstrap' and has, in case the chiral algebra is the Virasoro algebra with $c < 1$ lead to a complete classification of modular invariant partition functions [1]. For $c \geq 1$ modular invariance requires any CFT to contain an infinite number of representations of the Virasoro algebra. As we have seen before W algebras are a possible solution to this difficulty since even though the decomposition may by infinite w.r.t. the Virasoro algebra it can still be finite w.r.t. an extension of this algebra. However for general W algebras the modular program is more complicated and has not been completed sofar.

The construction of W algebras has proceeded roughly along three independent lines. Many explicit results were obtained by the so called 'direct method' which boils down to solving the Jacobi identies (for a review see [63]). This method however has the drawback that it does not yield any information on the representation theory or geometrical interpretation of W algebras. It does give explicit results on the algebraic structure however (i.e. structure constants).

Another approach to W algebras is the so called coset construction [13, 89]. In this approach one constructs the generators and representations of a W algebra in terms of the representation theory of an affine Lie algebra. The generators of the W algebra become composite fields in terms of the affine currents (or in a more mathematical language, one embeds the W algebra into the universal enveloping algebra of an affine Lie algebra). We have already seen an example of this in the case of the Virasoro algebra eq.(1.1.32). One can generalize the affine Sugawara construction as follows: Let g be a simple Lie algebra, $g' \subset g$ and let $\hat{g}, \hat{g}'$ be their affinizations. The central charge k' of $\hat{g}'$ will in general be equal to jk where k is the central charge of $\hat{g}$ and j is the index of the embedding $g' \hookrightarrow g$. In the coset construction one then starts with the affine Sugawara tensors $T(z)$ and $T'(z)$ associated to $\hat{g}$ and $\hat{g}'$ respectively and considers the field $\tilde{T} = T - T'$ which remarkably generates a Virasoro algebra of central charge $c(g,k) - c(g',k,)$. Now, the point is that the Virasoro algebra $\tilde{T}$, called the coset algebra, commutes with the subalgebra $\hat{g}'$ which means that a highest weight module $L(\Lambda)$ of $\hat{g}$ decomposes into a direct sum of $Vir \oplus \hat{g}'$ representations. Obviously the coset space obtained from $L(\Lambda)$ by identifying states in the same $\hat{g}'$ highest weight module is a (not necessarily irreducible) highest weight module of the coset algebra. Now, for certain g' it may happen that the commutant of $\hat{g}'$ in the universal enveloping algebra of $\hat{g}$ contains

primary fields of higher conformal dimension generating a W algebra. In that case we can obtain representations of this W algebra in the same way as we discussed above for the Virasoro algebra.

The coset construction has some drawbacks. For example, it is not clear that the coset algebra (i.e. the commutant of $\hat{g}'$) is finitely generated. In fact this is in general not the case except, possibly, on some very specific $\hat{g}$ modules. In the case of the algebra W_3 this was shown in [89]. This means that the coset construction does not work as a construction between abstract algebras but only on a subset of representations which is rather unappealing. Another drawback is that the set of W algebras that one has been able to treat in this way is limited and that it does not seem to lead to a general classification scheme. Furthermore it does not provide any new insights into the geometrical background of W algebras. On the other hand the coset construction does yield an explicit construction of unitary representations of W algebras and, as will be shown in the next chapter, it can be realized at the Lagrangian level by the theory of gauged WZW models.

The most powerful method for constructing W algebras, and the one that has lead to important progress in trying to understand them, is the so called Drinfeld-Sokolov (DS) reduction procedure [66, 7, 67]. This procedure is the subject of this thesis and we shall therefore only describe it very briefly here. Drinfeld-Sokolov reduction is in fact (classically) simply Poisson reduction in infinitely many dimensions where as Poisson structure one takes the Kirillov Poisson structure associated naturally to any Lie algebra. Quantization of this reduction procedure is performed by BRST methods [19, 20, 97]. The DS approach has lead to many insights into the structure and meaning of W algebras as well as to the construction of field theories with explicit W symmetry. Also DS reduction yields a functor (unlike the coset construction where only a small number of affine Lie algebra representations are allowed) from the category of representations of the affine Lie algebra to the category of W representations [20]. In particular it is possible to obtain free field realizations of arbitrary W algebras in this way [19, 20, 97]. Furthermore one can obtain a great deal of information on the theories with W symmetry (for example correlation functions etc.) using the fact that they are a reduction from a theory with affine symmetry.

This concludes our introduction into CFT and W algebras. To provide some additional motivation we shall give a quick overview of the different areas of theoretical physics where W algebras and CFT are of interest. At the end of this first and introductory chapter we briefly discuss the contents of the remaining chapters.

1.2 String theories of elementary particle physics

String theory is an attempt to unify all known fundamental forces and particles into a single consistent mathematical framework. At the microscopic level (at distances down to 10^{-15} cm) and for processes involving strong and electroweak interactions the quantum field theory called the 'standard model' (SM) is very successful. With it we have uncovered at least part of the truth. On the other hand general relativity (GR) has been very successful in describing the large scale behaviour of the universe, in particular all experimentally accessible phenomena involving gravitational interactions; from suble effects in the solar system such as the perihelion precession of Mercury and the bending of light by the sun, to the cosmological aspects of the universe as a whole. The interpretation that the SM and GR provide of the fundamental forces they describe are however completely different. GR describes gravity purely in geometrical terms as the curvature of spacetime, while in the SM forces are seen as the exchange of particles (gauge bosons). Nevertheless, lead by past experience, it is conjectured by most physicists that GR and the SM are really two 'low energy limits' of the same fundamental theory and in order to construct this theory it has been tried very hard to quantize GR. Unfortunately quantum gravity is a nonrenormalizable quantum field theory essentially due to the fact that the gravitational coupling constant (Newton's constant) is not dimensionless. Therefore it suffers incurable infinities which render the theory without predictive power..

String theory [29, 36, 32, 30] is a new approach to the unification of the SM and GR. The revolutionary aspect of string theory is that its basic assumption is that matter is made up of very tiny one dimensional objects (strings). While a particle looks like a point at 10^{-13} cm, if we magnify it to scales resolving 10^{-33} cm then we may see that it is in fact an extended object. Unification of the complete particle spectrum is achieved in string theory by interpreting different vibrational modes of a string as the different fundamental particles. In particular the known particles are believed to correspond to the massless sector of a (super)string theory (the particle masses are assumed to be spontaneously or dynamically generated by breaking of (super)symmetry). There is no a priori reason to assume that particles are really one dimensional objects, however it turns out that (if spacetime is flat) the string spectrum contains a particle like state which transforms as a symmetric rank 2 traceless tensor under Lorentz transformations of spacetime, which is massless and transverse. This state can be identified with the graviton. Calculating the scattering amplitudes involving external graviton states in the low energy limit (using the string prescription) one finds that they agree with the amplitudes of gravitons evaluated from the Einstein-Hilbert action. Thus quantum Einstein gravity appears as an effective low energy theory and questions like unitarity and renormalizability must be addressed not at the level of Einstein theory, but at the string level.

Now, string theory is expected to have improved ultraviolet behaviour to all order compared to point particle physics due to the inherent fuzzyness of the string. In point particle physics divergencies arise because when two vertices of a diagram get arbitrarily close the propagator connecting these vertices blows up. In string theory the interaction between strings is pictured as the joining and splitting of these strings. The difference with point particle physics is that, because of the fact that strings are extended objects, there is no Lorentz invariant notion of a spacetime point at which the joining or splitting occured. This means that the interaction vertex has become inherently fuzzy and there is no analogue of the region where different vertices come arbitrarily close. If one compares string theory with the point particle theory of gravity on the quantum level, one learns that string theory has a built in ultraviolet regulator provided by the tower of massive modes of the string. These beautifully conspire to produce a finite result for the generic higher order quantum corrections. This is certainly one of the main achievements of string theory so far. Also another plague that appears to kill most particle theories, known as anomalies, are remedied in string theory.

Now, it is important to note that since any part of a string diagram describing the joining and splitting of two strings locally looks like the propagation of a free string (again because of the fuzzyness of the string diagrams) one finds that once one has decided upon a set of rules for free string propagation this fixes the complete theory. In point particle physics one still has the choice of certain factors to associate to the vertices of a Feynman diagram.

At the present state of development the full underlying theory of strings, sometimes also called string field theory, is not known (there has been progress in this area however, see for example [55]). It *is* possible to develop a perturbative approach to string theory by expanding around a 'classical solution' of the theory. This means that in some way an indefinite number of gravitons have condensed into a classical groundstate described by a metric $\{G_{\mu\nu}\}_{\mu\nu=1}^{D}$, where D is the dimension of spacetime. The coupling of the string to this nontrivial background is described by a 2 dimensional σ-model. This can be seen by, in analogy with the action of a point particle, taking the action of the string to be proportional to the area of the worldsheet Σ it sweeps out in spacetime

$$S[X] = T \int_{\Sigma} d^2\sigma \, \sqrt{\det(G_{\mu\nu}(X)\partial_a X^\mu \partial_b X^\nu)} \tag{1.2.42}$$

where the Roman indices $a, b, \ldots$ run over the 'world sheet' coordinates $\sigma \equiv (\sigma^0, \sigma^1)$ and $X^\mu(\sigma)$ describes the position of the string. The constant of proportionality T is called the 'string tension' because it can be interpreted as the energy per unit length of the string [29].

Obviouly the action (1.2.42) is very difficult to work with due to the square root in the integrand. It is possible to circumvent this difficulty by considering the theory

(1.2.42) as a subsector of an enlarged theory. For this introduce extra degrees of freedom in the form of a world sheet metric $h_{ab}(\sigma)$ and consider the action

$$S_M[X;h] = \frac{T}{2}\int_\Sigma d^2\sigma\sqrt{h}h^{ab}\partial_a X^\mu \partial_b X^\nu G_{\mu\nu}(X) \tag{1.2.43}$$

which reduces to (1.2.42) after insertion of the equation of motion for h_{ab} into (1.2.43).

At this point we would like to make a remark on why one considers strings and not 2 or 3 dimensional objects (in principle 3 dimensional objects would be the most appealing intuitively). It turns out that it is easy to write down a general action for such objects, just as we did for strings, but that the resulting quantum theory is nonrenormalizable by power counting. Also such theories do not have Weyl symmetry which is special to strings. There has been some activity however in this area and maybe some day it will turn out to be possible after all to write down a renormalizable theory for such higher dimensional objects.

The theory of free strings discussed above is to be considered as a first quantized theory, which means that interactions will have to be put in by hand. The interaction between strings is, in analogy with point particle physics, considered to be the joining and splitting of strings. An N string scattering process will perturbatively be represented by a sum over all Riemann surfaces connecting the N external strings. The g^{th} order term in this perturbation series corresponds to a compact genus g Riemann surface with N punctures (the N tubes attached to the surface extending into the far past and future, depending on whether the string is incoming or outgoing, can be 'scaled down' by Weyl rescalings of the world sheet metric to punctures on a compact surface). The partition function, expressing the vacuum to vacuum amplitude, will obviously involve a sum over all topologies of compact Riemann sufaces (without punctures). Since compact and orientable Riemann surfaces are classified by their genus the sum over topologies can be implemented by a sum over genera. In Minkowski space and after Wick rotating both in spacetime and on the worldsheet the partition function becomes

$$Z = \sum_{g=0}^{\infty}\int [DX][Dh]\,\exp\left(-\int_{\Sigma_g} d^2\sigma\sqrt{h}h^{ab}\partial_a X^\mu\partial_b X_\mu\right) \tag{1.2.44}$$

where Σ_g is a compact genus g Riemann surface. Note that this approach is necessarily perturbative.

Remember that the action (1.2.43) is invariant under arbitrary diffeomorphisms and Weyl rescalings. The measure $[DX]$ in (1.2.44) is defined in the usual way by the metric

$$\|\delta X\|_h^2 = \int d^2\sigma\ \sqrt{h}\ \delta X\cdot\delta X \tag{1.2.45}$$

which obviously depends explicitly on the world sheet metric. It can now be shown that DX is invariant under diffeomorphisms but not under Weyl rescalings. Denoting the explicit dependence of DX on h by D_hX we find that [16]

$$[D_{e^{\phi}h}X] = e^{\frac{D}{48\pi}S_L[\phi;h]}[D_hX] \tag{1.2.46}$$

where

$$S_L[\phi;h] = \int d^2\sigma\ \sqrt{h}\left(\frac{1}{2}h^{ab}\partial_a\phi\partial_b\phi + R\phi + \mu e^{\phi}\right) \tag{1.2.47}$$

Here R is the 'world sheet curvature scalar' calculated using the metric h. From this follows that the Weyl invariance of (1.2.43) is anomalous.

The nonanomalous gauge invariance present in the partition function is the group of diffeomorphisms. The space of metrics on a genus g surface Σ_g is isomorphic to $Diff(\Sigma_g) \times Weyl(\Sigma_g) \times \mathcal{M}_g$, where $\mathcal{M}_g$ is the moduli space of metrics on Σ_g. The integration over all metrics in (1.2.44) can therefore roughly be written as an integration over the gauge orbit, an integration over the Weyl orbit and an integration over modulispace (of course there will also be a Jacobian J related to the change of variables). In order to apply the Faddeev-Popov procedure to extract the gauge volume one has to make a choice of gauge. Let τ denote a set of moduli (i.e. parameters on $\mathcal{M}_g$) and let $\hat{h}(\tau)$ be a representative of the equivalence class of metrics determined by τ (remember that an element τ of $\mathcal{M}_g$ is an equivalence class of metrics on Σ_g). Take as gauge slice the set of metrics of the form $e^{\phi(\sigma)}\hat{h}(\tau)$. It is important to note that a subgroup of $Diff(\Sigma_g)$, the subgroup of conformal diffeomorphisms $Diff_c(\Sigma_g)$, slides along this gauge slice. There will therefore be a residual Virasoro (gauge) symmetry left in the final answer.

Since the moduli space of metrics on a genus g Riemann surface is finite dimensional the integration over $\mathcal{M}_g$ is an ordinary integration over moduli τ rather than a functional integration. As usual one represents the Jacobian J as a functional integral over ghosts b, c

$$J = \int [D_hb\, D_hc]\ e^{-S_{gh}[b,c;h]} \tag{1.2.48}$$

The ghost action can be shown to be invariant under diffeomorphisms and Weyl rescalings but like $[D_hX]$ the ghost measure suffers a Weyl anomaly [16]

$$[D_{e^{\phi}h}b][D_{e^{\phi}h}c] = e^{-\frac{26}{48\pi}S_L[\phi;h]}[D_hb][D_hc] \tag{1.2.49}$$

Now, inserting all these results into (1.2.44) and extracting the gauge volume we find that the partition function reads

$$Z = \sum_{g=0}^{\infty}\int_{\mathcal{M}_g} d\tau \int [D_{\hat{h}}X][D_{\hat{h}}\phi][D_{\hat{h}}b\, D_{\hat{h}}c]\ e^{-S_M[X;\hat{h}]-S_{gh}[b,c;\hat{h}]-\frac{26-D}{48\pi}S_L[\phi;\hat{h}]} \tag{1.2.50}$$

From this expression follows that the worldsheet metric decouples in $D = 26$ whereas it becomes popagating in $D \neq 26$ (in particular $D = 4$). So even though Einstein gravity is trivial in $D = 2$, due to the fact that the Einstein-Hilbert action is proportional to the Euler characteristic, a nontrivial 2 dimensional (non Einsteinian) theory of gravity can be induced at the quantum level. This 'induced gravity' has turned out to be an extremely interesting but complicated subject. In order to see why consider the genus 0 term of (1.2.50). The moduli space $\mathcal{M}_0$ is a point and we can take $\hat{h}_{ab} = \delta_{ab}$. Now performing the integral over X one finds

$$\int [D\phi]\, e^{\frac{D-26}{48\pi}\int d^2\sigma\, (\frac{1}{2}(\partial_a\phi)^2+\mu^2 e^{\phi})} \tag{1.2.51}$$

This functional integral is known to be difficult due to the e^{ϕ} term in the Liouville action. For higher genus the situation is even further complicated by the integral over moduli space.

Note that eq.(1.2.51) was obtained after gauge fixing which means that the action

$$S[\phi] = \frac{26-D}{48\pi}\int d^2\sigma(\frac{1}{2}(\partial_a\phi)^2 + \mu^2 e^{\phi}) \tag{1.2.52}$$

is a gauge fixed version of some other (generally covariant) action. One can find this other action by performing the matter integral $[DX]$ before one gauge fixes, i.e. in the genus 0 part of formula (1.2.44). What one finds is the following nonlocal (induced) action for the world sheet metric h:

$$S[h] = \frac{D}{48\pi}\int d^2\sigma \left(\sqrt{h}\, R\frac{1}{\sqrt{h}\Box}\sqrt{h}R\right) \tag{1.2.53}$$

where R is the 2 dimensional curvature scalar associated to the metric h. In the conformal gauge ($h_{ab} \sim \delta_{ab}$) this action indeed reduces to (1.2.52).

In the critical dimension $D = 26$ the integral (1.2.50) simplifies considerably but remains highly nontrivial. The calculations above were made under the assumption that spacetime is flat Minkowski space and one might wonder whether they remain valid when spacetime is some nontrivial 26 dimensional manifold with metric $G_{\mu\nu}$ instead of Minkowski space. Again choosing a diffeomorphism invariant regulator one finds that for general $G_{\mu\nu}$ Weyl invariance is broken as a consequence of which the theory becomes unacceptable due to the emergence of unphysical modes. We conclude that we have to insist in this case on Weyl invariance which will obviously restrict the possible choices for the spacetime metric $G_{\mu\nu}$. For this note that $G_{\mu\nu}$, describing the graviton condensate, acts in (1.2.43) as a set of coupling constants (one for each index pair (μ, ν) and spacetime

point X) and for Weyl invariance to be conserved on the quantum level the β-function of these coupling constants must be zero. An explicit computation yields [29]

$$\beta_{\mu\nu} = -\frac{1}{4\pi}\left(R_{\mu\nu} + \frac{1}{4\pi T}R_{\mu\rho\kappa\lambda}R_{\nu}^{\mu\kappa\lambda} + \ldots\right) \tag{1.2.54}$$

Note that in the limit $T \to \infty$ (which corresponds to an infinite string tension and therefore a pointlike string) the vanishing of the β-function coincides with the Einstein equations in vacuum. The other terms are string theoretical corrections to Einstein gravity. This spectacular result shows that Einstein relativity and the equivalence principle may arize as consistency conditions on the underlying 2 dimensional world sheet theory. However, at this point it is not clear which solution of the Einstein equations is selected by string theory or to put it differently, due to the lack of a satisfactory string field theory the structure of spacetime does not emerge dynamically from the theory but has to be put in by hand as a particular background. The search for the fundamental theory underlying string theory is therefore equivalent to the search for a background independent formulation of string theory.

We have seen above that 2 dimensional local scale invariance plays an important role in string theory. It can easily be shown that any 2 dimensional quantum field theory with local scale invariance has a traceless stress energy tensor. On the other hand it is well known that a quantum field theory is invariant under conformal transformations if and only if its stress energy tensor is traceless. We conclude that the string groundstates are described by conformal field theories which are associated to certain sigma models. Having arrived at this conclusion one now turns the argument around and interprets any CFT as a perturbative string groundstate around which one can start a perturbation theory essentially in the same way as we did above. From this point of view *a classification of all* ($c = 26$) *conformal field theories would correspond to a classification of all possible perturbative string vacua* and as we have seen W algebras play an important role in this classification. However, since conformal field theories correspond to groundstates, or put differently, to 'classical solutions' of an underlying theory, it is probably not the list of all CFTs that is of fundamental importance to string theory but more the basic principles that lead to such a classification. It is hoped that a proper formulation of CFT probably together with a general scheme as to their construction might lead to some clues regarding the physical and geometrical principles underlying string theory.

The need for a formulation of string theory that goes beyond a perturbative expansion around a given groundstate, i.e. a background independent formulation, has become an urgent one recently. Namely, in [28] it was shown that the topological expansion of the partition function diverges with the factorial of the genus $Z \simeq \sum_{g=0}^{\infty}(2g)!$

and is therefore not Borel summable (remember that even if a series

$$A(\alpha) = \sum_{g=0}^{\infty} \alpha^g f(g) \tag{1.2.55}$$

formally diverges its Borel transform

$$A_B(\alpha) = \sum_{\alpha=0}^{\infty} \alpha^g \frac{f(g)}{g!} \tag{1.2.56}$$

may still converge. The original function can be retrieved by taking the inverse transformation

$$A(\alpha) = \int_0^{\infty} A_B(\alpha t) e^{-t}\, dt \tag{1.2.57}$$

However if $f(g) \simeq g!$ then even the Borel transformed function A_B is singular). This means that the string perturbation theory around a conformal field theory is at best an asymptotic expansion. Usually, in gauge theories, such behaviour is ascribed to nonperturbative effects which makes it all the more necessary to develop the nonperturbative approach to string theory.

In the formulation of string theory discussed above the fields representing the coordinates of the string define a matter CFT described by a nonlinear sigma model. Generalizing the theory (1.2.50) by taking an arbitrary CFT instead of a sigma model we are lead to study CFTs coupled to 2-dimensional quantum induced gravity (i.e. CFT on a randomly fluctuating surface). In recent years our understanding of minimal ($c < 1$) models coupled to 2 dimensional gravity has improved a great deal. A number of results (scaling and critical dimensions) were obtained by using the fact that 2 dimensional induced gravity formulated in the light cone gauge possesses a hidden affine $SL_2(\mathbf{R})$ symmetry (we will come back to this in the next chapter) [50]. It has also been tried to use the fact that Liouville theory (1.2.52) is completely integrable as a classical field theory to obtain results in the quantum case [46]. These efforts have become known under the name 'the continuum approach'.

It has turned out that 2 dimensional gravity can also very conveniently be studied in terms of discretized world sheets. In this approach the integration over metrics is replaced by a sum over triangulations [65]

$$\sum_g \int [Dh] \rightarrow \sum_{\text{random triangulations}} \tag{1.2.58}$$

The reason that one can get exact results from this approach is that the sum over triangulations can be represented by an integration over (hermitean) $N \times N$ matrices. These

(matrix) models can be solved exactly and in the 'double scaling limit' (which involves taking $N \to \infty$) shown to be equivalent to string theory at $c \leq 1$. An attractive feature of this approach, especially in the light of the non-Borel summability of perturbation theory, is that it provides nonperturbative information on string theory. A remarkable fact that was recently discovered is that the partition function Z of a matrix model can be characterized by so called W-constraints which are differential equations of the form

$$W_n^{(s)} \cdot Z = 0 \quad , \mathrm{n} > \Delta_s \tag{1.2.59}$$

where $W_n^{(s)}$ are certain differential operators [59] that satisfy a W-algebra with conformal weights $\{\Delta_s\}$ (which W algebra depends on the matrix model). Conversely it is conjectured that there is a matrix model associated to any W algebra. Furthermore, out of the W constraints came the realization that the partition function Z can be viewed as a τ function of a certain integrable hierarchy of KdV type [59].

Finally we mention another approach to 2 dimensional gravity is topological field theory (TFT). A TFT is a quantum field theory in which all correlation functions are independent of the metric [56] which means that the independence of the world sheet metric is built into the theory from the beginning. In the literature a large number of TFT's have been constructed each of which seems to be equivalent to $c < 1$ matter coupled to 2D gravity. As one could have hoped the discrete and topological approaches to string theory have turned out to be equivalent [45]. As with matrix models a strong relationship with the theory of integrable hierarchies has been discovered and in fact it has been possible, using this relation, to calculate intersection numbers on the moduli space of Riemann surfaces for arbitrary genus [41] which is an important mathematical result.

From the string theoretical point of view the results discussed above are unphysical due to the restriction $c \leq 1$. Unfortunatly the methods used to obtain these results are not applicable to the so called 'strong coupling regime' $1 < c < 25$ due to the appearance of tachyons rendering the whole theory ill defined. Also note that, as we have seen in the previous section, for $c < 1$ there are only a finite number of unitary lowest energy Virasoro representations. Since Virasoro symmetry is a residue of the gauge symmetry present in string theory it follows, that all Virasoro descendent states are gauge equivalent to their highest weight states and that the theory essentially has a finite number of degrees of freedom. This simplifies matters drastically. For $c > 1$ however there are always infinitely many degrees of freedom due to the fact that any CFT with $c > 1$ contains infinitely many Virasoro representations. Here again W algebras may prove a way out since CFTs with W symmetry may be finitely reducible even for $c > 1$. This means that the critical central charge $c = 1$ can be shifted to a higher value by considering a W generalization of ordinary 2 dimensional quantum gravity (W gravity). One is here of course faced with the completely new concept of a

gauge theory based on a nonlinear gauge symmetry. Some progress in this area has been made however [40, 31, 9]. Note that, like the Virasoro algebra (which arose as a residue of 2 dimensional diffeomorphism invariance), W symmetry is here a residual (gauge) symmetry of something else. What this larger symmetry principle is that reduces on partial gauge fixing to (chiral) W symmetry is not known at this time.

In this section we have discussed string theory as on of the most promising candidates for a theory of quantum gravity. We have seen that W algebras may play an important role in this theory at different levels: first of all in the classification of the perturbative vacua, and second in trying to generalize recent results on $c < 1$ string theories to the more realistic 'strong coupling regime'.

1.3 The Quantum Hall effect

The integer quantum Hall effect (IQHE) was discovered in 1980 by von Klitzing, Dorda and Pepper (von Klitzing was awarded a Nobel prize for this discovery in 1985). Since then there has been a large effort to develop a theoretical understanding of it (see for example [52]) and remarkable progress has been make. The Hall effect has drawn the interest of high energy theorists due to certain indications that the description of the quantum Hall system is intimately related to conformal field theory. In this section we shall briefly indicate how this relation comes about.

Consider some (effectively) 2 dimensional system of electrons in a strong perpendicular magnetic field and at low temperatures. The IQHE is the phenomenon that under certain circumstances the conductivity tensor σ is of the form [53]

$$\sigma = \begin{pmatrix} 0 & -\frac{\nu e^2}{h} \\ \frac{\nu e^2}{h} & 0 \end{pmatrix} \tag{1.3.60}$$

where h is Planck's constant and $\nu = 0, 1, 2, \ldots$, which means that the longitudinal or 'diagonal' resistance R_L is zero while the transverse or *Hall* resistance R_H is quantized. We see that in such a state the system is dissipationless which means that the QHE is closely related to superconductivity and superfluidity.

Note that above we did not mention what material we were actually dealing with. Acually one of the remarkable properties of the QHE is its universality or material independence. This indicates that quantities like the Hall conductance $\sigma_H = \frac{\nu e^2}{h}$, which characterize the quantum Hall state, only depend on large scale properties of the system and do not depend on the precise microscoping details. It is truly remarkable to see that such a directly measurable macroscopic quantity in a rather complicated and impure medium is exactly equal to an integer times an expression in terms of fundamental constants.

The standard picture of the quantum Hall state [52] is that of the 'incompressible ground state' and we shall now very briefly discuss its basic ideas. In what follows we always assume the electron-electron interactions to be extremely weak so that the N electron states can be built by filling up one electron states in accordance with the Pauli principle. Furthermore we take the sample to have the topology of a disk with radius R and denote it by Ω.

As was shown by Landau planar electrons in a perpendicular magnetic field can occupy a discrete set of highly degenerate so called Landau levels which are eigenspaces of the one particle Hamiltonian H. The degeneracy in the Landau levels is labelled by the angular momentum l of the energy eigenstates. The single particle angular momentum states with angular momentum l are peaked around a mean radius $r_l = l_B\sqrt{l}$, where l_B is the so called 'magnetic length' $l_B = \sqrt{hc/eB}$. of course r_l increases as the angular momentum increases and since the radius of the sample is finite there exists a maximal angular momentum $l = L$ above which there can exist no states. This means that in a finite sample the Landau levels are finitely degenerate. Let us denote this degeneracy by N_B and let n_B be the associated particle density. It is easy to derive an explicit expression for n_B and it reads $n_B = eB/hc$. Note that n_B is a linear function of the magnetic field strenth.

Now suppose all occupied Landau levels are completely filled, i.e. the particle density n is an integer multiple of n_B, then due to the lack of low lying conduction modes the diagonal conductivity vanishes. The semiclassical formula for the off diagonal element of the resistivity tensor is $\rho_H = B/nec$ and inserting $n = \nu n_B$ into this formula, where ν is some positive integer, we find $\rho_H = h/\nu e^2$ (also remember that the longitudinal resistivity, the diagonal element of the resistivity tensor, is zero). The conductivity tensor, which is the inverse of the resistivity tensor, then obtains the form (1.3.60). This is the standard picture of the IQHE and in what follows we consider the state in which the occupied Landau levels are all completely filled in a little more detail.

For simplicity consider the state $|\Omega\rangle$ where only the first Landau level is filled and all higher Landau levels are empty. Clearly such a state is incompressible since a compression would mean that at least one electron would have to go to a state with lower angular momentum. Since all states in the lowest Landau level with lower angular momentum are already occupied the electron would have to go to a lower angular momentum state in a higher Landau level. This process however faces an energy gap which means that the state is incompressible. It is possible to calculate the expectation value of the particle density operator ρ in the state $|\Omega\rangle$ and what one finds is that it is constant for radii $r << R$ and drops rapidly to zero in the neighborhood of the boundary of the sample. From these facts (i.e. the incompressibility and the uniform density) it follows that in the state $|\Omega\rangle$ the electrons behave like a rigid droplet of (quantum) fluid.

Another quantity of interest is the expectation value of the electron current operator

$J^i(x)$. Again a straighforward calculation yields

$$\langle\Omega|J^i(\mathbf{x})|\Omega\rangle \quad\sim\quad \varepsilon^{ij}\partial_j\langle\Omega|\rho(\mathbf{x})|\Omega\rangle \tag{1.3.61}$$

Obviously the current vanishes in the interior of the sample, where the density is constant, while it supports a transverse (chiral due to the magnetic field) current around the boundary of the droplet. This is the so called *edge current* and it indicates that all nontrivial dynamics of the system is concentrated around the boundary.

Recently there has been a great deal of activity on the relation between CFT and the QHE [35]. We shall now briefly discuss the basic ideas underlying this relation. As we mentioned above all nontrivial dynamics of a Hall system is restricted to the boundary. One might expect to be able to describe the system effectively as a 1+1 dimensional theory. In fact in [11] it was shown that the 2+1 dimensional Hamiltonian of the Hall sample reduces (in the scaling limit) to a 1+1 dimensional Hamiltonian for a system of excitations living on the boundary of the sample. Furthermore this theory turns out to be equivalent to a theory of relativistic chiral scalar fermions living on S^1 [23]. It is known that the spectrum of this theory can be described completely in terms of an affine $U(1)$ algebra [2] which, as we have seen in the first section, is a particular example of a W algebra.

In [23] the same result was derived in a slightly different way using classical arguments. Consider the current density $\vec{j}$ induced by an electric field $\vec{E}$, i.e. $j^i(\vec{x},t) = -\sigma_H\varepsilon^{ij}E_j(\vec{x},t)$, where $\vec{x}\in\Omega$ and $\sigma_H = \frac{\nu e^2}{h}$. Using Faraday's induction law we find $j^0 = -\sigma_H B$ and we can write these equations as follows

$$J_{\alpha\beta}(x) = \sigma_H F_{\alpha\beta}(x) \tag{1.3.62}$$

where $F_{0i} = E_i$, $F_{12} = -B$ and $J_{\alpha\beta} = \varepsilon_{\alpha\beta\gamma}j^\gamma$. In terms of differential forms this equation reads $J = \sigma_H F$ while the continuity equation for the current j and Faraday's law reduce to $dJ = 0$ and $dF = 0$ respectively (here d is the ordinary exterior differential). Now, since Ω is contractible we find (by the Poincare lemma) that $F = dA$ and $J = da$ where A is the 3 dimensional vector potential and a is a one form called the 'vector potential of the electric current'. In terms of these potentials eqn.(1.3.62) can be written as $d(a - \sigma_H A) = 0$ which can be derived from an action principle with action

$$S_{CS}[a-\sigma_H A] = \int_{\mathbf{R}\times\Omega}(a-\sigma_H A)\wedge d(a-\sigma_H A) \tag{1.3.63}$$

which we recognize as a $U(1)$ Chern-Simons gauge theory for the gauge field $a - \sigma_H A$. The Chern-Simons action is topological since it does not require a metric and it is shown

in [23] that this explains why σ_H does not depend on the microscopic disorder in the sample.

We conclude that the classical electrodynamical equations describing the quantum Hall sample are actually the equations of motion of a Chern-Simons action. This makes it possible to describe the large-scale properties of a Hall system in terms of a Chern-Simons gauge field theory [23]. From this picture the W algebra arises as follows: It is well known [60] that a Chern-Simons theory on a two dimensional manifold Ω is equivalent to a WZW model (or a chiral affine current algebra) on the boundary $\partial\Omega$ of Ω so again we find that, in the scaling limit, a quantum Hall system can be described by a chiral sector of a two dimensional conformal field theory.

Up to now we implicitly assumed that the magnetic field B is large enough to align all electron spins in the direction of B. In that case the spin wave function is totally symmetric while the orbital wave function is (as electrons are fermions) totally antisymmetric. In this approximation the electrons can be described as two dimensional scalar fermions. As we have seen this approximation leads to an affine $U(1)$ algebra. If the magnetic field is not large enough to align all the spins then one finds, due to the fact that the edge currents become spin carrying, that the CFT governing the edge theory has an underlying level 1 affine$SU(2)$ symmetry [23].

In the above we did not concern ourselves with the mechanism by which the bulk of the sample creates the incompressible groundstate. This would require taking into account the microscopic interactions between the electrons and probably most important of all a study of impurity effects, for it can be easily shown that there would be no QHE without impurities [52]. Rather we assumed that the incompressible groundstate is indeed formed. The theory of how this state is formed is still in its infancy.

1.4 Phase transitions in 2 dimensions

Phase transitions play an important role in many areas of theoretical physics. Roughly a phase transition is a sudden change in certain macroscopic properties of a many body system as a result of a slight change in some external parameters like the temperature. Mathematically a singularity in the free energy (or its derivatives) of a system is indicative for a phase transition. Here we shall be considering second order phase transitions in two dimensional lattice models like the Ising model.

As is well known a many body system (where we use the term 'body' in a very broad sense) is described in statistical mechanics (in the canonical formalism) by a map ρ from the phase space $\mathcal{M}$ of the system (i.e. the set of all microstates) to the interval $[0, 1]$ such that the integral (or sum) over phase space of ρ is equal to 1. Usually the map ρ

has the well known Boltzman form

$$\rho = \frac{1}{Z} e^{-\beta H} \tag{1.4.64}$$

where H is the Hamiltonian of the system and Z is the partition function

$$Z = \int_{\mathcal{M}} e^{-\beta H} \, d\mu \tag{1.4.65}$$

(here $d\mu$ is some measure on phase space). If Q is some measurable quantity (i.e. a function on phase space), then the average value of Q is given by

$$\langle Q \rangle = \int_{\mathcal{M}} Q \rho \, d\mu \tag{1.4.66}$$

As the system moves through phase space the observable quantities (observables) will fluctuate around their equilibrium values. In general these fluctuations will not be large and a measure for their size is given by the quantity

$$\langle \Delta Q \rangle^2 = \langle Q - \langle Q \rangle \rangle^2 \tag{1.4.67}$$

It can be shown [54] that the size of certain fluctuations and thermodynamical quantities like the specific heat are proportional. It is well known that near a phase transition many respons functions blow up which means that one expects to encounter abnormally large fluctuations near criticality. These large fluctuations are a reflection of the internal adjustments the system is making as it prepares to change to a new state.

In the case of lattice models it is possible to calculate the correlation functions between the dynamical variables on the vertices, edges or faces (depending on the model). For example, in the Ising model one can calculate the correlation function between two arbitrary spins s_i and s_j on the lattice. This correlation function, which is given by $\Gamma_{ij} = \langle (s_i - \langle s_i \rangle)(s_j - \langle s_j \rangle) \rangle$ clearly measures the correlation between fluctuations in the spins s_i and s_j. On a homogeneous lattice (like the square lattice) Γ_{ij} can only depend on the difference of the position vectors $\vec{r}_i, \vec{r}_j$ of the spins s_i, s_j. It is easy to show that Γ_{ij} is not equal to zero even if s_i and s_j are not nearest neighbours. Away from the critical point the correlation function Γ_{ij} decays exponentially in $r_{ij} = |\vec{r}_i - \vec{r}_j|$ while at criticality it has power law decay. Therefore at criticality fluctuations are correlated over large distances, or put differently, the correlation length becomes infinite. It is essentially this fact, i.e. the presence of fluctuations on all lenth scales, that made the problem of describing critical systems so hard. The crucial ingredient which made the

problem tractable is the property of scale invariance which is at the heart of the successfull renormalization group approach to critical phenomena. It means in particular that correlation functions tend to be scale invariant near criticality. Effectively one therefore expects to be able to describe a lattice system at criticality by a 2 dimensional quantum field theory. Indeed, in the scaling (or continuum) limit one interprets certain lattice correlation functions as Euclidean Green functions of some Euclidean field theory. This field theory will be scale invariant and can be explicitly constructed provided the lattice system admits a self-adjoint transfer matrix (it may be possible to do this even if there is no selfadjoint transfer matrix but it is not known how). Remember that a doubly periodic $N \times N$ lattice model is said to admit a transfer matrix if there exists a $2^N \times 2^N$ matrix T such that

$$Z = \mathrm{Tr}(T^N) \tag{1.4.68}$$

If a transfer matrix exists then obviously the problem of calculating the partition function reduces to finding the eigenvalues of the transfer matrix.

Selfadjointness of the transfermatrix is a frequent but not fundamental property from the point of view of statistical mechanics (for example it fails in the theory of self avoiding walks). It turns out however that if it is satisfied that one can use a procedure similar to Osterwalder-Schrader reconstruction [47] to associate to the set of Euclidean Green functions, obtained by taking the scaling limit of the lattice correlation functions a unitary relativistic quantum field theory [21]. First one constructs out of the set of Euclidean Green functions a separable Hilbert space $\mathcal{H}$ with a unique vacuum state Ω. Then it is shown that $\mathcal{H}$ carries a representation of the group of spacetime translations from which one can construct selfadjoint operators H and P satisfying the relativistic spectrum condition. Finally it is then shown that the thus obtained quantum field theory is scale invariant. Now, for homogeneous and isotropic systems scale invariance and the fact that interactions are local imply conformal invariance. We conclude that *conformal field theories can be interpreted as field theories describing statistical mechanical lattice models in the continuum limit and at criticality.* The classification of critical phenomena (or in the terminology of renormalization theory, all fixed points of the renormalization group) in 2 dimensions corresponds therefore (at least in part) to the classification of all 2D Euclidean conformal field theories. The central charge of the CFT is interpreted as a measure of the effective number of field degrees of freedom which have large scale fluctuations at criticality.

It has been possible to identify all minimal models with certain critical 2 dimensional lattice models. Before giving some examples let us note that to any CFT there is associated a whole universality class of statistical mechanical models so a given realization of a CFT as a lattice model is not by far unique.

The most famous identification is that of the $c = c(3) = \frac{1}{2}$ minimal model with the critical Ising model [8]. In fact it was shown in [85] that all minimal models correspond

to the critical points of exactly solvable restricted solid on solid (RSOS) models defined on Dynkin diagram A_n.

Remember that the minimal models all have $c < 1$. The value $c = 1$ corresponds to a free bosonic field, i.e. to a Gaussian fixed point. In this theory there is one parameter, which can be interpreted as the compactification radius of the free bosonic field, and the critical exponents depend continuously on this parameter. We are therefore dealing with a line of fixed points. This line can be shown to correspond to the critical line the 8 vertex model [4].

Statistical mechanical models with extended conformal symmetries (i.e. W symmetry) have also been found. In fact it turns out that the 3 states Potts model already mentioned above has W_3 symmetry at criticality. However, a systematic study of W symmetries in statistical mechanics has not been embarked on. This appears to be a worthwile problem for the future.

1.5 Solitary wave phenomena

Solitary waves were first discovered in 1834 by the Scottish engineer John Scott Russell when he observed a smooth and well rounded heap of water of approximately 10 meters length and 40 cm height that moved through a channel of water apparently without change of form or speed. Later, in an attempt to study this phenomenon, he successfully recreated many solitary waves in a laboratory setting. Partly due to scepticism and disbelief within the scientific community it was not until 1895 before any solid theoretical description of the phenomena observed by Russell was given. In that year D.J.Korteweg and G.de Vries derived their famous equation for one dimensional long, small amplitude water waves in a shallow channel. This equation, nowadays known as the KdV equation reads

$$u_t = u_{xxx} + 6uu_x \tag{1.5.69}$$

where $u = u(x,t)$ is the surface elevation above the equilibrium level at x at time t. This equation indeed has solutions of the type described by Russell as was shown by Boussinesq and Rayleigh

$$u(x,t) = s\kappa^2 \mathrm{sech}^2(\kappa(x + \kappa^2 t - x_0)) \tag{1.5.70}$$

where κ and x_0 are constants. A large part of the early scepticism towards Russell's discovery was that it was believed that a solitary wave could not be stable. With the discovery of the KdV equation it became clear that the usual dispersive effects of wave propagation cancel precisely against the nonlinear effects governed by the nonlinear term uu_x.

Solitary wave phenomena are not restricted to hydrodynamical systems however. In the 1960's E.Fermi, J.Pasta and S.Ulam accidentally ran into some solitary waves when they were investigating the heat conductivity of solids. The problem was this: if a solid is modelled by a one dimensional lattice of masses coupled by harmonic springs (i.e. springs for which the return force is linear in the displacement from the equilibrium) then energy is carried unhindered by the independent normal modes. This means that the effective thermal conductivity would be infinite or put differently that no thermal gradient is needed to push the heat through the lattice from one end to the other. Since this is certainly not the case Debye suggested that the finiteness of the thermal conductivity is caused by anharmonicity of the springs, i.e. nonlinearity in the return force. This suggestion motivated Fermi, Pasta and Ulam to undertake a numerical study of an anharmonic lattice. What they expected was that an initial state in which all energy was in the lowest mode would eventually relax into a state in which the energy would be distributed equally among all modes due to the nonlinearity of the spring couplings. The relaxation time would then be a measure of the diffusion coefficient. To their astonishment what happened is that the energy flowed back and forth among several low order modes eventually to be recollected almost entirely into the lowest mode. Then the whole process started all over again.

In an attempt to explain this phenomenon M.D.Kruskal and N.J.Zabusky again ran into the KdV equation. They also found numerically that the KdV equation has solutions in which there appear more than one solitary waves. Furthermore they found that if two solitary waves collided, then after some time they reappeared untouched, i.e. with the same height, width and speed. The only evidence of a collision was a phase shift whereby the solitary waves were not at the places they would have been had there not been a collision. This particle like behaviour lead to the name *solitons*. It was only after these discoveries by Kruskal and Zabusky that the theory of solitary waves became popular among mathematicians and physicists.

The next important step in the development of the theory of solitons was the realization that there are infinitely many quantities $\{H_i\}_{i=1}^{\infty}$ that are conserved under the KdV flow. This lead to the insight that the KdV equation was just part of an infinite dimensional completely integrable system. Remember that a finite dimensional classical mechanical system described by some Hamiltonian H_1 is called 'completely integrable' if there exist M functions $\{H_i\}_{i=1}^{M}$ of the coordinates and momenta (where M is the number of coordinates) that commute w.r.t. the Poisson bracket, i.e. $\{H_i, H_j\} = 0$. This means that all the quantities H_i are conserved w.r.t. the time evolution generated (through the Hamilton equations) by H_1 and furthermore that the conserved quantities Poisson commute among themselves. A theorem by Liouville then states that one can trivialize such a system by a canonical invertible coordinate transformation, i.e. there exist phase space coordinates Q_i and P_i (called action angle variables) such that Q_i is constant and P_i is a linear function of t. The transformation to the action angle

coordinates is called the action angle transform. Note that within the system $\{H_i\}$ the original Hamiltonian plays no prefered role which means that in fact we have obtained a set of N physical systems, described by the Hamiltonians $\{H_i\}$ which are completely integrable.

In infinitely many dimensions it is not so clear how to define complete integrability but it is clear that one should at least have infinitely many conserved quantities in involution. The point is now that the inverse scattering transformation used to solve many integrable hierarchies, can be interpreted as an infinite dimensional example of an action angle transform.

Let us now describe the KdV hierarchy, as well as a number of other hierarchies, as infinite dimensional integrable systems. For this consider the space D_N of differential operators L of the form

$$L = \partial^N + \sum_{i=0}^{N-2} u_i \partial^i \tag{1.5.71}$$

where u_i are smooth real valued functions of $x \in \mathbf{R}$ that converge quickly to 0 when $|x| \to \infty$. On this space the operation of taking the derivative is invertible so we can consider the space PD_N of so called 'pseudodifferential operators'

$$X = \sum_{i=-\infty}^{N-1} u_i \partial^i \tag{1.5.72}$$

From the fact that ∂ is a derivation it immediately follows that

$$\partial^{-1} u = \sum_{i=0}^{\infty} (-1)^i (\partial^i u) \partial^{-i-1} \tag{1.5.73}$$

One usually defines the algebra of pseudodifferential operators even when working with a space on which ∂ is not invertible. However in that case ∂^{-1} is a formal generator of the algebra satisfying the permutation relation (1.5.73). Before we define the hierarchies let us introduce a few more concepts that we shall need. Let X be a pseudodifferential operator, then we define the 'residue' of X to be the coefficient of ∂^{-1} and the 'trace' of X by the integral over the residue of X, i.e.

$$\mathrm{Tr}(X) = \int dx \, u_{-1}(x) \tag{1.5.74}$$

The space $F(D_N)$ is defined to be the space of functionals of the form

$$F[L] = \int dx \, \tilde{F}(u) \tag{1.5.75}$$

where $\tilde{F}$ is a function of $\{u_i\}$ and a finite number of its derivatives. Note that elements of $F(D_N)$ map differential operators to real numbers.

In the inverse scattering method a crucial role is played by the so called isospectral deformation problem. Consider the eigenvalue equation

$$L\psi = \lambda\psi \tag{1.5.76}$$

where $L \in D_N$. The question one asks is: suppose the functions u_i are time dependent, i.e. $u = u(x,t)$, what time evolution is such that the eigenvalues λ are time independent? In turns out (as was shown by Lax) that this is true iff the functions u_i satisfy a system of equations that can be written as

$$\frac{dL}{dt} = [M, L] \tag{1.5.77}$$

where M is again a differential operator (not necessarily of order N). The pair of operators (L, M) is called a Lax pair. Now, in order for the equation (1.5.77) to make sense M must have a very specific form. It can be shown [66] that M must be a linear combination of

$$M_k = (L^{k/N})_+ \tag{1.5.78}$$

where $L^{1/N}$ is the (unique) pseudodifferential operator such that $(L^{1/N})^N = L$ and the index + means that we only retain terms with positive powers of ∂. The hierarchy of evolution equations is now given by

$$\frac{dL}{dt_k} = [M_k, L] \tag{1.5.79}$$

It can proved by explicit calculation [66] that the different flows defined by (1.5.79) commute which means that it is legitimate to introduce an infinite number of times t_k and to consider u_i a function of all these times. The KdV hierarchy is obtained by taking $N = 2$ while $N = 3$ leads to the Boussinesq hierarchy.

We now describe the infinite set of conserved quantities and the (bi)Hamiltonian structures of the hierarchies constructed above. Let $F \in F(D_N)$ be a functional on D_N, then we define the pseudodifferential operator X_F by

$$X_F(\cdot) = \sum_{i=1}^{N-1} \partial^{-i} \frac{\delta F}{\delta_{u_{i-1}}} \cdot \tag{1.5.80}$$

On $F(D_N)$ there exist two coordinated Poisson brackets, called Gelfand-Dickii brackets given by

$$\begin{aligned} \{F,G\}_1(L) &= \mathrm{Tr}([L,X_F]_+X_G) \\ \{F,G\}_2(L) &= \mathrm{Tr}(\nabla_{X_F}(L)X_G) \end{aligned} \tag{1.5.81}$$

where

$$\nabla_{X_F}(L) = L(X_FL)_+ - (LX_F)_+L \tag{1.5.82}$$

Here 'coordinated' means that any linear combination of these two Poisson brackets is again a Poisson bracket. Now the point is that the Lax equation (1.5.79) is Hamiltonian w.r.t. both Poisson structures and the Hamiltonian functions are given by

$$H_k[L] = \mathrm{Tr}(L^{k/N}) \tag{1.5.83}$$

These quantities are conserved w.r.t. all flows in the hierarchy and commute among themselves w.r.t. both Gelfand-Dickii Poisson structures. We have therefore obtained an infinite dimensional integrable system.

Consider now the Poisson algebra $(F(D_N),\{.,.\}_2)$. Simply using the definitions it is easy to show that for $N=2$ the 'coordinate functional' u has the following Poisson bracket with itself

$$\{u(x),u(y)\}_2 = -\frac{1}{2}\delta'''(x-y) - 2u(y)\delta'(x-y) + u'(y)\delta(x-y) \tag{1.5.84}$$

which is the classical version of the Virasoro algebra. In fact it has been shown in [64] that in general the algebra $(F(D_N),\{.,.\}_2)$ is nothing but a classical version of the W_N algebra. In a similar way it can be shown that affine Lie algebras arise as the (second) Hamiltonian structures of certain matrix generalizations of the Lax equations. Thus we conclude that *(classical) W algebras arise in the theory of solitary waves as Hamiltonian structures*. Conversely one conjectures that to any classical W algebra there is associated a hierarchy of integrable evolution equations. If and how the special structure of W algebras is related to the integrability of the associated hierarchies is not known. Integrable hierarchies do however have another algebraic structure that is explicitly linked to the integrability, the so called 'weak action angle structure'. This is what we discuss next.

Remember that complete integrability implies the existence of so called action-angle variables. Let $\{f_n\}$ and $\{\phi_n\}$ denote the action and angle variables respectively, then their distinguishing property is that the action variables are constant w.r.t. all times while the angle variables depend linearly on all times. This implies that $\{f_n,f_m\}=0$ and $\{f_n,\phi_m\} = \omega_{nm}(\{f_k\})$. Transferring this algebraic structure to the Hamiltonian

vectorfields X_i, L_n associated to f_n and ϕ_n respectively we find that $[X_n, X_m] = 0$ while $[X_n, L_m] = C^k_{nm} X_k$, where $[.,.]$ denotes the usual commutator between vectorfiels. Note that there is a great deal of freedom in the choice for the angle variables since adding to them a function of action variables does not destroy their basic property, that of linear time dependence. It can be shown that this freedom is large enough to pick the variables ϕ_n (and therefore their vectorfields) such that $[L_n, L_m] = \tilde{C}^k_{nm} L_k$. The algebra spanned by $\{X_n\}$ and $\{L_n\}$ is called the 'weak action angle structure'.

Let us now return to the case of integrable hierarchies. Consider for simplicity hierarchies of equations in one field $u = u(x,t)$. The equations in the hierarchy are $u_{t_k} = X_k(u)$ where $\{X_k(u)\}_{k=0}^{\infty}$ are (finite) polynomials in $u, u_x, u_{xx}, \ldots$. These quantities can be seen as (the components of) vectorfields on an infinite dimensional Banach manifold and therefore we can define their commutator [48]. Assuming that the hierarchy is integrable we must have $[X_n, X_m](u) = 0$ for the different flows must commute.

A crucial role is played by the so called 'recursion operator', i.e. many hierarchies (including the KdV hierarchy as we shall see later) admit an operator Φ mapping elements of the hierarchy to higher elements, i.e. $X_{n+1} = \Phi X_n$. Such a recursion operator may even exist when the hierarchy doen not admit a Hamiltonian structure (see for example the Burgers hierarchy) and is therefore not completely integrable.

The next ingredient in the study of the weak action angle structure of integrable hierarchies is the so called 'scaling field'. The main idea is here that integrable hierarchies are usually invariant under scaling transformations of the type $u(x,t) \mapsto e^{\alpha\epsilon} u(e^{\beta\epsilon} x, e^{\gamma\epsilon} t)$ where α, β and γ are real numbers and ϵ is some parameter. For the KdV equation these numbers are $2, 1$ and 3 respectively. The scaling transformations are generated infinitesimally by vectorfields τ that have the property $[X_n, \tau] = \lambda X_n$. It is a remarkable fact that we can use these scaling fields to find the weak action angle structure of an integrable hierarchy. For this define $L_0 = \frac{1}{\lambda}\tau$ and $L_n = \Phi^n L_0$, where Φ is again the recursion operator. It can then be shown [48] that the vectorfields X_n and L_n form the algebra

$$\begin{aligned} [X_n, X_m] &= 0 \\ [L_n, X_m] &= (m+1) X_{n+m} \\ [L_n, L_m] &= (m-n) L_{n+m} \end{aligned} \tag{1.5.85}$$

We conclude that *the weak action angle structure of an integrable hierarchy is isomorphic to half an affine $U(1)$ algebra*. Obviously it is only *half* because the indices n and m run only over positive numbers. This also causes the absense of central terms in the relations (1.5.85).

Let us briefly give an example. The KdV hierarchy is the following set of evolution equations: $u_{t_k} = \Phi^k X_0$ where $X_0 = u_x$ and $\Phi = \partial^2 + 4u + 2u_x \partial^{-1}$. For $k = 1$ this leads to the KdV equation which is invariant under the scaling transformation with $\alpha = 2, \beta = 1$

and $\gamma = 3$. These scaling transformations are infinitesimally generated by the scaling field $\tau = xu_x + 2u = L_0$. From this it is easy to construct the complete weak action angle structure.

In this section we have clearly shown that W algebras play an important role in the theory of integrable hierarchies. We have shown that nonlinear W algebras appear as Hamiltonian structures while the action angle structure of the hierarchy is intimately related to an affine $U(1)$ algebra.

1.6 Outline of the thesis

In the previous sections we have seen that CFT and W algebras play a potentially important role in many areas of physics. Progress towards the application of W algebras in order to obtain physically significant results is however hampered by the fact that so little is known about them. Apart from some very special cases like W_3 little is known about the representation theory of nonlinear algebras and also it is not clear what their geometrical meaning is. Remember that the Virasoro algebra arises through unitary representations of the group of conformal diffeomorphisms which gives it a clear and well understood geometrical interpretation. It is natural to ask what the analogous interpretation is for a given W algebra. This has turned out to be a very difficult problem due to the nonlinear nature of a generic W algebra. It is only one of the many mysteries still surrounding the concept of nonlinear symmetries in 2 dimensions.

In this thesis we are going to present new steps towards the understanding of W algebras. Since most of the material presented in chapters 3 4 and 5 will be rather technical and algebraic we are going to ease into this 'mathematical state of mind' by providing in chapter 2 an intuitive motivation for the constructions presented in the following chapters. In chapter 2 we consider how Liouville theory arises as a reduction of a WZW model and also how this translates into a reduction of the WZW Noether current algebra. For this it will be necessary to use a piece of mathematical machinery called 'Poisson reduction' and since this is not common knowledge among physicists we briefly outline its main ideas. A slightly different, but equivalent, construction of Liouville theory through 'gauged WZW models' is also discussed. The natural framework for the quantization of Poisson reductions is the BRST formalism which will also be considered.

In chapter 3 we then start the main analysis of W theory. One of the reasons that this has been so hard is that, unlike the theory of affine Lie algebras, the underlying finite structures (or even the existence of these) were not known. The theory of affine Lie algebras would have much harder to develop if finite dimensional simple Lie algebras were not so well understood. It is with this in mind that we introduce and develop in chapter 3 the theory of 'finite W algebras'. Classically these are nonlinear but finitely

generated Poisson algebras that can be obtained from simple Lie algebras by Poisson reduction. The quantization of these algebras, consisting of an assignment of an appropriate noncommutative 'operator' algebra to a given classical finite W algebra, will be performed by BRST methods. We also develop the classical and quantum representation theory of finite W algebras. It turns out that apart from ordinary finite dimensional representations finite W algebras also admit Fock representations in terms of harmonic oscillators. We also calculate the symplectic leaves, or what could be called the W coadjoint orbits, of arbitary finite W algebras.

Having developed the theory of finite W algebras in detail we turn to the study of their infinite dimensional counterparts. For this we generalize the so called Drinfeld-Sokolov reduction procedure in order to obtain a very large class of W algebras. In fact it is a conjecture that essentially all W algebras can be obtained by (possibly repeated) use of this procedure. The classical theory will lead to an elegant new formula for the Poisson structure of arbitrary W algebras which is, in a very special case, equivalent to the Gelfand-Dickii Poisson brackets discussed in section 5 of the introduction. Again the quantization is carried out by BRST methods and an explicit algorithm is developed in order to construct the W generators and their relations. Furthermore it is shown that any W algebra can be embedded into a semisimple affine Lie algebra and also that all W algebras admit a free field realization. All through chapter 4 we purposely follow the same lines as we did in chapter 3 in order to illustrate the formal similarity between the theories of finite and infinite W algebras.

In the fifth and final chapter we consider some applications of W theory to integrable models. First it is shown that finite dimensional completely integrable Toda theories are reductions of a system describing free particles moving on group manifolds. Furthermore using results obtained in chapter 3 the set of Toda theories is extended greatly and it is shown that these models have finite W symmetry. Next we consider the infinite dimensional version of this reduction procedure. This will lead to generalizations of the construction discussed in chapter 2 of Liouville theory as a reduction of a WZW model. The models thus obtained are generalized Toda theories in which the W algebras of chapter 4 arise as Noether symmetry algebras. Classical W algebras can also arise as Hamiltonian structures of integrable hierarchies however as we have seen in this introduction. Therefore we investigate what hierarchies are associated to the the aforementioned W algebras in the last section of the final chapter..

The results of this thesis shed a new light on the theory of W algebras since it becomes clear that they can be viewed as reductions of Lie algebras. We believe that this point of view is very promising and may lead to many interesting new developments both in mathematics and physics.

Chapter 2

Classical and quantum reduction of WZW models

In classical mechanics it is well known that some highly nontrivial systems can be described as reductions of simpler ones. The general scheme behind this is reduction of a mechanical system with symmetry: suppose the system has a certain symmetry then the number of 'essential' degrees of freedom can be reduced. However, the equations of motion in terms of the smaller number of degrees of freedom will in general look much more complicated and be harder to investigate then the original equations of motion. One could now turn this around and ask whether a given complicated system can be described as a simpler system with more degrees of freedom. In finite dimensions systematic use was made of this idea to integrate explicitly many more or less complicated systems (for a review see [49]).

In field theory a similar construction was used recently in the study of Toda field theories in general and Liouville theory in particular. As we will show in this chapter Liouville theory is in fact nothing but a reduction of an $SL_2(\mathbf{R})$ Wess-Zumino-Witten (WZW) model which is well studied and understood. Again, in the WZW variables the Liouville eqn becomes extremely simple and the well known Leznov-Savaliev solutions arise simply as reductions of the WZW solutions. As one could have expected the WZW model has much more symmetry than the Liouville model and as we will show the WZW-Liouville reduction also manifests itself on the level of their Noether symmetry algebras. This will lead us to the concept of Poisson reduction and, in the quantum case, BRST quantization.

2.1 Constrained WZW models

The progress made towards solving 2 dimensional induced quantum gravity using the so called 'continuum approach' is for a large part due to the observation by Polyakov that the induced gravity action possesses in the chiral (or light cone) gauge a hidden affine $SL_2(\mathbf{R})$ symmetry [50]. Motivated by the insight that such an underlying symmetry has to manifest itself in one form or another in all gauges Forgac et al [22] showed that Liouville theory

$$S[\phi] = \int d^2x \, (\frac{1}{2}(\partial_a \phi)^2 + Me^{\phi}), \tag{2.1.1}$$

describing induced gravity in the conformal gauge (i.e. $ds^2 = e^{\phi} dx^+ dx^-$), is in fact nothing but a constrained $SL_2(\mathbf{R})$ WZW model. This opens up the possibility to study Liouville theory, which is very complicated, using the more natural WZW variables. In fact it turns out that field configurations which are singular in the Liouville variables, but which do have regular energy-momentum densities (i.e. they are physically regular), are regular in the corresponding WZW variables. As we shall see the singularities in the Liouville variables arise, in the WZW picture, from the fact that Gauss decompositions of group elements are only defined on a dense subset of the corresponding group (in the Liouville case $SL_2(\mathbf{R})$).

The WZW model is described by the following action functional for (smooth) embeddings g of a 2 dimensional Riemann manifold (Σ, h) into a finite dimensional Lie group G.

$$S[g] = -\frac{k}{8\pi} \int d^2x \, \sqrt{h} h^{\alpha\beta}(x) \, \mathrm{Tr}(g^{-1}\partial_\alpha g \, g^{-1}\partial_\beta g) + \frac{k}{12\pi} \int_{B_3} \mathrm{Tr}(\tilde{g}^{-1} d\tilde{g})^3 \tag{2.1.2}$$

where B_3 is a 3 manifold whose boundary $\partial B_3 = \Sigma$ and $\tilde{g}$ is an extension of g from Σ to B_3. of course this action will depend on the choice of B_3 and on the extension $\tilde{g}$ but it can be shown that if k is integer then the Euclidean partition function is independent w.r.t. these choices. From now on we shall simply take $\Sigma = \mathbf{R} \times S^1$ equipped with the Minkowski metric.

The WZW action possesses left and right affine Lie symmetry, i.e. it is invariant under arbitrary transformations of the form [33, 25]

$$g(x^+, x^-) \mapsto \Omega(x^+) g(x^+, x^-) \bar{\Omega}(x^-)^{-1} \tag{2.1.3}$$

where Ω and $\bar{\Omega}$ are arbitrary group valued functions of the lightcone coordinates $x^{\pm}$. The Noether currents associated to this symmetry are

$$J = k\partial_+ g g^{-1} \quad \text{and} \quad \bar{J} = k g^{-1} \partial_- g \tag{2.1.4}$$

while the field equations, coinciding with the conservation laws of these currents, read

$$\partial_- J = \partial_+ \bar{J} = 0 \tag{2.1.5}$$

The currents J and $\bar{J}$ are Lie algebra valued functions on Σ which means that if $\{t_a\}$ is a basis of the Lie algebra of G then we can write $J = J^a t_a$ and $\bar{J} = \bar{J}^a t_a$ and the equations of motion are equivalent to $\partial_- J^a = \partial_+ \bar{J}^a = 0$ (for all a).

Let us now describe how Liouville theory arises from the $G = SL_2(\mathbf{R})$ WZW model [22]. For this we use the Gauss decomposition $g = ABC$ of an arbitrary group element, where $A = \exp(aE),\ B = \exp(\frac{1}{2}\phi H)$ and $C = \exp(bF)$. Here $\{H, E, F\}$ is the standard Cartan-Weyl basis of $SL_2(\mathbf{R})$, i.e.

$$H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \tag{2.1.6}$$

The Gauss decomposition is local in the sense that it is only well defined on a dense subset of $SL_2(\mathbf{R})$. Let us for the moment ignore this subtlety however. Also note that the arguments that follow below are not valid for the compact group $SU(2)$ due to the fact that it allows for no Gauss decomposition.

Inserting $g = ABC$ into the WZW action and using the well known 'Polyakov-Wiegman identity'

$$\begin{aligned} S[ABC] &= S[A] + S[B] + S[C] - \frac{k}{4\pi}\int d^2x\, \mathrm{Tr}((A^{-1}\partial_- A)(\partial_+ B)B^{-1} + \\ &\quad + (B^{-1}\partial_- B)(\partial_+ C\, C^{-1}) + (A^{-1}\partial_- A)B(\partial_+ C\, C^{-1})B^{-1}) \end{aligned}$$

we find that $S[g]$ reduces to the following action for the fields a, b and ϕ

$$S[a,b,\phi] = \frac{k}{8\pi}\int d^2x\, \left(\frac{1}{2}\partial_+\phi\partial_-\phi + 2(\partial_- a)(\partial_+ b)e^{-\phi}\right) \tag{2.1.7}$$

The equations of motion of this action are given by

$$\partial_-(\partial_+ b\, e^{-\phi}) = 0 \tag{2.1.8}$$

$$\partial_+(\partial_- a\, e^{-\phi}) = 0 \tag{2.1.9}$$

$$\partial_+\partial_-\phi + 2(\partial_- a)(\partial_+ b)e^{-\phi} = 0 \tag{2.1.10}$$

Obviously from the first two equations follows that

$$\partial_+ b = \mu(x^+)e^{\phi} \quad \text{and} \quad \partial_- a = \nu(x^-)e^{\phi} \tag{2.1.11}$$

where μ and ν are arbitary functions of x^+ and x^- respectively while the third equation reduces to

$$\partial_+\partial_-\phi(x) + 2M(x)e^{\phi(x)} = 0 \tag{2.1.12}$$

where $M(x) = 2\mu(x^+)\nu(x^-)$. Note that if M was a constant function then this equation would be nothing but the Liouville equation.

In all steps made above we have retained the full content of the WZW model. Now, the phase space of any mechanical model (i.e. the space of 'positions and momenta') can be described, at least locally, as the space of solutions of the equations of motion because through every point p in phase space goes one unique solution such that $t = 0$ at the time when this solution passes through p (put differently, the space of boundary conditions is equal to the phase space). From the above therefore follows that Liouville theory is a reduction of the WZW model in the sense that the Liouville phase space is equal to the subspace of the WZW phase space determined by the constraints that μ and ν are constant. On this reduced phase space the WZW equations of motion reduce to the Liouville equation.

A nice byproduct of this approach to Liouville theory is that the well known Leznov-Savaliev solutions of the Liouville equation can be derived easily from the general classical solution of the WZW model [22]

$$g(x^+, x^-) = g_L(x^+)g_R(x^-) \tag{2.1.13}$$

where g_L and g_R are arbitrary group valued functions constrained only by boundary conditions. The Leznov-Savaliev solutions can now be obtained from (2.1.13) by imposing the constraints and then inserting the Gauss decomposition.

As we have discussed above a WZW model has left and right affine Lie symmetry and one might wonder what happens to this symmetry after reduction. For this we have to consider the Noether currents associated to this symmetry a little bit closer. As we have seen the conservation laws induced by the affine symmetry are given by $\partial_- J^a = \partial_+ \bar{J}^a$. A standard calculation reveals that the (equal time) Poisson brackets between these currents read

$$\begin{aligned}
\{J^a(x), J^b(y)\} &= f_c^{ab} J^c(y)\delta(x-y) + kg^{ab}\delta'(x-y) \\
\{\bar{J}^a(x), \bar{J}^b(y)\} &= f_c^{ab} \bar{J}(y)^c\delta(x-y) + kg^{ab}\delta'(x-y) \\
\{J^a(x), \bar{J}^b(y)\} &= 0
\end{aligned} \tag{2.1.14}$$

where g^{ab} is the inverse of $g_{ab} = \mathrm{Tr}(t_a t_b)$. This current algebra will obviously be changed by the reduction process because some currents will be constrained. In order to see this consider again our example above. In terms of the Cartan-Weyl basis the current J can be written as $J(x) = J^H(x)H + J^E(x)E + J^F(x)F$ (and a similar equation for the

current $\bar{J}(x)$). Inserting the Gauss decomposition into the definition of J one finds that $J^E = k(\partial_+ b)\, e^{-\phi}$ and $\bar{J}^F = -k(\partial_- a)\, e^{-\phi}$ from which follows that after imposing the constraints we have $J^E(x) = \mu$ and $\bar{J}^F(x) = -\nu$. From a mathematical point of view one therefore has a Poisson algebra (2.1.14) subject to a set of constraints. Intuitively it is clear that the reduced current algebra will be equal to the Poisson algebra induced by (2.1.14) on the constrained phase space. Note that this algebra is *not* obtained by simpy inserting the constraints into (2.1.14). Let's illustrate this again with an example. Consider the Poisson bracket between J^H and J^E. From (2.1.14) follows that

$$\{J^H(x), J^E(y)\} \sim J^E(y)\delta(x-y) \tag{2.1.15}$$

On the constraint manifold $J^E = \mu$ and obviously a constant commutes with everything, so we must have $\{J^H(x), J^E(y)\} = \{J^H(x), \mu\} = 0$. The right hand side however of equation (2.1.15) is, after imposing the constraints, equal to $\mu\delta(x-y)$ which is nonzero. Blindly inserting the constraints therefore leads to inconsistencies. The proper way to deal with the construction of the reduced current algebra (2.1.14) is by 'Poisson reduction'. This is the subject of the next section.

2.2 Reduction of Poisson manifolds

In this section we shall briefly discuss the reduction of Poisson manifolds [77, 44]. This procedure will be used extensively in the next chapters in order to construct the Noether current algebras associated to certain reduced WZW models.

Let $(M, \{.,.\})$ be a Poisson manifold, i.e. M is a manifold and $\{.,.\}$ is an antisymmetric bilinear map from $C^\infty(M) \times C^\infty(M) \to C^\infty(M)$ which satisfies the Jacobi identities

$$\{f, \{g, h\}\} + \{g, \{h, f\}\} + \{h, \{f, g\}\} = 0 \tag{2.2.16}$$

where $f, g, h \in C^\infty(M)$. It is well known that any symplectic manifold is a Poisson manifold, but the converse need not be true. It *is* true however that a Poisson manifold is the disjoint union of symplectic submanifolds (we shall come back to this in the next chapter).

Any Poisson structure $\{.,.\}$ induces a map

$$P : T^*M \to TM \tag{2.2.17}$$

which is defined as follows: A given vectorfield $X \in \Gamma(TM)$ (= the space of smooth sections of TM) can be written as $X = g_i df^i$ where $g_i, f_i \in C^\infty(M)$. P is therefore uniquely

defined by putting $P(df) = X_f$ where X_f is the 'Hamiltonian vectorfield associated to f', i.e. $X_f = \{f, .\}$.

Now, let M_0 be a closed and embedded submanifold $i : M_0 \hookrightarrow M$ of M of codimension k. The manifold M_0 is completely characterized by $C^\infty(M_0)$, the space of smooth functions on M_0 and the relation between $C^\infty(M)$ and $C^\infty(M_0)$ can be described as follows: let I be the ideal in $C^\infty(M)$ of functions that vanish on M_0, i.e.

$$I = \{f \in C^\infty(M) \mid f(x) = 0, \text{ for all } x \in M_0\}, \tag{2.2.18}$$

then

$$C^\infty(M_0) \simeq C^\infty(M)/I \tag{2.2.19}$$

A similar construction holds for the space of vectorfields $\Gamma(TM_0)$: it is clear that any vectorfield on M_0 arises as the restriction of some vectorfield on M. However, the restriction of a vectorfield on M to M_0 is not, in general, a vectorfield on M_0 since it need not be tangent to M_0. Obviously the space of vectorfields on M which restrict to vectorfields on M_0 is given by

$$\mathcal{N}(I) = \{X \in \Gamma(TM) \mid X(f) \in I, \text{ for all } f \in I\} \tag{2.2.20}$$

because an element f of I is constant on M_0 which means that the derivative of f in a direction tangent to M_0 must be zero. of course two vectorfields $X_1, X_2 \in \mathcal{N}(I)$ may restrict to the same vectorfield on M_0. This happens when their difference vanishes on M_0, or put differently when $X_1 - X_2 \in I\mathcal{N}(I)$. We therefore find

$$\Gamma(TM_0) \simeq \frac{\mathcal{N}(I)}{I\mathcal{N}(I)} \tag{2.2.21}$$

It is also easy to see that the set of functions $f \in C^\infty(M)$ which have the property that their Hamiltonian vectorfields X_f are tangent to M_0 is given by

$$N(I) = \{f \in C^\infty(M) \mid \{f, g\} \in I, \text{ for all } g \in I\} \tag{2.2.22}$$

Note that $N(I)$ is a Poisson subalgebra of $C^\infty(M)$ and that it is in fact the normalizer of I (I is an ideal w.r.t. the multiplicative structure on $C^\infty(M)$ but not necessarily w.r.t. the Poisson structure). Elements of $N(I)$ are called 'first class'.

It is well known in physics that first class elements generate gauge invariances [18]. These gauge invariances cause degeneracies which have the result that the Poisson structure on M does *not* induce a Poisson structure on M_0 but on a certain coset space of M_0. Let us make this statement more precise.

Let i^*TM and i^*T^*M denote the pullbacks of the tangent cotangent bundle to M_0 via the embedding i. TM_0 and T^*M_0 are subbundles of i^*TM and i^*T^*M respectively. Define now another subbundle, called the annihilator bundle $\mathrm{Ann}(TM_0)$, of i^*T^*M as follows: The fiber over $p \in M_0$ is given by

$$\mathrm{Ann}_p(TM_0) = \{\alpha \in i^*T_p^*M \mid \alpha(v) = 0, \text{ for all } v \in T_pM_0\} \tag{2.2.23}$$

The image under P of the annihilator bundle is denoted by $TM_0^\perp$, i.e.

$$T_pM_0^\perp = \{p(\alpha) \mid \alpha \in \mathrm{Ann}_p(TM_0)\} \tag{2.2.24}$$

Now, using $TM^\perp$ we can define the so called *null distribution* V of M_0 which is the intersection of TM_0 and $TM_0^\perp$

$$V_p = T_pM_0 \bigcap T_pM_0^\perp \tag{2.2.25}$$

Let us explain these definitions. The annihilator bundle consists roughly of those one forms that are exterior differentials of elements of I, for let $f \in I$ and $v \in TM_0$ then $df(v) \equiv v(f) = 0$ because f is constant on M_0. $TM_0^\perp$ is then nothing but the span of the Hamiltonian vectorfields of elements of I. Not all elements of I are however first class and since the Hamiltonian vectorfields of first class constraints are tangent to M_0 we can identify the gauge directions by considering the intersection of $TM_0^\perp$ and TM_0. The null distribution is therefore nothing but an indication as to what directions on M_0 are gauge directions.

It can be shown [44] that the null distribution V is an involutive distribution which means, by Frobenius' theorem, that M_0 foliates into maximal connected submanifolds or leaves of 'gauge equivalent states'. One can eliminate all gauge freedom by identifying all points on a leaf or equivalently by 'gauge fixing'. Gauge fixing amounts to finding a submanifold of M_0 (called the 'gauge slice') that has exactly one point in common with every leaf. On the gauge slice, denoted by $\tilde{M}$, the Poisson structure $\{.,.\}$ then induces a reduced Poisson structure $\{.,.\}^*$ [76, 44].

Let us now make the above somewhat more concrete. Again consider a Poisson manifold $(M, \{.,.\}$ and let the submanifold M_0 be given as the zero set of a collection of 'constraints' $\{\phi^i\} \subset C^\infty(M)$, i.e.

$$M_0 = \{p \in M \mid \phi^i(p) = 0, \text{ for all } i\} \tag{2.2.26}$$

I is then the ideal in $C^\infty(M)$ generated by the constraints

$$I = \{f_i\phi^i \mid f_i \in C^\infty(M) \text{ for all } i\} \tag{2.2.27}$$

Suppose the gauge invariances generated by the first class constraints (not all constraints need to be first class) can be completely fixed by adding some 'gauge fixing constraints', i.e. constraints that determine the gauge slice. Denoting the *total* set of constraints by $\{\phi^\mu\}$ (i.e. with Greek indices) the gauge slice is given by

$$\tilde{M} = \{p \in M \mid \phi^\mu(p) = 0, \text{ for all } \mu\} \tag{2.2.28}$$

and the reduced Poisson structure is [18, 78]

$$\{\bar{f}, \bar{g}\}^* = \overline{\{f, g\} - \{f, \phi^\mu\}\Delta_{\mu\nu}\{\phi^\nu, g\}} \tag{2.2.29}$$

where $f, g \in C^\infty(M)$, the bar denotes the restriction to $\tilde{M}$ (which is a submanifold of M) and $\Delta_{\mu\nu}$ is the inverse of $\Delta^{\mu\nu} \equiv \{\phi^\mu, \phi^\nu\}$. The bracket (2.2.29) was originally discovered by Dirac and is therefore called the Dirac bracket.

Let us now return to our example of Liouville theory. The Poisson algebra we are dealing with in this case is (2.1.14) for $SL_2(\mathbf{R})$ subject to the constraint $\phi^1(x) \equiv J^E(x) - \mu = 0$. Since the Poisson bracket of $\phi(x)$ with itself is zero this constraint is first class and therefore generates gauge invariances. This gauge invariance has to be fixed and it will be shown in the next chapters that one can fix it completely by adding the constraint $\phi^2(x) \equiv J^H = 0$. Calculating the Poisson relation between $J^E(x)$ and $J^E(y)$ with respect to (a field theoretical version of) the Dirac bracket one finds that they satisfy the Virasoro algebra

$$\{J^E(x), J^E(y)\}^* = \partial J^E(y)\delta(x-y) - 2J^E(y)\delta'(x-y) - \frac{1}{2}\delta'''(x-y) \tag{2.2.30}$$

This is then the current algebra associated to Liouvilly theory.

Above we have seen that the affine $SL_2(\mathbf{R})$ algebra leads, on reduction, to the Virasoro algebra. One might wonder if it is possible to obtain more complicated W algebras in this way by reducing affine $SL_N(\mathbf{R})$ algebras (or more generally affine Lie algebras over arbitrary simple Lie algebras). This is precisely the question we address in chapter 4. It is clear that affine $SL_2(\mathbf{R})$ subalgebras will play an important role in this theory because any W algebra must contain a Virasoro algebra and we have just seen that this algebra is a reduction of the affine $SL_2(\mathbf{R})$ algebra. However, there are in general many inequivalent affine $SL_2(\mathbf{R})$ subalgebras into a given affine Lie algebra and it is therefore natural to conjecture that there is a W algebra associated to every $SL_2(\mathbf{R})$ embedding. This conjecture will be proven in chapter 4.

2.3 Gauged WZW models

Above we have constructed Liouville theory as a constrained WZW model. The procedure we followed was that of contraining the WZW phase space by essentially specifying the solution space of the reduced theory. In order to quantize Liouville theory it would however be much more useful to have available a Langrangian realization of the WZW-Liouville reduction. The natural framework for this is the theory of gauged WZW models [7].

We have seen above that the WZW action (2.1.2) is invariant under left and right affine transformations (2.1.3). Note that the group valued functions Ω and $\bar{\Omega}$ are functions of one lightcone coordinate only. Consider now two isomorphic subgroups H and $\bar{H}$ of G. The affinizations $\hat{H}$ and $\hat{\bar{H}}$ of these groups (consisting of H ($\bar{H}$) valued functions of the coordinates $x^+(x^-)$) are subgroups of the left-right affine symmetry of the WZW lagrangian which means that $S[g]$ certainly has $\hat{H}\times\hat{\bar{H}}$ symmetry. Gauging this symmetry means, in the WZW context, making the action invariant under transformations

$$g(x^+x^-)\mapsto \alpha(x^+,x^-)\,g(x^+,x^-)\,\beta^{-1}(x^+,x^-) \tag{2.3.31}$$

where α and β are H respectively $\bar{H}$ valued functions *of both* x^+ *and* x^-. For this one introduces $\hat{h}$ respectively $\hat{\bar{h}}$ valued gauge fields A_+ and A_- (where $\hat{h}$ ($\hat{\bar{h}}$) is the Lie algebra of $\hat{H}$ ($\hat{\bar{H}}$) which transform as usual under gauge transformations

$$\begin{aligned} A_+ &\mapsto \alpha A_+\alpha^{-1}+\partial_+\alpha\,\alpha^{-1} \\ A_- &\mapsto \beta A_-\beta^{-1}+\beta^{-1}\partial_-\beta \end{aligned} \tag{2.3.32}$$

A 'gauged' WZW action that is invariant (or more accurately, invariant up to a total derivative) under the transformations (2.3.31) and (2.3.32) is

$$\begin{aligned} I[g,A_+,A_-] &= S[g]+\int d^2x\,(\mathrm{Tr}(A_-(\partial_+g\,g^{-1}-M))+ \\ &\quad +\mathrm{Tr}(A_+(g^{-1}\partial_-g-\tilde{M}))+\mathrm{Tr}(A_-gA_+g^{-1})) \end{aligned} \tag{2.3.33}$$

where M and $\tilde{M}$ are the constraints one wishes to impose on the system (in the case of Liouville theory discussed above $M=\mu E$ and $\tilde{M}=\nu F$).

It is clear that one can use the gauge invariance 2.3.31 of the action (2.3.33) to put A and B in the Gauss decomposition $g=ABC$ equal to the unit element e of G. For the $SL_2(\mathbf{R})$ case this implies $g=\exp(\frac{1}{2}\phi H)$. From the equations of motion of the fields $A_\pm$ one then finds that $A_+=B^{-1}MB$ and $A_-=B\tilde{M}B^{-1}$. Substituting this, together with the fact that $g=B$ back into the action (2.3.33) one finds that it reduces to the Liouville action. What happens in the general case (i.e. for arbitary algebras) will be examined in chapter 5.

In the literature there are two approaches to gauged WZW models corresponding (on the algebraic level) to the reduction approach discussed above and the coset construction respectively. Let H be a subgroup of G and let $A = A_+dx^+ + A_-dx^-$ be an h valued gauge field (where again h is the Lie algebra of H). Consider instead of (2.3.33) the action

$$\tilde{I}[g, A_-, A_+] = S[g] - \frac{k}{4\pi}\int d^2x \,\mathrm{Tr}(A_-(\partial_+ g)g^{-1} + (g^{-1}\partial_- g)A_+ + A_- g A_+ g^{-1} - A_- A_+) \tag{2.3.34}$$

This is called the action of a G/H WZW model and it has been shown in [14] that it provides a Lagrangian realization of the Goddard-Kent-Olive coset construction. Note the difference with the action (2.3.33): the fields A_-, A_+ are now components of an h valued gauge field while in (2.3.33) they are completely independent.

2.4 Quantum framework for constrained WZW models

In the previous section we have considered the classical problem of reduction of the WZW model to Liouville theory. As we have seen this reduction has an algebraic counterpart relating the chiral Noether symmetry algebras associated to these two theories. In the quantum case the Hilbert space of the WZW model will carry a unitary representation of the affine Lie algebra while the Liouville Hilbert space will decompose into Virasoro representations. The problem is to construct the Liouville Hilbert space as a reduction of the WZW Hilbert space and to calculate the Liouville spectrum from the WZW spectrum. The formalism ideally suited to handle problems like this one is the BRST formalism [34].

Suppose one is given any (classical) physical system together with a set of first class constraints $\{\phi_i\}$, i.e.

$$\{\phi_i, \phi_j\} = C_{ij}^k \phi_k \tag{2.4.35}$$

where C_{ij}^k may in principle be arbitrary functions on the phase space of the system. In what follows we will however always assume these 'structure constants' to be constant. The meaning of this relation is that the Poisson bracket between first class constraints vanishes on the constrained manifold from which follows that the Hamiltonian vector fields of first class constraints are tangent to it. As Dirac showed the directions on phase space specified by these first class constraints via their Hamiltonian vectorfields are gauge directions and have to be eliminated. In principle one could first reduce the system to a gauge slice and then quantize it but in practise it is easier to do the opposite, i.e. quantize first and reduce later. of course one has to assume that one is able to quantize the unconstrained theory (which in the WZW case is certainly the case [25]).

Suppose therefore that we have quantized our original system yielding a Hilbert space $\mathcal{H}$. BRST quantization now involves introducing a set of 'ghosts' c^i and 'antighosts' b_i which obey canonical anticommutation relations $c^i b_j + b_j c^i = \delta^i_j$. Mathematically one one has extended the phase space by adding (Grassman) degrees of freedom. In the end one will have to remove all the Grassman variables from the theory together with the redundant degrees of freedom associated to the gauge directions. The way one goes about this is as follows: Introduce the operator

$$Q = c^i \phi_i - \frac{1}{2} C^k_{ij} c^i c^j b_k \tag{2.4.36}$$

which is known as the BRST operator. In the second term one needs, in the infinite dimensional case, a normal ordening prescription but we will ignore this subtlety in this chapter (we do come back to this in chapter 4 however). Also it should be mentioned that if the structure constants would have been structure functions (i.e. funcions on phase space) then the BRST operator could have picked up higher order terms. How to construct the BRST operator in the general case can be found in [5].

The basic property of the BRST operator is that $Q^2 = 0$ and that it maps the ghost number n subspace $\mathcal{H}_n$ of the (extended) Hilbert space (i.e. the eigenspace of the ghost number operator $N = c^i b_i$ with eigenvalue n) to the ghost number $n+1$ subspace, i.e.

$$Q : \mathcal{H}_n \to \mathcal{H}_{n+1} \tag{2.4.37}$$

A state is now called 'BRST invariant' if $Q|\psi\rangle = 0$. Since the ghosts were artificially put into the theory it is clear that the BRST invariant states of ghost number 0 are of special interest. For such a state we have $N|\psi\rangle = 0$ which means that $b_i|\psi\rangle = 0$ for all i which means that the second term in the BRST operator vanishes on the ghost number zero subspace. In fact one has

$$Q|\psi\rangle = c^i f_i|\psi\rangle \tag{2.4.38}$$

and since a state annihilated by all b_i cannot be annihilated by all c^i we find that the state is BRST invariant iff $\phi_i|\psi\rangle = 0$. We conclude that a state of ghost number 0 is BRST invariant iff it is annihilated by the (quantum) constraints. Therefore BRST invariant states are of great interest.

There are however states that are trivially BRST invariant, namely those states $|\psi\rangle$ for which there exists a state $|\chi\rangle$ such that $|\psi\rangle = Q|\chi\rangle$ (these states are called BRST exact). Let us therefore consider two states to be equivalent if their difference is BRST exact. Heuristically one could say that BRST equivalence corresponds to gauge equivalence which means that one should not distinguish physically between BRST equivalent states.

From the above we find that the true Hilbert space is the space of BRST invariant states at ghost number zero modulo the set of BRST exact states. Mathematically this is called the zeroth cohomology of the BRST complex. of course similar arguments are in force with respect to the observables in the theory, i.e. the physically interesting observables $\mathcal{O}$ of the reduced theory are those which are BRST invariant $[Q, \mathcal{O}] = 0$ but not BRST exact $\mathcal{O} \neq [Q, \mathcal{O}_2]$.

Let us again apply this on Liouville theory. As we have said the WZW Hilbert space carries a (reducible) representation of the underlying affine Lie algebra. This algebra is given in terms of operator product expansions by

$$J^a(z)J^b(w) = \frac{kg^{ab}}{(z-w)^2} + \frac{f_c^{ab}J^c(w)}{z-w} + \dots \tag{2.4.39}$$

and as we have seen the constraints are given in the Liouville case by $\phi(z) = J^E(z) - 1$. Introducing the ghost field $c(z)$, the antighost field $b(z)$ satisfying the OPE

$$c(z)b(w) = \frac{1}{z-w} + \dots \tag{2.4.40}$$

and using the prescription outlined above we find that the BRST operator is given by

$$Q = (J^E(z) - 1)c(z) \tag{2.4.41}$$

The total Hilbertspace is then given by

$$\mathcal{H}_{tot} = \mathcal{H}_{WZW} \otimes \mathcal{H}_{ghost} \tag{2.4.42}$$

and what we need to do is calculate the zeroth cohomology of Q. In the case of Liouville theory this was done in [6] using free field resolutions of affine and Virasoro representations. They also showed that the cohomology space carries a Virasoro representation of central charge

$$c(k) = 13 - 6(k+2) - \frac{6}{k+2} \tag{2.4.43}$$

This Virasoro algebra is the chiral current algebra of Liouville theory.

In this chapter we have seen how Liouville theory arises as a (conformally) reduced WZW model. In chapter 4 we are going to generalize this construction to general affine Lie algebras and arbitrary reductions. The methods we are going to use for this are however essentially the same as the ones that we explained in this chapter.

Chapter 3

Finite W algebras

Introduction

Starting with the seminal work of Weyl and Wigner symmetry principles have been playing an ever increasing role in theoretical physics. The simplest symmetries are those in which the physical system is symmetric under a finite number of transformations or when it is invariant under displacement in a finite number of 'directions'. The natural mathematical structures describing such symmetries are discrete groups and finite dimensional Lie groups respectively. In this century the theory of these mathematical structures has, motivated by physics, been brought to a high level of development. This has lead to many new insights into the principles of nature and in fact nowadays symmetry principles play a role in physics that is as fundamental as for example the action principle. A paramount example of this is invariance under the Poincare group, which is at the heart of Einstein's special theory of relativity and hence is a necessary criterion for any realistic theory of nature.

Another sacred principle of fundamental physics is locality or causality. Combining symmetry principles with locality one is naturally lead to the study infinite dimensional groups of smooth maps from some spacetime manifold X to a finite dimensional Lie group G. The group multiplication in such a group is just pointwise multiplication, i.e. if $f, g \in Map(X, G)$ and $a \in G$ then $(f.g)(a) = f(a)g(a)$. In quantum field theory groups of the form $Map(X, G)$ and their Lie algebras $Map(X, g)$ (where g is the Lie algebra of G) arise essentially in two different ways: through the principle of local gauge invariance, which is at the heart of modern high energy physics (i.e. the standard model of the fundamental interactions between quarks and leptons), and through the theory of current groups and algebras.

Unfortunately for generic manifolds X surprisingly little is known about the group $Map(X, G)$. Especially the representation theory of these groups is still almost unex-

plored. The exception to this is the case $X = S^1$ where $Map(S^1, G)$ and $Map(S^1, g)$ are called 'loop groups' and 'loop algebras'. Loop groups and algebras arise in simplified models of quantum field theory in which space is taken to be 1-dimensional and therefore also in string models of elementary particles. The study of loop groups and algebras is much simpler than when X is some more complicated manifold. This is caused by the fact that they behave much like the ordinary finite dimensional Lie groups and algebras that underly them. This remarkable fact makes knowledge of the finite dimensional theory essential for the study of the infinite dimensional theory.

Compared to Lie groups and algebras the theory of W algebras has evolved backwards, for up to now infinite dimensional W algebras have received the bulk of the attention. One reason for this is that it was not entirely clear what the finite algebras underlying W algebras were and whether there was a finite version of W theory at all. Considering the way in which W algebras were first introduced into physics they really don't seem to have any relation to the theory of loop groups and algebras. Therefore finite structures underlying W algebras have not been considered. In this chapter we shall address the question: does the theory of W algebras have a finite counterpart? The answer to this question turns out to be affirmative. In fact it will be shown that this finite theory is remarkably rich and, as with loop algebras, contains already most of the essential features of infinite dimensional W algebra theory.

In this chapter we approach the problem of constructing finite W theory in the same way as we approach the theory of infinite dimensional W algebras, that is we consider W algebras (finite or infinite) to be reductions of Lie algebras. In the next chapter we will show that infinite W algebras are reductions of affine Lie algebras (i.e. central extensions of certain loop algebras). Our approach to constructing finite W theory is by mimicing the reduction procedure on finite dimensional Lie algebras. As we shall see this leads to a rich theory of finitely generated nonlinear algebras.

This chapter consists of three parts. The first part deals with the classical theory of finite W algebras. Classical finite W algebras will be constructed as Poisson reductions of the Kirillov Poisson structure on the dual of a simple Lie algebra. In this section finite W algebras are therefore finitely generated nonlinear Poisson algebras. The classical representation theory of these algebras is considered, i.e. we calculate the symplectic leaves associated to arbitrary finite W algebras. In the second part of this chapter we quantize these classical finite W algebras. This is done by the well known BRST quantization procedure which is ideally suited for situations like this. In the third and last part of this chapter we construct the representation theory of quantum finite W algebras. As it turns out the representation theory of semisimple Lie algebras is in essence all one needs to construct these representations. We also give a general procedure to obtain Fock realizations of arbitrary finite W algebras.

3.1 The Classical Theory of Finite W algebras

In this section we introduce and develop the theory of finite W algebras. As we already mentioned, we will show in the next chapter that many classical W algebras can be obtained by Poisson reduction of the Kirillov Poisson algebras naturally associated to affine Lie algebras. Given a certain affine Lie algebra there are several possible reductions all leading to different W algebras. In fact we will show that to any sl_2 embedding into the simple Lie algebra underlying the affine Lie algebra one can associate such a Poisson reduction. Motivated by this we consider in this chapter analogous Poisson reductions of Kirillov Poisson algebras associated to finite dimensional simple Lie algebras. Obviously this is the natural finite dimensional analogue of the procedure that leads to infinite W algebras. The finitely generated Poisson algebras one obtains in this way we define to be finite W algebras. In order to make the presentation more or less selfcontained we shall first briefly review some basic aspects of Kirillov Poisson structures.

3.1.1 Kirillov Poisson structures

Let $(M, \{.,.\})$ be a Poisson manifold, that is $\{.,.\}$ is a Poisson bracket on the space $C^\infty(M)$ of C^∞ functions on M, and G a Lie group. Also let $\Phi : G \times M \to M$ be a smooth and proper action of G on M which preserves the Poisson structure, i.e.

$$\Phi_g^*\{\phi, \psi\} = \{\Phi_g^*(\phi), \Phi_g^*(\psi)\} \tag{3.1.1}$$

where Φ_g^* is the pullback of $\Phi_g : M \to M$. Physically this implies that if $\gamma(t)$ is a solution of the equations of motion w.r.t. some G-invariant Hamiltonian H (i.e. $\Phi_g^* H = H$ where as usual $\Phi_g^* H = H \circ \Phi_g$), then $(\Phi_g \circ \gamma)(t)$ is again a solution. If we do not want to consider solutions that can be transformed into each other in this way to be essentially different we are led to consider the space M/G as the 'true' phase space of the system. The only observables (i.e. functions on M) that one is concerned with are those which are themselves G-invariant and therefore descend to observables on M/G. Denote the set of smooth G-invariant functions on M by O, and let then $\phi, \psi \in O$. Using the fact that Φ_g preserves the Poisson structure for all $g \in G$ and that ϕ and ψ are G-invariant we can easily deduce that $\{\phi, \psi\}$ is also an element of O, i.e. O is a Poisson subalgebra of $C^\infty(M)$. Let $\pi : M \to M/G$ be the canonical projection. The pullback map $\pi^* : C^\infty(M/G) \to O$ is in fact an isomorphism. It assigns to a function $\hat{\phi}$ on M/G the function $\hat{\phi} \circ \pi$ on M which is constant along the G-orbits. What one wants is now to define a Poisson structure $\{.,.\}^*$ on $C^\infty(M/G)$ such that the Poisson algebras $(C^\infty(M/G), \{.,.\}^*)$ and $(O, \{.,.\})$ are isomorphic. This would mean that all (G-invariant) information of the original phase space is transferred to the Poisson algebra $(C^\infty(M/G), \{.,.\}^*)$. Therefore define

$$\{\hat{\phi}, \hat{\psi}\}^* = (\pi^*)^{-1}\{\pi^*\hat{\phi}, \pi^*\hat{\psi}\} \tag{3.1.2}$$

for all $\hat{\phi}, \hat{\psi} \in C^\infty(M/G)$. Obviously π^* is a Poisson algebra isomorphism between the two Poisson algebras. So starting from a G-invariant theory we have arrived at a formulation in which all essential information is contained and all redundancy has been eliminated.

Consider now a G-invariant theory in which the group manifold itself is the configuration space (for example a particle moving on G with a Hamiltonian that is invariant under G). The phase space of such a system is of course T^*G, the cotangent bundle of G. The left action of the group on itself, denoted by L_g, induces a left $G-$action L_g^* on T^*G which is in fact a Poisson action w.r.t. the canonical Poisson structure on T^*G (T^*G carries a natural Poisson structure because it is a cotangent bundle). Therefore we can do what we did before and extract the G-invariant piece of the theory by going to the quotient manifold

$$T^*G/G \tag{3.1.3}$$

Obviously this space is nothing but the space of left invariant one forms which means that it is isomorphic to g^*, the dual of the Lie algebra of G. The canonical projection

$$\pi : T^*G \to g^* \tag{3.1.4}$$

is therefore simply the pullback to the fiber over the identity element of G by means of left translation. As before the Poisson structure on T^*G induces a Poisson structure on g^*. This Poisson structure on g^* is called the 'Kirillov' Poisson structure.

It is possible to obtain (by explicit calculation) a more concrete formula for the Kirillov Poisson structure [76]. It reads

$$\{F, G\}(\xi) = \xi([\mathrm{grad}_\xi F, \mathrm{grad}_\xi G]) \tag{3.1.5}$$

where F and G are smooth functions on g^*, $\xi \in g^*$ and $\mathrm{grad}_\xi F \in g$ is determined uniquely by

$$\frac{d}{d\epsilon} F(\xi + \epsilon\eta)|_{\epsilon=0} = \eta(\mathrm{grad}_\xi F) \tag{3.1.6}$$

for all $\eta \in g^*$. In this paper we will always take G to be a simple group which means that the Cartan-Killing form $(.,.)$ on its Lie algebra is non-degenerate. This allows us to identify the Lie algebra g with its dual, i.e. all $\xi \in g^*$ are of the form $(\alpha, .)$ where $\alpha \in g$. The above formulas then become

$$\{F, G\}(\alpha) = (\alpha, [\mathrm{grad}_\alpha F, \mathrm{grad}_\alpha G]) \tag{3.1.7}$$

and

$$\frac{d}{d\epsilon} F(\alpha + \epsilon\beta)|_{\epsilon=0} = (\beta, \mathrm{grad}_\alpha F) \tag{3.1.8}$$

where $\alpha, \beta \in g$ and $F, G \in C^\infty(g)$. This is the form that we will use.

It is possible to give an even more detailed description of the Kirillov bracket. Let $\{t_a\}$ be a basis of g and let $\{J^a\}$ be the dual basis, i.e. $J^a(t_b) = \delta^a_b$. of course the functions J^a are smooth functions on g and we can calculate the Kirillov Poisson bracket between them. What one finds is

$$\{J^a, J^b\} = f^{ab}_c J^c \tag{3.1.9}$$

where f^c_{ab} are the structure constants of g in the basis $\{t_a\}$ and indices are raised and lowered by the Cartan-Killing metric. Note the resemblance with the Lie bracket

$$[t_a, t_b] = f^c_{ab} t_c \tag{3.1.10}$$

However we have to remember that the Poisson bracket (3.1.9) is defined on the space of smooth functions on g which is a Poisson algebra, i.e. it also carries a commutative multiplication compatible with the Poisson structure. This means for example that the expression $J^a J^b J^c$ makes sense, while $t_a t_b t_c$ does not (remember that if F, G are two functions on g then their product is defined by $(FG)(\alpha) = F(\alpha)G(\alpha)$).

3.1.2 sl_2-embeddings

As we have mentioned before the inequivalent Poisson reductions of a Kirillov Poisson algebra are labelled by the equivalence classes of sl_2 embeddings into the simple Lie algebra in question. Let us therefore take a moment to recall some basic facts on sl_2 embeddings [79] and prove some lemmas that we shall need later on.

Let $i : sl_2 \hookrightarrow g$ be an embedding of sl_2 into a simple Lie algebra g and let $L(\Lambda; n)$ be the irreducible finite dimensional representation of g with highest weight Λ. As a representation of the subalgebra $i(sl_2)$ of g, $L(\Lambda; n)$ need not be irreducible. In general it decomposes into a direct sum of irreducible sl_2 representations

$$L(\Lambda; n) \simeq \bigoplus_{j \in \frac{1}{2}\mathbf{N}} n_j(\Lambda; i) \cdot \underline{2j+1}_2 \tag{3.1.11}$$

where $\underline{2j+1}_2$ denotes the $(2j+1)$-dimensional irreducible representation of sl_2 and $n_j(\Lambda; i)$ are the multiplicities of these representations in the decomposition. The set of numbers $\{n_j(\Lambda; i)\}_j$ is called the 'branching rule' of the representation $L(\Lambda; n)$ under the sl_2 embedding i. In what follows $L(\Lambda; n)$ will always either be the ($n-$dimensional) fundamental or the ($n^2 - 1$ dimensional) adjoint representation of sl_n which we simply denote by $\underline{n}_n$ and $\underline{ad}_n$ respectively.

Two sl_2 embeddings i and i' are called equivalent if there exists an inner automorphism ψ of g such that

$$i' = \psi \circ i \tag{3.1.12}$$

It is easy to show that if $i \sim i'$ then $n_j(\Lambda; i) = n_j(\Lambda; i')$ for all representations $L(\Lambda; n)$ and $j \in \frac{1}{2}\mathbf{N}$. The converse is not so clear. However, for $g = sl_n$ it can be shown that [79]

1. The inequivalent sl_2 embeddings into sl_n are completely characterized by the branching rule of the fundamental representation, i.e. two sl_2 embeddings are equivalent if and only if the branching rules of the fundamental representation w.r.t. these two embeddings are equal.

2. There exist $P(n)$ inequivalent sl_2 embeddings into sl_n, where $P(n)$ is the number of partitions of the number n.

The set of equivalence classes of sl_2 embeddings into sl_n is therefore parametrized by the branching rules of the fundamental representation, while the different branching rules that are possible are simply given by the partitions of n.

Example: Let $g = sl_4$. There are 5 equivalence classes of sl_2 embeddings into sl_4 characterized by the following decompositions

1. $\underline{4}_4 \simeq \underline{4}_2$
2. $\underline{4}_4 \simeq \underline{3}_2 \oplus \underline{1}_2$
3. $\underline{4}_4 \simeq 2 \cdot \underline{2}_2$
4. $\underline{4}_4 \simeq \underline{2}_2 \oplus 2 \cdot \underline{1}_2$
5. $\underline{4}_4 \simeq 4 \cdot \underline{1}_2$

of the fundamental representation $\underline{4}_4$ of sl_4.

The embeddings that are characterized by the fact that the fundamental representation becomes an irreducible representation of the embedded algebra are called 'principal embeddings'. If the fundamental representation branches into singlets only the embedding is called trivial.

In what follows we shall also need the centralizer $C(i)$ of $i(sl_2)$ in $g = sl_n$. It corresponds to the singlets of the adjoint action of $i(sl_2)$ in g. In order to describe $C(i)$ we have to take a look at the branching rule of the fundamental representation

$$\underline{n}_n \simeq \bigoplus_{j \in \frac{1}{2}\mathbf{N}} n_j \cdot \underline{2j+1}_2 \tag{3.1.13}$$

where for convenience we simply denote $n_j \equiv n_j(\Lambda; i)$ for the fundamental representation. Let q be the number of different values for j appearing in this decomposition for which n_j is non-zero, then $C(i)$ is given by

$$C(i) \simeq \bigoplus_{j \in \frac{1}{2}\mathbf{N}} sl_{n_j} \bigoplus (q-1)u(1) \tag{3.1.14}$$

so in essence it is a direct sum of sl_{n_j} algebras.

From the branching of the fundamental representation one can deduce the branching of the (n^2-1)-dimensional adjoint representation $\underline{ad}_n$ of $g = sl_n$. It reads

$$\begin{aligned} \underline{ad}_n \oplus \underline{1}_n \quad \simeq \quad & \bigoplus_{j \in \mathbf{N}} n_j^2 \cdot (\underline{1}_2 \oplus \underline{3}_2 \oplus \ldots \oplus \underline{4j+1}_2) \\ & \bigoplus_{j \neq j'} n_j n_{j'} \cdot (\underline{2|j-j'|+1}_2 \oplus \ldots \oplus \underline{2|j+j'|+1}_2) \end{aligned} \tag{3.1.15}$$

where the singlet on the left hand side was added to indicate that in order to obtain the branching rule of the adjoint representation $\underline{ad}_n$ we still have to 'subtract' a singlet from the righthand side.

Let $\{t_0, t_+, t_-\}$ be the standard generators of $i(sl_2)$. The Cartan element t_0, called the defining vector of the embedding, can always be chosen to be an element of the Cartan subalgebra of g [79]. Therefore it defines a $\frac{1}{2}\mathbf{Z}$ gradation of g given by

$$g = \bigoplus_{m \in \frac{1}{2}\mathbf{Z}} g^{(m)} \ ; \tag{3.1.16}$$

where $g^{(m)} = \{x \in g \mid [t_0, x] = mx\}$. We can choose a basis

$$\{t^{(\mu)}_{j,m}\}_{j \in \frac{1}{2}\mathbf{N};\ -j \leq m \leq j;\ 1 \leq \mu \leq n_j^{ad}} \tag{3.1.17}$$

for $g = sl_n$ (where n_j^{ad} is the multiplicity of the spin j representation of sl_2 in the branching of the adjoint representation) such that

$$\begin{aligned} [t_3, t^{(\mu)}_{j,m}] &= m t^{(\mu)}_{j,m} \\ [t_\pm, t^{(\mu)}_{j,m}] &= c(j,m) t^{(\mu)}_{j,m\pm 1} \end{aligned} \tag{3.1.18}$$

where $c(j,m)$ are standard normalization factors. We will always take the labeling to be such that $t^{(1)}_{1,\pm 1} \equiv t_\pm$ and $t^{(1)}_{1,0} \equiv t_0$. It is clear that

$$g^{(m)} = \bigoplus_{j,\mu} \mathbb{C} t^{(\mu)}_{j,m} \tag{3.1.19}$$

Note that the spaces $g^{(m)}$ and $g^{(-m)}$ are always of the same dimension. We have the following

Lemma 1 *The spaces $g^{(m)}$ and $g^{(n)}$ are orthogonal w.r.t. the Cartan-Killing form on g, i.e.*

$$(g^{(m)}, g^{(n)}) = 0 \tag{3.1.20}$$

iff $m \neq -n$.

Proof: Let $x \in g^{(m)}$ and $y \in g^{(n)}$, then obviously

$$([t_0, x], y) = m(x, y) \tag{3.1.21}$$

but also

$$([t_0, x], y) = -(x, [t_0, y]) = -n(x, y) \tag{3.1.22}$$

where we used the invariance property of the Cartan Killing form. It follows immediately that $(n + m)(x, y) = 0$. Therefore if $n \neq -m$ we must have $(x, y) = 0$. This proves the lemma .

For notational convenience we shall sometimes denote the basis elements $t^{(\mu)}_{j,m}$ simply by t_a where a is now the multi-index $a \equiv (j, m; \mu)$. Let K_{ab} denote the matrix components of the Cartan-Killing form in this basis, i.e. $K_{ab} = (t_a, t_b)$ and let K^{ab} denote its matrix inverse.

From the above lemma and the fact that the Cartan-Killing form is non-degenerate on g follows immediately that $g^{(k)}$ and $g^{(-k)}$ are non-degenerately paired. This implies that if $t_a \in g^{(k)}$ then $K^{ab}t_b \in g^{(-k)}$ (where we used summation convention).

We then have the following lemma which we shall need later.

Lemma 2 *If t_a is a highest weight vector (i.e. $a = (j, j; \mu)$ for some j and μ) then $K^{ab}t_b$ is a lowest weight vector (and vica versa). In particular if $t_a \in C(i)$ then $K^{ab}t_b \in C(i)$.*

Proof: Since t_a is a highest weight vector we have

$$\begin{aligned} 0 &= ([t_a, t_+], x) \\ &= (t_a, [t_+, x]) \end{aligned} \tag{3.1.23}$$

for all $x \in g$ which means that t_a is orthogonal to $\mathrm{Im}(ad_{t_+})$ or put differently

$$\mathrm{Ker}(ad_{t_+}) \perp \mathrm{Im}(ad_{t_+}) \tag{3.1.24}$$

It is easy to do the same thing for t_-. One therefore has the following decomposition of g into mutually orthogonal spaces

$$g = \Big(\mathrm{Ker}(ad_{t_+}) + \mathrm{Ker}(ad_{t_-})\Big) \oplus \Big(\mathrm{Im}(ad_{t_+}) \cap \mathrm{Im}(ad_{t_-})\Big) \tag{3.1.25}$$

where

$$\Big(\mathrm{Ker}(ad_{t_+}) + \mathrm{Ker}(ad_{t_-})\Big) \perp \Big(\mathrm{Im}(ad_{t_+}) \cap \mathrm{Im}(ad_{t_-})\Big) \tag{3.1.26}$$

Also one has the following decomposition

$$\mathrm{Ker}(ad_{t_+}) + \mathrm{Ker}(ad_{t_-}) = C(i) \bigoplus_{k>0} \mathrm{Ker}(ad_{t_+})^{(k)} \bigoplus_{k>0} \mathrm{Ker}(ad_{t_-})^{(-k)} \tag{3.1.27}$$

As we saw above $K^{ab}t_b \in g^{(-k)}$ iff $t_a \in g^{(k)}$. Therefore

$$\begin{array}{rcll} \mathrm{Ker}(ad_{t_\pm})^{(\pm k)} & \perp & \mathrm{Ker}(ad_{t_\pm})^{(\pm l)} & \text{for all } k,l>0 \\ \mathrm{Ker}(ad_{t_+})^{(k)} & \perp & \mathrm{Ker}(ad_{t_-})^{(-l)} & \text{for } k \neq l \\ \mathrm{Ker}(ad_{t_+})^{(k)} & \perp & C(i) & \text{for all } k>0 \\ \mathrm{Ker}(ad_{t_-})^{(-k)} & \perp & C(i) & \text{for all } k>0 \end{array}$$

from which the lemma follows .

Note that from the proof of this lemma follows that the spaces $\mathrm{Ker}(ad_{t_+})^{(k)}$ and $\mathrm{Ker}(ad_{t_-})^{(-k)}$ are nondegenerately paired. The same is true for the centralizer $C(i)$ with itself. We shall use these results in the next section when we start reduction of the Kirillov Poisson structure.

3.1.3 Reductions associated to sl_2 embeddings

We are now set up to describe the Poisson reductions that lead to finite W algebras. The procedure is to impose certain constraints, or in other words to restrict oneself to a certain submanifold of a simple Lie algebra g(= M), then to construct the reduced phase space $\bar{M}$ and finally to calculate the Poisson structure on this reduced phase space. This Poisson structure then corresponds to a finite W algebra. The different sets of constraints that lead to consistent reductions are determined by the sl_2 embeddings. Let us now describe how precisely this works.

Let there be given a certain sl_2 subalgebra $\{t_0, t_+, t_-\}$ of a simple Lie algebra g. Under the adjoint action of this sl_2 subalgebra g branches into a direct sum of irreducible sl_2 multiplets. Let $\{t^{(\mu)}_{j,m}\}$ be the basis of g introduced in the previous section. Associated with this basis is a set of C^∞ functions $\{J^{j,m}_{(\mu)}\}$ on g (the dual basis) with the property $J^{j,m}_{(\mu)}(t^{(\mu')}_{j',m'}) = \delta^{\mu'}_{\mu}\delta^{j}_{j'}\delta^{m}_{m'}$. These can be called the (global) coordinate functions on g in the basis $\{t^{(\mu)}_{j,m}\}$ because they associate to an element $\alpha \in g$ its (j,m,μ) component.

Let's now impose the following set of constraints

$$\{\phi^{j,m}_{(\mu)} \equiv J^{j,m}_{(\mu)} - \delta^{j}_{1}\delta^{m}_{1}\delta^{1}_{\mu}\}_{j\in\frac{1}{2}\mathbf{N}\,;\,m>0} \tag{3.1.28}$$

(remember that $t^{(1)}_{1,\pm 1} \equiv t_{\pm}$; $t^{(1)}_{1,0} \equiv t_0$). Denote the 'zero set' in g of these constraints by g_c. Its elements have the form

$$\alpha = t_+ + \sum_j \sum_{m \leq 0} \sum_{\mu} \alpha^{j,m}_{(\mu)} t^{(\mu)}_{j,m} \tag{3.1.29}$$

where $\alpha^{j,m}_{(\mu)}$ are real or complex numbers (depending on which case we want to consider).

As we shall see in the next chapter the constraints postulated above are motivated in the infinite dimensional case by the requirement that the Poisson algebra which we obtain after reduction must be a W algebra [7, 67]. In principle, from a mathematical point of view, one could consider more general sets of constraints, however these will, if consistent, in general not lead to W algebras. This is the reason why we define finite W algebras only to be the reductions associated to sl_2 embeddings. In principle other reductions could also be handled however by the formalism that we will develop in this thesis.

As discussed in the previous chapter we need to find out which of the constraints (3.1.28) are first class for these will generate gauge invariances on g_c. This is the subject of the next lemma.

Lemma 3 *The constraints $\{\phi^{j,m}_{(\mu)}\}_{m \geq 1}$ are first class.*

Proof: First we show that

$$\{J^{j,m}_{(\mu)}, J^{j',m'}_{(\mu')}\} = \sum_{j'',\mu''} \alpha^{\mu''}_{j''} J^{j'',m+m'}_{(\mu'')} \tag{3.1.30}$$

for some coefficients $\alpha^{(\mu)}_j$. Let $x \in g^{(m'')}$, then

$$\{J^{j,m}_{(\mu)}, J^{j',m'}_{(\mu')}\}(x) = (x, [\mathrm{grad}_x J^{j,m}_{(\mu)}, \mathrm{grad}_x J^{j',m'}_{(\mu')}]) \tag{3.1.31}$$

Now let y be an element of $g^{(k)}$, then

$$(y, \mathrm{grad}_x J^{j,m}_{(\mu)}) = \frac{d}{d\epsilon} J^{j,m}_{(\mu)}(x + \epsilon y)|_{\epsilon=0} \tag{3.1.32}$$

which is zero except when $k = m$ (see definition of $J^{j,m}_{(\mu)}$). From this and the lemmas of the previous section one concludes that $\mathrm{grad}_x J^{j,m}_{(\mu)} \in g^{(-m)}$ and that the Poisson bracket (3.1.31) is nonzero if and only if $m'' = m + m'$. From this eq.(3.1.30) follows.

Consider now the Poisson bracket between two constraints in the set $\{\phi^{j,m}_{(\mu)}\}_{m \geq 1}$

$$\begin{aligned} \{\phi^{j,m}_{(\mu)}, \phi^{j',m'}_{(\mu')}\} &= \{J^{j,m}_{(\mu)}, J^{j',m'}_{(\mu')}\} \\ &= \sum_{j'',\mu''} \alpha_{j'',\mu''} J^{j'',m+m'}_{(\mu'')} \\ &= \sum_{j'',\mu''} \alpha_{j'',\mu''} \phi^{j'',m+m'}_{(\mu'')} \end{aligned}$$

which is obviously equal to zero on g_c Note that the fact $m, m' \geq 1$ was used in the last equality sign. This proves the lemma .

Note that in general the set $\{\phi^{j,m}_{(\mu)}\}_{m\geq 1}$ is not equal to the total set of constraints since the constraints with $m = \frac{1}{2}$ are not included. These constraints will turn out to be second class.

Let us now determine the group of gauge transformations on g_c generated by the first class constraints. Again we use the multi-index notation where now Roman letters $a, b, \ldots$ run over all j, m and μ, and Greek letters $\alpha, \beta \ldots$ over all j, μ but only $m > 0$ (those are the indices associated with the constraints). Let ϕ^α be one of the first class constraints (i.e. $\alpha \equiv (j, m; \mu)$ with $m \geq 1$), then the gauge transformations associated to it are generated by its Hamiltonian vectorfields

$$X^\alpha \equiv \{\phi^\alpha, \cdot\} \tag{3.1.33}$$

Let $x = x^a t_a \in g$, then the change of x under a gauge transformation generated by ϕ^α is given by

$$\begin{aligned} \delta_\alpha x &= \epsilon\{\phi^\alpha, J^a\}(x)t_a \\ &= \epsilon\{J^\alpha, J^a\}(x)t_a \\ &= \epsilon g^{\alpha a} f^c_{ab} x^b t_c \\ &= [\epsilon g^{\alpha a} t_a, x] \end{aligned} \tag{3.1.34}$$

Since $g^{\alpha a} t_a \in g^{(-k)}$ iff $t_\alpha \in g^{(k)}$ (see lemmas in the previous section) we find that the Lie algebra of the group of gauge transformations is given by

$$h = \bigoplus_{k\geq 1} g^{(-k)} \tag{3.1.35}$$

This is obviously a nilpotent Lie subalgebra of g and can be exponentiated to a group H. This is the gauge group generated by the first class constraints. Note that from eq.(3.1.34) follows that H acts on g_c in the adjoint representation, i.e the gauge orbit of a point $x \in g_c$ is given by

$$\mathcal{O} = \{gxg^{-1} \mid g \in H\} \tag{3.1.36}$$

Now that we have identified the gauge group we can come to the matter of constructing the space g_c/H, or equivalently, gauge fixing. Of course this can be done in many ways, however in [7, 67] it was argued that there are certain gauges which are the most convenient from the conformal field theoretic point of view. These are the so called 'lowest weight gauges'. Define the subset g_{fix} of g_c as follows

$$g_{fix} = \{t_+ + \sum_{j,\mu} x^j_{(\mu)} t^{(\mu)}_{j,-j} \mid x^j_{(\mu)} \in \mathbb{C}\} \tag{3.1.37}$$

We then have the following

Theorem 1

$$H \times g_{fix} \simeq g_c \tag{3.1.38}$$

Proof: Note first the obvious fact that $g^{(-l)} = g^{(-l)}_{lw} \oplus g^{(-l)}_0$, where $g^{(-l+1)}_0 = [t_+, g^{(-l)}]$ and $[t_-, g^{(-l)}_{lw}] = 0$. Let now $x \in g_c$ which means that $x = t_+ + x^{(0)} + \ldots + x^{(-p)}$ where $x^{(-k)} \in g^{(-k)}$ and $p \in \frac{1}{2}\mathbf{N}$ is the largest j value in the decomposition of the adjoint representation w.r.t. the embedding i. Of course each $x^{(-k)}$ can be written as a sum $x^{(-k)}_0 + x^{(-k)}_{lw}$ where $x^{(-k)} \in g^{(-k)}_0$ and $x^{(-k)} \in g^{(-k)}_{lw}$. Let also $\alpha^{(-k)}$ be an element of grade $-k$ in h, i.e. $\alpha^{(-k)} \in g^{(-k)} \subset h$. Then

$$\begin{aligned}
\left(e^{\alpha^{(-k)}} x e^{-\alpha^{(-k)}}\right)^{(-k+1)} &= (x^{(-k+1)}_0 + x^{(-k+1)}_{lw}) - ad_{t_+}(\alpha^{(-k)}) \\
\left(e^{\alpha^{(-k)}} x e^{-\alpha^{(-k)}}\right)^{(-k+i)} &= x^{(-k+i)} \qquad \text{for } 1 < i \leq k
\end{aligned}$$

i.e. only the elements $x^{(-k+1)}, \ldots, x^{(-p)}$ are changed by this gauge transformation. Now, since $ad_{t_+} : g^{(-k)} \to g^{(-k+1)}_0$ is bijective by definition, there exists a unique $\alpha^{(-k)} \in g^{(-k)}$ such that $ad_{t_+}(\alpha^{(-k)}) = x^{(-k+1)}_0$, i.e. the element $x^{(-k+1)}_0$ can be gauged away by choosing $\alpha^{(-k)}$ appropriately. From this follows immediately that there exist unique elements $\alpha^{(-l)} \in g^{(-l)}$ $(l = 1, \ldots, p)$ such that

$$e^{\alpha^{(-p)}} \ldots e^{\alpha^{(-1)}} x\, e^{-\alpha^{(-1)}} \ldots e^{-\alpha^{(-p)}} = y \quad \in g_{fix} \tag{3.1.39}$$

This provides, as one can now easily see, a bijective map between $H \times g_{fix}$ and g_c given by

$$\left(e^{-\alpha^{(-1)}} \ldots e^{-\alpha^{(-p)}}, y\right) \to x \tag{3.1.40}$$

This proves the theorem .

So starting from g, after imposing constraints and fixing gauge invariances we have arrived at a submanifold g_{fix} of g. We now want to determine the Poisson algebra structure of $C^\infty(g_{fix})$. For this we need to calculate the Dirac brackets

$$\{J^{j,-j}_{(\mu)}, J^{j',-j'}_{(\mu')}\}^* \tag{3.1.41}$$

between the generators $\{J^{j,-j}_{(\mu)}\}$ of $C^\infty(g_{fix})$. of course one could do this directly by using the formula for the Dirac bracket, but this is rather cumbersome. We shall therefore use another method which in the end will lead to a very elegant and general formula for the Poisson structure on $C^\infty(g_{fix})$. Unfortunately for this it will be necessary to go through some rather technical lemmas and theorems that lay the mathematical groundwork for this method. The reader may wish to skip right to formula (3.1.72) on first reading.

Consider the following general situation: Let $\{t_i\}_{i=1}^{dim(g)}$ be a basis of the Lie algebra g, k be a positive integer smaller or equal to $\dim(g)$ and $\Lambda \equiv t_{k+1}$. Consider then the following subset of g

$$g_f = \{\Lambda + \sum_{i=1}^{k} \alpha^i t_i \mid \alpha^i \in \mathbb{C}\} \tag{3.1.42}$$

which can be seen as the zero set of the constraints $\phi^1 = J^{k+1} - 1$ and $\phi^i = J^{k+i}$ for $1 < i \leq \dim(g) - k$. Also suppose that the Kirillov bracket on g induces a Dirac bracket $\{.,.\}^*$ on g_f (i.e. all constraints are second class).

Denote by $\mathcal{R}$ the set of smooth functions

$$R : \mathbb{C}^k \times g_f \longrightarrow g \tag{3.1.43}$$

of the form

$$R(\vec{z}; y) = \sum_{i=1}^{k} z_i R^i(y) \tag{3.1.44}$$

where $\vec{z} \equiv \{z_i\}_{i=1}^{k} \in \mathbb{C}^k$, $y \in g_f$ and $R^i(y) \in g$. To any element $R \in \mathcal{R}$ one can associate a map

$$Q_R : \mathbb{C}^k \times g_f \longrightarrow C^\infty(g) \tag{3.1.45}$$

defined by

$$Q_R(\vec{z}; y) = \sum_{i=1}^{dim(g)} (R(\vec{z}; y), t_i) J^i \tag{3.1.46}$$

(where as before $J^i \in C^\infty(g)$ is defined by $J^i(t_j) = \delta^i_j$). We are going to use certain elements of the set $\mathcal{R}$ in order to explicitly calculate the Dirac brackets on g_f. We have the following theorem.

Theorem 2 *If there exists a unique $R \in \mathcal{R}$ such that for all $\vec{z} \in \mathbb{C}^k$ and $y \in g_f$ we have*

$$\Lambda + [R(\vec{z}; y), y] \quad \in g_f \tag{3.1.47}$$

and

$$Q_R(\vec{z}; y)|_{g_f} = \sum_{i=1}^{k} z_i J^i + (R(\vec{z}; y), \Lambda) \tag{3.1.48}$$

(i.e. the restriction of $Q_R(\vec{z}; y)$ to g_f is equal to the right hand side of (3.1.48)) then

$$\sum_{ij=1}^{k} z_i \, \{J^i, J^j\}^*(y) \, t_j = [R(\vec{z}; y), y] \tag{3.1.49}$$

for all $\vec{z} \in \mathbb{C}^k$ and $y \in g_f$.

Note that from eq.(3.1.49) one can read off *all* Dirac brackets between the generators $\{J^i\}_{i=1}^k$ of $C^\infty(g_f)$ since the formula holds for all $\vec{z} \in \mathbb{C}^k$ and the elements t_j are all independent. From this follows that calculating the whole set of Dirac brackets has been brought down to determining the map R. This is, in practical situations, easily done using eqs.(3.1.47) and (3.1.48).

Also note that from equation (3.1.48) follows that within the Dirac bracket $\{.,.\}^*$ the function $Q_R(\vec{z};y)$ is equal to $\sum_{i=1}^k z_i J^i$, i.e.

$$\{Q_R(\vec{z};y),\cdot\}^* = \sum_{i=1}^k z_i \{J^i,\cdot\}^* \tag{3.1.50}$$

since constants commute with everything and restriction to g_f is always implied within the Dirac bracket. This explains the origin of the formula (3.1.48).

The proof of the theorem is based on the following two lemmas.

Lemma 4 *Let M be a Poisson manifold, $Q \in C^\infty(M)$ and X_Q its Hamiltonian vector-field, i.e. $X_Q = \{Q,.\}$. Let there also be given a set of second class constraints and let Z be the submanifold where they all vanish. If for some $p \in M$ we have $X_Q|_p(\phi_i) = 0$ for all i (i.e. $X_Q|_p$ is tangent to Z), then*

$$\{Q,F\}(p) = \{\bar{Q},\bar{F}\}^*(p) \quad \textit{for all } F \in C^\infty(M) \tag{3.1.51}$$

where $\{.,.\}^*$ is the Dirac bracket.

Proof: Simply calculating the Dirac bracket in p gives

$$\begin{aligned} \{\bar{Q},\bar{F}\}^*(p) &\equiv \{Q,F\}(p) - \{Q,\phi_i\}(p)\Delta^{ij}(p)\{\phi_j,F\}(p) \\ &= \{Q,F\}(p) \end{aligned}$$

because by hypothesis $\{Q,\phi_i\}(p) = 0$ for all i. This proves the lemma .

The origin of equation (3.1.47) in the theorem becomes clear in the following lemma.

Lemma 5 *If for all $y \in g_f$ and $\vec{z} \in \mathbb{C}^k$*

$$\Lambda + [R(\vec{z};y), y] \quad \in g_f \tag{3.1.52}$$

then

$$\{Q_R(\vec{z};y),\phi^i\}(y) = 0 \tag{3.1.53}$$

for all $1 \le i \le p-k$; $y \in g_f$ and $\vec{z} \in \mathbb{C}^k$.

Proof: By definition of J^i we have $y = t_i J^i(y)$ which gives us

$$\begin{aligned}
\Lambda + [R(\vec{z};y),y] &= \Lambda + \sum_{ij=1}^{dim(g)} [R(\vec{z};y)^i t_i, t_j J^j](y) \\
&= \Lambda + \sum_{ijlm=1}^{dim(g)} K_{ij} f_m^{jl} R(\vec{z};y)^i J^m(y) t_l \\
&= \Lambda + \sum_{ijl=1}^{dim(g)} \{K_{ij} R(\vec{z};y)^i J^j, J^l\}(y) t_l \\
&\equiv \Lambda + \sum_{i=1}^{dim(g)} \{Q_R(\vec{z};y), J^i\}(y) t_i
\end{aligned}$$

but by hypothesis this was an element of g_f. Therefore $\{Q_R(\vec{z};y), J^i\}(y) = 0$ for $i \geq k+1$. This proves the lemma .

Putting together these two lemmas we find

Proof of theorem: From the previous lemmas follows that if eqns. (3.1.47) and (3.1.48) are satisfied, then

$$\begin{aligned}
\sum_{ij=1}^{k} z_i \{J^i, J^j\}^*(y) t_j &= \sum_{i=1}^{k} \{Q_R(\vec{z};y), J^i\}^*(y) t_i \\
&= \sum_{i=1}^{k} \{Q_R(\vec{z};y), J^i\}(y) t_i \\
&= \sum_{i=1}^{dim(g)} \{Q_R(\vec{z};y), J^i\}(y) t_i \\
&= [R(\vec{z};y), y] \qquad (3.1.54)
\end{aligned}$$

where eqn.(3.1.48) was used in the first step. This proves the theorem .

Note that conversely in order to show the existence of the Dirac bracket on g_f it is sufficient to prove that equations (3.1.47) and (3.1.48) of the above theorem are solvable within $\mathcal{R}$. We will now show that this is the case when $g_f \equiv g_{fix}$ associated to an arbitrary sl_2 embedding.

Theorem 3 *Let $|i|$ be the total number of sl_2 multiplets in the branching of the adjoint representation of g. There is a unique $R \in \mathcal{R}$ such that*

$$t_+ + [R(\vec{z};y), y] \quad \in g_{fix} \qquad (3.1.55)$$

and

$$Q_R(\vec{z};y)|_{g_{fix}} = \sum_{j;\mu} z_j^{(\mu)} J_{(\mu)}^{j,-j} + (R(\vec{z};y), t_+) \tag{3.1.56}$$

for all $\vec{z} \in \mathbb{C}$ *and* $y \in g_{fix}$.

Proof: First solve the second equation.

$$\begin{aligned} Q_R(\vec{z};y)(y) &= \left(R^{(p)} + \ldots + R^{(-p)}, t_+ + y^{(0)} + \ldots + y^{(-p)}\right) \\ &= (R^{(-1)}, t_+) + (R^{(p)}, y^{(-p)}) + \ldots + (R^{(0)}, y^{(0)}) \end{aligned}$$

Now, $y^{(-j)} \in g_{lw}^{(-j)}$ which means that $y^{(-j)} = \sum_\mu y_{(\mu)}^{j,-j} t_{j,-j}^{(\mu)}$. Also $R^{(j)} = R_0^{(j)} + R_{hw}^{(j)}$ where $R_0^{(j)} \in \mathrm{Im}(ad_{t_+})$ and $R_{hw}^{(j)} \in \mathrm{Ker}(ad_{t_+})$. Let $\{t_{(\mu)}^{j,-j}\}_\mu \subset g^{(j)}$ be such that $(t_{j,-j}^{(\mu)}, t_{(\mu')}^{j',-j'}) = \delta_j^{j'} \delta_{\mu'}^{\mu}$ (remember that $g^{(j)}$ and $g^{(-j)}$ are non-degenerately paired). Then take

$$R_{hw}^{(j)} = \sum_\mu z_j^{(\mu)} t_{(\mu)}^{j,-j} \tag{3.1.57}$$

Obviously the second equation of the theorem is then satisfied.

Let's now solve the first equation. For $k > 0$ it reads $[R,y]^{(k)} = 0$. Writing this out gives

$$[R_0^{(k-1)}, t_+] = [y^{(0)}, R^{(k)}] + \ldots + [y^{(k-p)}, R^{(p)}] \tag{3.1.58}$$

Remember that $ad_{t_+} : g_0^{(k-1)} \to g^{(k)}$ is bijective which means that if we already know $R^{(k)}, \ldots, R^{(p)}$ then we can uniquely solve this equation for $R_0^{(k-1)}$ (remember that for $l \geq 0$ we have already determined the quantities $R_{hw}^{(l)}$). The initial term of this sequence of equations is obviously $R^{(-p)} = R_{hw}^{(-p)}$ (i.e. $R_0^{(-p)} = 0$). In this way we can completely determine the positive and zero grade piece of $R(\vec{z};y)$.

For $-k \geq 0$ the situation is similar but slightly different. What we now have to demand is that $[R,y]_0^{(-k)} = 0$. This leads to the equation

$$[t_+, R^{(-k-1)}] = [R^{(-k)}, y^{(0)}]_0 + \ldots + [R^{(-k+p)}, y^{(-p)}]_0 \tag{3.1.59}$$

where we used that $[t_+, R^{(-k-1)}]$ is already an element of $g_0^{(-k)}$. Remember that $ad_{t_+} : g^{(-k-1)} \to g_0^{(-k)}$ is bijective so there exists a unique $R^{(-k-1)}(z)$ such that the equation is solved, assuming we know $R^{(-k)}, \ldots, R^{(-k+p)}$ already. In this way we can grade by grade determine the negative grade piece of the matrix R. It is also clear from the above construction that $R \in \mathcal{R}$. This proves the theorem .

From theorem 2 therefore follows that

$$\sum_{j,j';\mu,\mu'} z_j^{(\mu)} \{J_{(\mu)}^{j,-j}, J_{(\mu')}^{j',-j'}\}^*(y)\, t_{j',-j'}^{(\mu')} = [R(\vec{z};y), y] \tag{3.1.60}$$

from which we can derive all relations.

It is possible to derive a general formula for $R(\vec{z};y)$ using arguments similar to those used in [69]. Let again

$$g_{lw} = \bigoplus_{j\in\frac{1}{2}\mathbf{N}} g_{lw}^{(-j)} = \mathrm{span}(\{t_{j,-j}^{(\mu)}\}_{j;\mu} \tag{3.1.61}$$

and let Π be the orthogonal (w.r.t. the Cartan-Killing form) projection onto $\mathrm{Im}(ad_{t_+})$. Obviously the map $ad_{t_+} : \mathrm{Im}(ad_{t_-}) \to \mathrm{Im}(ad_{t_+})$ is invertible. Denote the inverse of this map, extended by 0 to g by L. As before what we want to do is solve the equation

$$[R,y] = x \ \in g_{lw} \tag{3.1.62}$$

for $y \in g_{fix}$. Noting that $y = t_+ + w$ where $w \in g_{lw}$ and applying Π this equation reduces to

$$\Pi \circ ad_{t_+} R(\epsilon) = \epsilon\Pi([R(\epsilon), w]) \tag{3.1.63}$$

where we introduced a parameter ϵ in the right hand side which we want to put to 1 later. Note that the left hand side is equal to $ad_{t_+}R(\epsilon)$ since this is already an element of $\mathrm{Im}(ad_{t_+})$. Assume now that $R(\epsilon)$ can be (perturbatively) written as

$$R(\epsilon) = \sum_{k=0}^{\infty} R_k \epsilon^k \tag{3.1.64}$$

(we shall have to justify this later). Consider now the zeroth order part of the equation (3.1.63). It reads

$$ad_{t_+} R_0 = 0 \tag{3.1.65}$$

This means that $R_0 \equiv F(\vec{z})$ which is an arbitrary element of $\ker(ad_{t_+})$. The first order equation is equal to

$$ad_{t_+} R_1 = \Pi([F,w]) \tag{3.1.66}$$

Obviously this equation is solved by

$$R_1 = -L \circ ad_w F \tag{3.1.67}$$

Proceeding with higher orders we find

$$R_{k+1}(\vec{z};y) = -L \circ ad_w \left(R_k(\vec{z};y)\right) \tag{3.1.68}$$

which means that

$$R(\vec{z};y) = \frac{1}{1+\epsilon L \circ ad_w} F(\vec{z}) \tag{3.1.69}$$

There are no convergence problems with this series since the operator L lowers the degree by one which means that after $2p$ steps it must cancel. We therefore find

$$\sum_{j,j';\mu\mu'} z_j^{(\mu)} \{J_{(\mu)}^{j,-j}, J_{(\mu')}^{j',-j'}\}^*(y)\, t_{j',-j'}^{(\mu')} = -ad_y \left(\frac{1}{1+Lad_w} F(\vec{z})\right) \tag{3.1.70}$$

Now, let $w \in g_{lw}$ and $Q \in C^\infty(g_{lw})$. We then define $\mathrm{grad}_w Q \in \mathrm{Ker}(ad_{t_+}) \equiv g_{hw}$ by

$$(x, \mathrm{grad}_w Q) \equiv \frac{d}{d\epsilon} Q(w+\epsilon x)|_{\epsilon=0} \quad \text{for all } x \in g_{lw} \tag{3.1.71}$$

Note that this uniquely defines $\mathrm{grad}_w Q$ since, as we saw before, g_{lw} and g_{hw} are non-degenerately paired. Using equation (3.1.70) we can now give the following general description of the classical finite W algebra: it is nothing but the Poisson algebra $(C^\infty(g_{lw}), \{.,.\}^*)$ where

$$\{Q_1, Q_2\}^*(w) = \left(w, [\mathrm{grad}_w Q_1, \frac{1}{1+L\circ ad_w} \mathrm{grad}_w Q_2]\right) \tag{3.1.72}$$

for all $Q_1, Q_2 \in C^\infty(g_{lw})$ and $w \in g_{lw}$. For the so called trivial embedding ($t_0 = t_\pm = 0$) the map L is equal to the zero map and the above formula reduces to the ordinary Kirillov bracket as it should (because in that case $g_{fix} \equiv g$). For nontrivial sl_2 embeddings however (3.1.72) is a new and highly nontrivial Poisson structure.

From now on denote the Kirillov Poisson algebra associated to a Lie algebra g by $K(g)$ and the finite W algebra $(C^\infty(g_{lw}), \{.,.\}^*)$ by $\mathcal{W}(i)$ where i is again the sl_2 embedding in question.

In general $\mathcal{W}(i)$ is a non-linear Poisson algebra as we shall see when we consider examples. However it can happen that $\mathcal{W}(i)$ contains a subalgebra that is isomorphic to a Kirillov algebra. The next theorem deals with this.

Theorem 4 *The algebra $\mathcal{W}(i)$ has a Poisson subalgebra which is isomorphic to $K(C)$ where C is the centralizer of the sl_2 subalgebra of g.*

Proof: Since all elements of the centralizer are lowest (and highest) weight vectors w.r.t. $i(sl_2)$ the function J^a is not constrained if $t_a \in C(i)$, i.e. all the elements J^a associated to the centralizer survive the reduction. It is not obvious however that they still form the same algebra w.r.t. the Poisson bracket (3.1.72). This is what we have to show.

The part of the equation (3.1.60) that determines the Poisson relations between the currents associated to the centralizer is

$$\sum_{\mu,\mu'} z_0^{(\mu)} \{J_{(\mu)}^{0,0}, J_{(\mu')}^{0,0}\}^*(y)\, t_{0,0}^{(\mu')} = [R(\vec{z};y), y]^{(0)} \tag{3.1.73}$$

The right hand side of this equation reads in more detail

$$[R,y]^{(0)} = [R^{(0)}_{lw}, y^{(0)}] + [R^{(0)}_0, y^{(0)}] + \ldots + [R^{(p)}, y^{(-p)}] + [R^{(-1)}, t_+] \tag{3.1.74}$$

Note that $R^{(0)}_{lw}$ and $y^{(0)}$ are both elements of $C(i)$ which means that the first term in the right hand side is also an element of $C(i)$. We will now show that $R^{(0)}_0, R^{(1)}, \ldots, R^{(p)}$ and $R^{(-1)}$ do not depend on $\{z^{(\mu)}_0\}_\mu$ which means that all but the first term in the right hand side of equation (3.1.74) are irrelevant for the Poisson brackets $\{J^{0,0}_{(\mu)}, J^{0,0}_{(\mu')}\}^*$.

From eq.(3.1.58) follows that $R^{(k)}_0$ is only a function of $z^{(\mu)}_p, \ldots, z^{(\mu)}_{k+1}$ for $k \geq 0$ and therefore $R^{(k)} = R^{(k)}(z^{(\mu)}_p, \ldots, z^{(\mu)}_k)$. Now, $R^{(-1)}$ is determined by equation (3.1.59) for $k = 0$

$$[t_+, R^{(-1)}] = [R^{(0)}, y^{(0)}]_0 + \ldots + [R^{(p)}, y^{(-p)}]_0 \tag{3.1.75}$$

Note that the terms $[R^{(l)}, y^{(-l)}]$ for $l > 0$ certainly do not contain $z^{(\mu)}_0$ as we just saw. Also note that

$$[R^{(0)}, y^{(0)}]_0 = [R^{(0)}_0 + R^{(0)}_{lw}, y^{(0)}]_0 = [R^{(0)}_0, y^{(0)}] \tag{3.1.76}$$

because of the reason we mentioned earlier that $R^{(0)}_{hw}$ and $y^{(0)}$ are both in $C(i)$. However, as we have seen above $R^{(0)}_0$ does not contain $z^{(\mu)}_0$. From this we conclude that the only term in $[R,y]^{(0)}$ that contains $z^{(\mu)}_0$ is $[R^{(0)}_{lw}, y^{(0)}]$, i.e.

$$\begin{aligned} \sum_{\mu\mu'} z^{(\mu)}_0 \{J^{0,0}_{(\mu)}, J^{0,0}_{(\mu')}\}^*(y) t^{(\mu)}_{0,0} &= [R^{(0)}_{lw}, y^{(0)}] \\ &= [R^{(0)}_{lw}, \sum_\mu J^{0,0}_{(\mu)}(y) t^{(\mu)}_{0,0}] \end{aligned} \tag{3.1.77}$$

The generators $\{t^{(\mu)}_{0,0}\}$ are a basis of $C(i)$. From this, eq.(3.1.77) and theorem 2 follows immediately that the Poisson algebra generated by $\{J^{0,0}_{(\mu)}\}$ w.r.t. the Dirac bracket is isomorphic to Kirillov algebra of $C(i)$. This proves the theorem .

3.1.4 Generalized finite Miura transformations

In this section we present a generalized version of the Miura transformation. Roughly this is a Poisson homomorphism of the finite W algebra $\mathcal{W}(i)$ in question to a certain Kirillov algebra. In order to be able to describe this map for arbitrary embeddings however we first have to concern ourselves with the cases when in the decomposition of g into sl_2 multiplets there appear half integer grades. As we have seen, in those cases the constraints $\phi^{j,\frac{1}{2}}_{(\mu)}$ are second class. In what follows it will be necessary to be able to replace the usual set of constraints by an alternative set which contains only first class constraints but which gives rise to the same reduction [80]. Roughly what one does is

impose only half of the constraints that turned out to be second class in such a way that they become first class. The other constraints that were second class can then be obtained as gauge fixing conditions. In this way g_{fix} stays the same but g_c is different. Since the resulting Poisson algebra only depends on g_{fix} it is clear that we obtain the same algebra $\mathcal{W}(i)$.

Let's now make all of this more precise. We describe the sl_n algebra in the standard way by traceless $n \times n$ matrices; E_{ij} denotes the matrix with a one in its (i,j) entry and zeroes everywhere else. As we said earlier embeddings of sl_2 into sl_n are in one-to-one correspondence with partitions of n. Let $(n_1, n_2, \ldots)$ be a partition of n, with $n_1 \geq n_2 \geq \ldots$. Define a different partition $(m_1, m_2, \ldots)$ of n, with m_k equal to the number of i for which $n_i \geq k$. Furthermore let $s_t = \sum_{i=1}^{t} m_i$. Then we have the following

Lemma 1 *An embedding of sl_2 in sl_n under which the fundamental representation branches according to $n \to \oplus n_i$ is given by*

$$\begin{aligned} t_+ &= \sum_{l\geq 1} \sum_{k=1}^{n_l - 1} E_{l+s_{k-1}, l+s_k}, \\ t_0 &= \sum_{l\geq 1} \sum_{k=1}^{n_l} (\frac{n_l+1}{2} - k) E_{l+s_{k-1}, l+s_{k-1}}, \\ t_- &= \sum_{l\geq 1} \sum_{k=1}^{n_l - 1} k(n_l - k) E_{l+s_k, l+s_{k-1}}. \end{aligned} \tag{3.1.78}$$

The proof is by direct computation*. If the fundamental representation of sl_n is spanned by vectors $v_1, \ldots v_n$, on which sl_n acts via $E_{ij}(v_k) = \delta_{jk} v_i$, then the irreducible representations n_l of sl_2 to which it reduces are spanned by $\{v_{l+s_{k-1}}\}_{1 \leq k \leq n_l}$. Previously we decomposed the lie algebra g into eigenspaces of ad_{t_0}. However, for our present purposes it is convenient to introduce a different grading of the Lie algebra g, that we need to describe the generalized Miura map and the BRST quantization. The grading is defined by the following element of the Cartan subalgebra of sl_n:

$$\delta = \sum_{k\geq 1} \sum_{j=1}^{m_k} \left(\frac{\sum_l l m_l}{\sum_l m_l} - k \right) E_{s_{k-1}+j, s_{k-1}+j}. \tag{3.1.79}$$

This leads to the alternative decomposition $g = g_- \oplus g_0 \oplus g_+$ of g into spaces with negative, zero and positive eigenvalues under the adjoint action of δ respectively (note that in case the grading of ad_{t_0} is an integral grading then we have $t_0 = \delta$ so in those

*The commutation relations are $[t_0, t_\pm] = \pm t_\pm$, and $[t_+, t_-] = 2t_0$.

cases nothing happens. In general however $t_0 \neq \delta$ and also $g^{(m)} \neq g_m$). The subalgebra g_0 consists of matrices whose nonzero entries are in square blocks of dimensions $m_1 \times m_1$, $m_2 \times m_2$, etc. along the diagonal of the matrix and is therefore a direct sum of sl_{m_k} subalgebras (modulo $u(1)$ terms). The nilpotent subalgebra g_+ is spanned by $\{E_{l+s_{k-1},r+s_k}\}_{l\geq 1;\, 1\leq k\leq n_l-1;\, r>0}$, and the nilpotent subalgebra g_- by the transpose of these. Let $\pi_\pm$ denote the projections onto $g_\pm$. Then the following theorem describes the replacement of the mixed system of first and second class constraints by a system of first class constraints only.

Theorem 1 *The constraints* $\{J^{l+s_{k-1},r+s_k} - \delta^{r,l}\}_{l\geq 1;\, 1\leq k\leq n_l-1;\, r>0}$ *are first class. The gauge group they generate is* $\hat{H} = \exp(g_-)$ *acting via the adjoint representation on* g*. The resulting finite* W *algebra is the same as the one obtained by imposing the constraints (3.1.28).*

Proof: decompose g in eigenvalues of ad_δ, $g = \oplus_k g_k$. Note that ad_δ has only integral eigenvalues. Using the explicit form of t_+ in (3.1.78), one easily verifies that $[\delta, t_+] = t_+$. Thus, $t_+ \in g_1$. Again since $[g_+, g_+] = [g_{\geq 1}, g_{\geq 1}] \subset g_{\geq 2}$ it follows (exactly like in lemma 3) that the constraints $\{J^{l+s_{k-1},r+s_k} - \delta^{r,l}\}_{l\geq 1;\, 1\leq k\leq n_l-1;\, r>0}$ are first class. The gauge group can be determined similarly as in section 3.2, and the analogue of theorem 1 of section 3.2 can be proven in the same way with the same choice of g_{fix}, if one uses the decomposition of g in eigenspaces of ad_δ rather than ad_{t_0}. Therefore the resulting finite W algebra is the same, because theorem 2 and 3 of section 3.2 show that it only depends on the form of g_{fix}.

Let us explain this theorem in words. The number of second class constraints in any system is necessarily even. If one switches to the ad_δ grading the set of second class constraints is split into half: one half gets grade 1 w.r.t. ad_δ while the other half gets grade 0. Now what one does is impose only that half that has obtained grade 1 w.r.t. the ad_δ grading. These constraints are then first class. The gauge transformations they generate can be completely fixed by imposing the constraints that were in the other half. Having done that we are back in exactly the same situation as before. The only difference is that we now know of a system of first class constraints that in the end leads to the same reduction.

Note that the number of generators of the finite W algebra is equal to $\dim(g_0) = (\sum_i m_i^2) - 1$. This is indeed the same as the number of irreducible representations of sl_2, minus one, one obtains from $(\oplus_i n_i) \otimes (\oplus_i n_i)$, as the latter number equals $(\sum_j (2j-1)n_j) - 1$, and one easily checks that $\sum_i m_i^2 = \sum_j (2j-1)n_j$.

The generalized Miura transformation can now be formulated as

Theorem 5 *There exists an injective Poisson homomorphism from* $\mathcal{W}(i)$ *to* $K(g_0)$*.*

Proof: First we show that for every element $x \in t_+ + g_0$ there exist a unique element h in the gauge group $\hat{H}$ such that $h \cdot x \cdot h^{-1} \in g_{fix}$. For this note that there exists a unique element $h' \in \hat{H}$ such that $h'.x.h'^{-1} \in g_c$. This follows from the previous theorem and the remarks made after it. In fact it follows from those remarks that h' is an element of that subgroup of $\hat{H}$ that is generated by the 'ex' second class constraints (the ones that were made into first class constraints by not imposing the other half). It was shown earlier however that

$$g_{fix} \times H \simeq g_c \tag{3.1.80}$$

which means that there exists a unique element $h'' \in H$ such that

$$h''h'.x.(h''h')^{-1} \in g_{fix} \tag{3.1.81}$$

We conclude from this that there is a surjective map from $x \in t_+ + g_0$ to g_{fix} given by (3.1.81). The pull back of this map $C^\infty(g_{fix}) \equiv \mathcal{W}(i) \to C^\infty(t_+ + g_0)$ (which will therefore be injective) is then the Miura map. Of course we still have to check whether this map is a Poisson homomorphism. This we address next.

What is the Poisson structure on $C^\infty(t_+ + g_0)$? Since the constraints of which the space $t_+ + g_0$ is the zero set are obviously second class the Kirillov bracket on g induces a Dirac bracket on it. It is not difficult to see that the Dirac term in the Dirac bracket cancels in this case which means that the Poisson algebra $C^\infty(t_+ + g_0)$ with the induced Poisson structure is isomorphic to the Poisson algebra $K(g_0)$. Since the transformation from $\mathcal{W}(i)$ to $K(g_0)$ corresponds to a gauge transformation (3.1.81) the map is necessarily a homomorphism.

3.1.5 Examples

The simplest examples of finite W algebras are those associated to the so called 'principal sl_2 embeddings'. These embeddings are associated to the trivial partition of the number n: $n = n$. The fundamental representation of $g = sl_n$ therefore becomes an irreducible representation of the sl_2 subalgebra, i.e.

$$\underline{n}_n \simeq \underline{n}_2 \tag{3.1.82}$$

The branching rule for the adjoint representation of g therefore reads

$$\underline{ad}_n \simeq \underline{3}_2 \oplus \underline{5}_2 \oplus \ldots \oplus \underline{2n-1}_2 \tag{3.1.83}$$

From this follows immediately that the finite W algebra will have $n-1$ generators (since there are $n-1$ sl_2 multiplets). Without going into details we can immediately predict

the Poisson relations between these generators from the generalized Miura transformation for in this case the subalgebra $g^{(0)} = g_0$ coincides with the Cartan subalgebra of sl_n. Since the Cartan subalgebra is an abelian algebra, and since the Kirillov algebra of an abelian Lie algebra is a Poisson commutative algebra we find that a finite W algebra associated to a principal embedding must also be Poisson commutative since it is isomorphic to a Poisson subalgebra of $K(g_0)$. We conclude therefore that the principal sl_2 embedding into sl_n leads to the abelian Poisson algebras with $(n-1)$ generators .

The simplest nontrivial case of a finite W algebra is associated to the (only) nonprincipal embedding of sl_2 into sl_3. This embedding is associated to the following partition of 3: $3 \to 2+1$. The branching rule of the fundamental representation of sl_3 is therefore

$$\underline{3}_3 \simeq \underline{2}_2 \oplus \underline{1}_2 \tag{3.1.84}$$

From this we find the following branching rule for the adjoint representation

$$\underline{ad}_3 \simeq \underline{3}_2 \oplus 2.\underline{2}_2 \oplus \underline{1}_2 \tag{3.1.85}$$

from which follows immediately that the finite W algebra associated to this embedding will have 4 generators. We shall go through the construction of this finite W algebra in some detail in order to illustrate the theory discussed above.

The explicit form of the sl_2 embedding is

$$\begin{aligned} t_+ &= E_{1,3} \\ t_0 &= \mathrm{diag}(\frac{1}{2}, 0, -\frac{1}{2}) \\ t_- &= E_{3,1} \end{aligned} \tag{3.1.86}$$

where as before E_{ij} denotes the matrix with a one in its (i,j) entry and zeros everywhere else. The (sl_3 valued) function $J = t^{(\mu)}_{j,m} J^{j,m}_{(\mu)}$ (where we used summation convention) reads

$$J = \begin{pmatrix} \frac{1}{2}J^{1,0}_{(1)} + J^{0,0}_{(1)} & J^{\frac{1}{2},\frac{1}{2}}_{(1)} & J^{1,1}_{(1)} \\ J^{\frac{1}{2},-\frac{1}{2}}_{(2)} & -2J^{0,0}_{(1)} & J^{\frac{1}{2},\frac{1}{2}}_{(2)} \\ J^{1,-1}_{(1)} & J^{\frac{1}{2},-\frac{1}{2}}_{(1)} & J^{0,0}_{(1)} - \frac{1}{2}J^{1,0}_{(1)} \end{pmatrix} \tag{3.1.87}$$

According to the general prescription the constraints are

$$J^{1,1}_{(1)} - 1 = J^{\frac{1}{2},\frac{1}{2}}_{(1)} = J^{\frac{1}{2},\frac{1}{2}}_{(2)} = 0 \tag{3.1.88}$$

the first one being the only first class constraint. As was shown earlier the gauge invariance generated by this constraint can be completely fixed by adding the 'gauge fixing condition'

$$J_{(1)}^{1,0} = 0 \tag{3.1.89}$$

The Dirac brackets between the generators $\{J_{(1)}^{0,0}, J_{(1)}^{\frac{1}{2},-\frac{1}{2}}, J_{(2)}^{\frac{1}{2},-\frac{1}{2}}\}$ and $J_{(1)}^{1,-1}$ can now easily be calculated. In order to describe the final answer in a nice form introduce

$$\begin{aligned} C &= -\frac{4}{3}(J_{(1)}^{1,-1} + 3(J_{(1)}^{0,0})^2) \\ E &= J_{(1)}^{\frac{1}{2},-\frac{1}{2}} \\ F &= \frac{4}{3} J_{(2)}^{\frac{1}{2},-\frac{1}{2}} \\ H &= 4J_{(1)}^{0,0} \end{aligned} \tag{3.1.90}$$

(note that this is an invertible basis transformation). The Dirac bracket algebra between these generators reads [74]

$$\begin{aligned} \{H, E\}^* &= 2E \\ \{H, F\}^* &= -2F \\ \{E, F\}^* &= H^2 + C \end{aligned} \tag{3.1.91}$$

and C Poisson commutes with everything. This algebra which is called $\bar{W}_3^{(2)}$ was first constructed in [72] as a nonlinear deformation of $su(2)$. In [74] it was shown to be a reduction of sl_3 and its representation theory was explicitly constructed.

Let's now consider the finite Miura transformation for this algebra. Since the grading of sl_3 by ad_{t_0} is half integer we have to switch to the grading by ad_δ. The explicit form of δ is

$$\delta = \frac{1}{3}\mathrm{diag}(1, 1, -2) \tag{3.1.92}$$

It is easily checked that this defines an integer grading of sl_3. The crucial change is that the elements E_{23} and E_{12} have grade 1 and 0 w.r.t. the ad_δ grading while they have grade $\frac{1}{2}$ w.r.t. ad_{t_0}. According to the prescription given in the previous section the alternative set of constraints that one now imposes in order to reduce the mixed system of first and second class constraints to a system of first class constraints only is

$$J_{(1)}^{1,1} - 1 = J_{(2)}^{\frac{1}{2},\frac{1}{2}} = 0 \tag{3.1.93}$$

As we already mentioned in the previous section, what has happened here is that one has now imposed only half of the constraints that turned out to be second class. The

result is that both constraints in (3.1.93) are first class. The point however is that the gauge symmetry induced by the second constraint in eq. (3.1.93) can be completely fixed by adding the gauge fixing condition

$$J_{(1)}^{\frac{1}{2},\frac{1}{2}} = 0 \tag{3.1.94}$$

which then leaves us with exactly the same set of constraints and gauge invariances as before.

We can now describe the generalized Miura map for this case. An arbitrary element of $t_+ + g_0$ is given by

$$J_0 \equiv \begin{pmatrix} h+s & e & 1 \\ f & s-h & 0 \\ 0 & 0 & -2s \end{pmatrix} \tag{3.1.95}$$

Note that $g_0 \simeq sl_2 \oplus u(1)$ which means that the Poisson relations in $K(g_0)$ between the generators h, e, f and s (viewed as elements of $C^\infty(g_0)$) are given by

$$\begin{aligned} \{h, e\} &= e \\ \{h, f\} &= -f \\ \{e, f\} &= 2h \end{aligned} \tag{3.1.96}$$

and s commutes with everything. As shown in the previous section the equation that we have to solve in order to get explicit formulas for the Miura map is is the following equation for $h \in \hat{H}$ (where $\hat{H}$ is again the group of gauge transformations generated by the two first class constraints (3.1.93)).

$$hJ_0h^{-1} = \begin{pmatrix} J_{(1)}^{0,0} & 0 & 1 \\ J_{(2)}^{\frac{1}{2},-\frac{1}{2}} & -2J_{(1)}^{0,0} & 0 \\ J_{(1)}^{1,-1} & J_{(1)}^{\frac{1}{2},-\frac{1}{2}} & J_{(1)}^{0,0} \end{pmatrix} \tag{3.1.97}$$

The unique solution of this equation is given by

$$h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{3}{2}s + \frac{1}{2}h & e & 1 \end{pmatrix} \tag{3.1.98}$$

Inserting this back into eqn.(3.1.99) one finds certain expressions for $J_{(1)}^{0,0}, J_{(1)}^{\frac{1}{2},-\frac{1}{2}}, J_{(2)}^{\frac{1}{2},-\frac{1}{2}}$ and $J_{(1)}^{1,-1}$ in terms of the functions $\{e, f, h, s\}$. In terms of the generators (3.1.90) these

read

$$\begin{aligned} H &= 2h - 2s \\ E &= (3s - h)e \\ F &= \frac{4}{3} f \\ C &= -\frac{4}{3}(h^2 + 3s^2 + fe) \end{aligned} \tag{3.1.99}$$

It is easy to check, using the relations (3.1.96) that these satisfy the algebra (3.1.91). Therefore we find that indeed (3.1.99) provides an injective Poisson homomorphism from $W_3^{(2)}$ into the Kirillov Poisson algebra of the Lie algebra $g_0 = sl_2 \oplus u(1)$.

In the appendix we consider more examples of finite W algebras and their quantum versions. In particular we construct all finite W algebras associated to sl_4 (to get the classical versions of the algebras displayed in the appendix just replace commutators by Poisson brackets and forget about all terms of order $\hbar^2$ or higher).

3.1.6 An algebraic description of finite W algebras

Putting together the results obtained in the previous sections we can describe finite W algebras in an elegant and purely algebraic manner.

As before latin indices will be supposed to run over the entire basis t_a of the Lie algebra g, greek indices run over the basis of g_+ and barred greek indices (like $\bar{\alpha}$) run over the basis of $g_- \oplus g_0$. Indices can again be raised and lowered by use of the Cartan Killing metric.

Again let $K(g) = (C^\infty(g), \{.,.\})$ denote the Kirillov Poisson algebra associated to g. The constraints $\{\phi^\alpha\}$ are of course simply elements of $C^\infty(g)$. Denote by I the subset

$$I = \{f_\alpha \phi^\alpha \mid f_\alpha \in C^\infty(g)\} \tag{3.1.100}$$

of $C^\infty(g)$. Obviously I is an ideal w.r.t. the (abelian) multiplication map in $K(g)$ since $h f_\alpha \phi^\alpha = \tilde{f}_\alpha \phi^\alpha \in I$ for all $h \in C^\infty(g)$, where $\tilde{f}_\alpha = h f_\alpha$. In this terminology an element $f \in C^\infty(g)$ is called 'first class' if

$$\{f, \phi^\alpha\} \in I \tag{3.1.101}$$

It is now easy to verify that I is a Poisson subalgebra of $K(g)$. In order to see this let $f = f_\alpha \phi^\alpha$ and $h = h_\alpha \phi^\alpha$ be elements fo I, then

$$\begin{aligned} \{f, g\} &= \{f_\alpha, h_\beta\}\phi^\alpha \phi^\beta + \{f_\alpha, \phi^\beta\} h_\beta \phi^\alpha + \{\phi^\alpha, h_\beta\} f_\alpha \phi^\beta \\ &\quad + f_\alpha h_\beta \{\phi^\alpha, \phi^\beta\} \end{aligned}$$

Obviously the first three terms are again elements of I. That the last term is in I follows from the fact that the constraints $\{\phi^\alpha\}$ are all first class which means that there exist elements $\Phi^{\alpha\beta}_\gamma \in C^\infty(g)$ such that

$$\{\phi^\alpha, \phi^\beta\} = \Phi^{\alpha\beta}_\gamma \phi^\gamma \tag{3.1.102}$$

The interpretation of the ideal I is that it is the set of all smooth functions on g which are zero on the set

$$C = \{x \in g \mid \phi^\alpha(x) = 0, \text{ for all } \alpha\} \tag{3.1.103}$$

Obviously the space $C^\infty(C)$ of smooth functions on C is therefore isomorphic to the set

$$A \equiv C^\infty(g)/I \tag{3.1.104}$$

In the terminology that we used before this is the space of smooth functions on the zero set of the constraints. As we have seen the Kirillov Poisson structure on g does not induce a Poisson structure on C since C is plagued by gauge invariances generated by the first class constraints $\{\phi^\alpha\}$. We solved this problem before by gauge fixing. The algebraic version of this procedure will now be described.

Define the maps

$$X^\alpha : C^\infty(g) \to C^\infty(g) \tag{3.1.105}$$

by

$$X^\alpha(f) = \{\phi^\alpha, f\} \tag{3.1.106}$$

In geometric terms the X^α are of course the 'Hamiltonian vectorfields' associated to the constraints $\{\phi^\alpha\}$. Now, let $[f]$ denote the equivalence class $f + I$, i.e. $[f] \in A$, then

$$\begin{aligned} X^\alpha(f + I) &= \{\phi^\alpha, f\} + \{\phi^\alpha, I\} \\ &\subset X^\alpha(f) + I \end{aligned} \tag{3.1.107}$$

where we used the fact that I is a Poisson subalgebra of $K(g)$. Put differently we therefore have

$$X^\alpha[f] = [X^\alpha(f)] \tag{3.1.108}$$

which means that the maps X^α descend to well defined maps

$$X^\alpha : A \to A \tag{3.1.109}$$

In geometric terms this is nothing but the statement that the Hamiltonian vectorfields of the constraints are tangent to C.

Now define

$$A_{inv} = \{[f] \in A \mid X^\alpha[f] = 0, \text{ for all } \alpha\} \tag{3.1.110}$$

Interpretation of this space is that it is the space of smooth functions on $C/\hat{H}$ (which we showed to be isomorphic to g_{fix}), where as in the previous section $\hat{H}$ is the gauge group on C generated by the first class constraints $\{\phi^\alpha\}$.

The point is now that the Kirillov Poisson structure on $C^\infty(g)$ induces naturally a Poisson structure $\{.,.\}^*$ on A_{inv}. Let $[f], [h] \in A_{inv}$ then this Poisson structure is simply given by

$$\{[f], [h]\}^* = [\{f, h\}] \tag{3.1.111}$$

of course one has to show that this Poisson structure is well defined. For this we have to check two things: first we have to check whether $[\{f, h\}] \in A_{inv}$ whenever $[f], [h] \in A_{inv}$, and second we have to show that the definition does not depend on the choice of representants f and h. Both these checks are easily carried out and one finds that indeed the Poisson bracket $\{.,.\}^*$ turns A_{inv} into a Poisson algebra. The whole point is now that by construction the Poisson algebra

$$W(g; t_0) \equiv (A_{inv}, \{.,.\}^*) \tag{3.1.112}$$

is Poisson isomorphic to the finite W algebra $\mathcal{W}(i)$ where i is the sl_2 embedding associated to t_0. The whole construction described above is a purely algebraic description of finite W algebras.

Let us now consider how to construct the Poisson relations in $W(g, t_0)$ using the algebraic formalism described above. For this one has to construct the space A_{inv}. of course the algebra $C^\infty(g)$ is generated as an (abelian) associative algebra by the functions J^a. Also the constraints ϕ^α are of the form

$$\phi^\alpha = J^\alpha - \chi(J^\alpha) \tag{3.1.113}$$

where χ is defined by $\chi(J^\alpha) = 1$ if $t_\alpha = t_+$ and $\chi(J^\alpha) = 0$ otherwise. Therefore we immediately find that $A = C^\infty(g)/I$ is isomorphic to the abelian associative algebra generated by the functions $\{J^{\bar{\alpha}}\}$.

The next step is to construct A_{inv}. By definition this space correponds to those elements w of A such that

$$[\{\phi^\alpha, w\}] \equiv [\{J^\alpha, w\}] = 0 \tag{3.1.114}$$

or in geometric terms, the gauge invariant elements of C. In order to construct a set of generators of A_{inv} we can use theorems 1 and 5 which say that for any C valued function on g of the form

$$t_+ + \sum_{\bar{\alpha}} J^{\bar{\alpha}} t_{\bar{\alpha}} \tag{3.1.115}$$

there exists a unique element $a : g \to \hat{H}$ such that

$$a(t_+ + \sum_{\bar{\alpha}} J^{\bar{\alpha}} t_{\bar{\alpha}})a^{-1} = t_+ + \sum_{t_{\bar{\alpha}} \in g_{lw}} W^{\bar{\alpha}} t_{\bar{\alpha}} \tag{3.1.116}$$

The functions $W^{\bar{\alpha}}$ will in general be polynomials in the functions $\{J^{\bar{\alpha}}\}$ of the form

$$W^{\bar{\alpha}} = \sum_{l=0}^{p} W_l^{\bar{\alpha}} \tag{3.1.117}$$

where $\deg(W_l^{\bar{\alpha}}) = l$, $W_p^{\bar{\alpha}} = J^{\bar{\alpha}}$ and $p = \deg(J^{\bar{\alpha}})$. Obviously they are gauge invariant (i.e. invariant under $\hat{H}$) and generate the algebra A_{inv}. The Poisson relations in $W(g; t_0)$ can now easily be calculated using (3.1.111)

$$\{W^{\bar{\alpha}}, W^{\bar{\beta}}\}^* = \{W^{\bar{\alpha}}, W^{\bar{\beta}}\} \tag{3.1.118}$$

Let us give two examples of this construction. First we consider the (rather trivial) case of $g = sl_2$. The basis of sl_2 is given by

$$J^a t_a = \begin{pmatrix} J^2 & J^1 \\ J^3 & -J^2 \end{pmatrix} \tag{3.1.119}$$

of course $t_+ = t_1$ which means that the constraint is $\phi = J^1 - 1$. We therefore find that A is generated by J^2 and J^3. In order to construct the generator W^3 of A_{inv} we solve the equation

$$\begin{pmatrix} 1 & 0 \\ a_1 & 1 \end{pmatrix} \begin{pmatrix} J^2 & 1 \\ J^3 & -J^2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -a_1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ W^3 & 0 \end{pmatrix} \tag{3.1.120}$$

which yields $a_1 = J^2$ and $W^3 = J^3 + J^2 J^2$. Note that indeed we have

$$[\{W^3, \phi\}] \equiv [\{W^3, J^3\}] = 0 \tag{3.1.121}$$

verifying the gauge invariance of W^3. of course the relation of W^3 with itself is trivial $\{W^3, W^3\} = 0$.

In order to illustrate the construction in case of a slightly less trivial case consider again the nonprincipal sl_2 embedding into sl_3. For sl_3 we again choose the basis

$$J^a t_a = \begin{pmatrix} J^4 + \frac{1}{2}J^5 & J^2 & J^1 \\ J^6 & -2J^4 & J^3 \\ J^8 & J^7 & J^4 - \frac{1}{2}J^5 \end{pmatrix} \tag{3.1.122}$$

The constraints are again $\phi^3 = J^3$ and $\phi^1 = J^1 - 1$. Obviously A is generated by $\{J^2, J^4, \ldots, J^8\}$. In order to construct the generators $\{W^4, W^6, W^7, W^8\}$ (remember that $g_{lw} = \mathrm{Span}(t_4, t_6, t_7, t_8)$) we solve the equation

$$a(t_+ + \sum_{\bar{\alpha}} J^{\bar{\alpha}} t_{\bar{\alpha}})a^{-1} = t_+ + \sum_{t_{\bar{\alpha}} \in g_{lw}} W^{\bar{\alpha}} t_{\bar{\alpha}} \tag{3.1.123}$$

where

$$a = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a_1 & a_2 & 1 \end{pmatrix} \tag{3.1.124}$$

From this equation follows that $a_1 = \frac{1}{2}J^5$, $a_2 = J^2$ and

$$\begin{aligned} W^4 &= J^4 \\ W^6 &= J^6 \\ W^7 &= J^7 + \frac{1}{2}J^2J^5 - 3J^2J^4 \\ W^8 &= J^8 + \frac{1}{4}J^5J^5 + J^2J^6 \end{aligned} \tag{3.1.125}$$

(Remember that since $\{J^{\bar{\alpha}}\}$ are functions the ordering in the products is irrelevant. It only becomes relevant after quantization.) Again introducing the generators

$$\begin{aligned} C &= -\frac{4}{3}(W^8 + 3W^4W^4) \\ E &= W^7 \\ F &= \frac{4}{3}W^6 \\ H &= 4W^4 \end{aligned} \tag{3.1.126}$$

we find, using formula (3.1.111) that these form the algebra (3.1.91).

It is also possible to give a very simple description of the generalized Miura transformation in this algebraic framework. For this note that the Poisson bracket of a grade l_1 element with a grade l_2 element is an element of grade $l_1 + l_2$. From this it immediately follows that the Poisson brackets between the elements $W_0^{\bar{\alpha}}$ and $W_0^{\bar{\beta}}$ must have precisely the same form as the Poisson brackets between the elements $W^{\bar{\alpha}}$ and $W^{\bar{\beta}}$, i.e. the grade 0 pieces of the generators must form the same algebra. Therefore

$$W^{\bar{\alpha}} \mapsto W_0^{\bar{\alpha}} \tag{3.1.127}$$

is a Poisson algebra isomorphism. This is again the Miura map. In order to see this remember that the elements $W_0^{\bar{\alpha}}$ are polynomials in the functions $\{J^{\bar{\beta}}\}_{t_{\bar{\beta}} \in g_0}$. Also remember that these generate a Poisson algebra that is the Kirillov Poisson algebra of a semisimple Lie algebra, namely g_0. Therefore the map (3.1.127) is an embedding of the classical finite W algebra into a Kirillov Poisson algebra.

3.1.7 The symplectic geometry of finite W algebras

Suppose we are given a Poisson manifold, that is a manifold M together with a Poisson bracket $\{.,.\}$ on the space of smooth functions on M. In general such a Poisson structure is not associated to a symplectic form on M since the Poisson bracket can be degenerate, i.e. there may exist functions that Poisson commute with all other functions. The Poisson bracket $\{.,.\}$ does however induce a symplectic form on certain submanifolds of M. In order to see this consider the set of Hamiltonian vectorfields

$$X_f(\cdot) = \{f, .\} \qquad \text{for } f \in C^\infty(M) \tag{3.1.128}$$

Note that the set of functions on M that Poisson commute with all other functions are in the kernel of the map $f \mapsto X_f$. From this follows that in every tangent space the span of all Hamiltonian vectorfields is only a subspace of the whole tangentspace. We say that a Poisson structure is regular if the span of the set of Hamiltonian vectorfields has the same dimension in every tangentspace. Obviously a regular Poisson structure defines a tangent system on the manifold M and from the well known relation

$$[X_f, X_g] = X_{\{f,g\}} \tag{3.1.129}$$

it then immediately follows that this system is integrable (in the sense of Frobenius). Therefore M foliates into a disjoint union of integral manifolds of the Hamiltonian vectorfields. Obviously the restriction of the Poisson bracket to one of these integral manifolds is nondegenerate and therefore associated to a symplectic form. They are therefore called symplectic leaves.

The symplectic leaves play an important role in the representation theory of Poisson algebras. A representation of a Poisson algebra is a symplectic manifold S together with a map π from the Poisson algebra to the space of smooth functions on S that is linear and preserves both the multiplicative and Poisson structure of the Poisson algebra. Also the Hamiltonian vectorfield associated to $\pi(f) \in C^\infty(S)$ must be complete. A representation is called *irreducible* if span of the set $\{X_{\pi(f)}(s) \mid f \in C^\infty(M)\}$ is equal to the tangent space of S in s for all $s \in S$. The role of the symplectic leaves becomes clear from the following theorem [38]: If a representation of the Poisson algebra $C^\infty(M)$ is irreducible then S is symplectomorphic to a covering space of a symplectic leaf of M.

From the above it becomes clear that it is rather important to construct the symplectic leaves associated to finite W algebras. Looking at eq.(3.1.72) it is clear that constructing these symplectic orbits from scratch could be rather difficult. Luckily we can use the fact that the symplectic orbits of the Kirillov Poisson structure are known (by the famous Kostant-Souriau theorem they are nothing but the coadjoint orbits and

their covering spaces) to construct these. The answer turns out to be extremely simple and is given in the following theorem:

Theorem 2 *Let $\mathcal{O}$ be a coadjoint orbit of the Lie algebra g, then the intersection of $\mathcal{O}$ and g_{lw} is a symplectic orbit of the finite W algebra $W(g;t_0)$.*

Proof: Let again $C = \{x \in g \mid \phi^\alpha(x) = 0\}$. The Hamiltonian vectorfields $X^\alpha = \{\phi^\alpha, .\}$ form an involutive system tangent to C and therefore C foliates. g_{lw} has one point in common with every leaf and denote the canonical projection from C to g_{lw}, which projects an element $x \in C$ to the unique point $x' \in g_{lw}$, by π. This map induces a map π_* from the tangent bundle TC of C to the tangent bundle Tg_{lw} of g_{lw}. Let f be a 'gauge invariant' function on C, i.e. $\{f, \phi_\alpha\}|_C = 0$ then obviously $X_f = \{f, .\}$ is a section of TC (i.e. as a vectorfield it is tangent to C in every point of C). It need not be an element of Tg_{lw} however but using the gauge invariance we can project it back onto the gauge slice. This projection of X_f on the the gauge slice is given by $\pi_*(X_f) \in Tg_{lw}$. By construction the Dirac bracket now has the property that $\{f, .\}^* = \pi_*(X_f)$. What we now need to show is that $\pi_*(X_f)$ is tangent to the coadjoint orbit $\mathcal{O}$. Well, obviously we have $\pi_*(X_f) = X_f + Y$ where Y is a tangent to the gauge orbit. Since the spaces tangent to the gauge orbits are spanned by the Hamiltonian vectorfields X^α we find that Y is a tangent vector of $\mathcal{O}$. Since X_f is by definition a tangent vector of $\mathcal{O}$ we find that $\pi_*(X_f)$ is also tangent to the coadjoint orbit $\mathcal{O}$. What conclude that the Hamiltonian vectorfields on g_{lw} w.r.t. the reduced Poisson bracket are all tangent to the coadjoint orbit and therefore the symplectic orbits are submanifolds of the coadjoint orbits. The theorem now follows immediately .

Let us now recall some basic facts on coadjoint orbits of simple Lie algebras. Let f be an adinvariant function on g, i.e. $f(axa^{-1}) = f(x)$ for all $a \in G$ and $x \in g$, then we have

$$\begin{aligned} 0 &= \frac{d}{dt} f(e^{tx} y e^{-tx})|_{t=0} = \frac{d}{dt} f(y + t[x,y])|_{t=0} \\ &= \Big(\mathrm{grad}_y f, [x,y]\Big) = \Big([\mathrm{grad}_y f, x], y\Big) \end{aligned} \tag{3.1.130}$$

which means that

$$\Big([\mathrm{grad}_y f, \mathrm{grad}_y h], y\Big) = \{f,h\}(y) = 0 \tag{3.1.131}$$

We conclude that any adinvariant function Poisson commutes with all other functions. Conversely any function that Poisson commutes with all others is adinvariant because $\{f,h\} = 0$ for all h implies that $X_h(f) = 0$ for all h, where X_h is the Hamiltonian vectorfield associated to h, which means that the derivative of f in all directions tangent to a symplectic orbit are zero.

Consider now the Casimir functions $\{C_i\}_{i=1}^{\text{rank}(g)}$ of the Lie algebra g. Certainly these Poisson commute with all other functions and it can in fact be shown that the (co)adjoint (since we have identified the Lie algebra with its dual the adjoint and coadjoint orbits coincide) orbits of maximal dimension of the Lie algebra are given by their constant sets

$$C_{\vec{\mu}} = \{x \in g \mid C_i(x) = \mu_i\,;\ \mu_i \in \mathbf{R}\,;\ i = 1, \ldots, \text{rank}(g)\} \tag{3.1.132}$$

Their dimension is therefore $\dim(g) - \text{rank}(g)$.

Coadjoint orbits are very important to the representation theory of (semi)simple Lie groups. In order to see why let us recall the Borel-Weil-Bott (BWB) theorem [10]. Let G be a compact semisimple Lie group with maximal torus T and let R be a finite dimensional irreducible representation of G. The space of highest weight vectors of R is 1 dimensional (call it V) and furnishes a representation of T. The product space $G \times V$ is a trivial line bundle over G and its quotient by the action of T (where $(a, v) \sim (at, t^{-1}v)$ for $a \in G$ and $t \in T$) is a holomorphic line bundle over G/T (which is a complex manifold). Now, L admits a G action $(a, v) \to (a'a, v)$, where $a, a' \in G$ and $v \in V$, so the space of holomorphic sections of L is naturally a G representation. The BWB theorem now states that this representation is isomorphic to R. The BWB construction has however the restriction that it applies only to compact groups. There does exist however a generalization of the BWB construction called the coadjoint method of Kirillov. In this method one generalizes G/T to certain homogeneous spaces G/H which can be realized as coadjoint orbits. As we have mentioned above, a coadjoint orbit carries a natural (G invariant) symplectic form ω inherited from the Kirillov Poisson structure and can therefore be seen as a phase space of some classical mechanical system. One then attempts to quantize this symplectic manifold using the methods of geometric quantization [24]. For this one is supposed to construct a holomorphic line bundle L over the coadjoint orbit such that the first Chern class of L is equal to the (cohomology class of) ω. If such a line bundle exists the coadjoint orbit is called quantizable. For a quantizable orbit the bundle L can then be shown to admit a hermitean metric whose curvature is equal to ω. The space of sections of L thus obtains a Hilbert space structure and is interpreted as the physical Hilbert space of the quantum system associated to the coadjoint orbit. It carries a unitary representation of the group G and in fact one attempts to construct all unitary representations in this way. Obviously this construction is a generalization of the BWB method. Note that the fact that generic groups have unitary irreducible representations only for a discrete set of highest weights is translated into the fact that only a discrete set of orbits is quantizable.

From the above it is clear that the symplectic (or W-coadjoint) orbits of finite W algebras are extremely important especially in the study of global aspects of W algebras. In order to see this note that coadjoint orbits are homogeneous spaces of the Lie group G in question, and it is therefore tempting to interpret the symplectic orbits of W

algebras as *W-homogeneous spaces*. These naturally carry much information on the global aspects of W transformations. In the notation used above the symplectic orbits of the finite W algebra $W(g;t_0)$ are given by

$$\mathcal{O}_{\vec{\mu}} = C_{\vec{\mu}} \bigcap g_{lw} \tag{3.1.133}$$

which are therefore (generically) of dimension $\dim(g_{lw}) - \mathrm{rank}(g)$.

Let us give some examples. As we have seen the finite W algebras associated to principal sl_2 embeddings are trivial in the sense that they are Poisson abelian. This means that all Hamiltonian vectorfields are zero and that all symplectic orbits are points. Again the simplest nontrivial case is the algebra $W_3^{(2)}$. The 2 independent Casimir functions of sl_3 are

$$C_1 = \mathrm{Tr}(J^2) \quad \text{and } C_2 = \mathrm{Tr}(J^3) \tag{3.1.134}$$

where $J = J^a t_a$. The spaces $\mathcal{O}_{\vec{\mu}}$ can be constructed by taking for J the constrained and gauge fixed matrix

$$J = \begin{pmatrix} j & 0 & 1 \\ G_+ & -2j & 0 \\ T & G_- & j \end{pmatrix} \tag{3.1.135}$$

One then easily finds

$$\begin{aligned} C_1 &= 3j^2 + T \\ C_2 &= -6j^3 + 3G_+G_- + 6jT \end{aligned} \tag{3.1.136}$$

Obviously $\dim(g_{lw}) = 4$ and introducing the variables $x_1 = j;\ x_2 = G_+;\ x_3 = G_-;\ x_4 = T$ we find that

$$\mathcal{O}_{\mu_1,\mu_2} = \{\vec{x} \in \mathbf{R}^4 \mid 3x_1^2 + x_4 = \mu_1 \ \text{ and } \ 2x_1x_4 + x_2x_3 - 2x_1^3 = \mu_2\} \tag{3.1.137}$$

We can investigate the topological nature of these symplectic orbits by inserting $x_4 = \mu_1 - 3x_1^2$ into the second equation. We find that topologically $\mathcal{O}_{\mu_1,\mu_2}$ is equivalent to the 2 dimensional surface in $\mathbf{R}^3$ determined by the equation

$$x_2x_3 = \mu_2 - 2\mu_1x_1 + 8x_1^3 \tag{3.1.138}$$

In the plane $x_1 = \alpha$ this is generically a hyperbola but it degenerates if x_1 is a zero of the polynomial $\mu_2 - 2\mu_1x_1 + 8x_1^3$. For example the orbit $\mathcal{O}_{0,0}$ degenerates into the two lines $x_1 = x_3 = 0$ and $x_1 = x_2 = 0$ for $x \to 0$ and has a singular point in the origin. In fact this phenomenon also occurs in ordinary Lie group coadjoint orbits. Consider

for example the case of $SL_2(\mathbf{R})$ where the coadjoint orbits are given by the following hyperplanes in $\mathbf{R}^3$ [61]

$$x_2 x_3 = x_1^2 + \mu \tag{3.1.139}$$

where $\mu \in \mathbf{R}$. Here also for $\mu = 0$ the coadjoint orbit degenerates if $x_1 \to 0$. This coadjoint orbit is in fact conelike and for arbitrary non-compact semisimple Lie groups there are always just a finite number of them (infinite dimensional Lie groups may have infinitely many of them). The group representations that one can associate to them using the Kirillov method are called unipotent representations. We therefore interpret the $W(g; t_0)$ symplectic orbits that are singular as unipotent finite W representations. Also it should be noted that, in the $SL_2(\mathbf{R})$ case, the origin itself is also a coadjoint orbit, i.e. it is a fixed point of the group action. In the same way the singular points in the unipotent W orbits are fixed points of global W transformations.

This concludes our presentation of the theory of classical finite W algebras.

3.2 Quantum finite W Algebras

Having developed the classical theory of finite W algebras we now turn to their quantization. In quantum mechanics, quantization amounts to replacing Poisson brackets by commutators, or in a mathematically more sophisticated language, replacing a Poisson algebra $(\mathcal{A}_0, \{.,.\})$ by an associative algebra $\mathcal{A}$ depending on a parameter $\hbar$ such that

- $\mathcal{A}$ is a free $\mathbb{C}[[\hbar]]$ module,
- $\mathcal{A}/\hbar\mathcal{A} \simeq \mathcal{A}_0$
- if π denotes the natural map $\pi : \mathcal{A} \to \mathcal{A}/\hbar\mathcal{A} \simeq \mathcal{A}_0$, then $\{\pi(X), \pi(Y)\} = \pi((XY - YX)/\hbar)$.

In most cases one has a set of generators for $\mathcal{A}_0$, and $\mathcal{A}$ is completely fixed by giving the commutation relations of these generators.

For example, let $\mathcal{A}_0$ be the Kirillov Poisson algebra $K(g)$ associated to g, then a quantization of this Poisson algebra is the algebra $\mathcal{A}$ generated by the J^a and $\hbar$, subject to the commutation relations $[J^a, J^b] = \hbar f^{ab}_{\ c} J^c$ (note that here the J^a are no longer functions on g but the quantum objects associated to them). Obviously, the Jacobi identities are satisfied. Specializing to $\hbar = 1$, this algebra is precisely the universal enveloping algebra $\mathcal{U}g$ of g.

To find quantizations of finite W algebras, one can first reduce the sl_n Kirillov Poisson algebra, and then try to quantize the resulting algebras that we studied in the previous sections. On the other hand, on can also first quantize and then constrain. We

will follow the latter approach, and thus study the reductions of the quantum Kirillov algebra

$$[J^a, J^b] = \hbar f^{ab}_{\ c} J^c. \tag{3.2.140}$$

We want to impose the same constraints on this algebra as we imposed previously on the Kirillov Poisson algebra, to obtain the quantum versions of the finite W algebras related to sl_2 embeddings. Imposing constraints on quantum algebras can be done using the BRST formalism [86].

3.2.1 BRST quantization of finite W algebras

Consider again the map χ defined by $\chi(J^\alpha) = 1$ if $t_\alpha = t_+$ and $\chi(J^\alpha) = 0$ otherwise. In terms of χ, the constraints can be written as $\pi_+(J) = \chi(\pi_+(J))$, where π_+ denotes the projection $g \to g_+$ and $J = J^a t_a$ It is this form of the constraints that we will use. Furthermore we will take $\hbar = 1$ for simplicity; the explicit $\hbar$ dependence can be determined afterwards.

To set up the BRST framework we need to introduce anticommuting ghosts and antighosts c_α and b^α, associated to the constraints that we want to impose [86]. They satisfy $b^\alpha c_\beta + c_\beta b^\alpha = \delta^\alpha_\beta$ and generate the Clifford algebra $Cl(g_+ \oplus g_+^*)$. The quantum Kirillov algebra is just the universal enveloping algebra $\mathcal{U}g$, and the total space on which the BRST operator acts is $\Omega = \mathcal{U}g \otimes Cl(g_+ \oplus g_+^*)$. A $\mathbf{Z}$ grading on Ω is defined by $\deg(J^a) = 0$, $\deg(c_\alpha) = +1$ and $\deg(b^\alpha) = -1$, and we can decompose $\Omega = \oplus_k \Omega^k$ accordingly. The BRST differential on Ω is given by $d(X) = [Q, X]$, where Q is the BRST charge

$$Q = (J^\alpha - \chi(J^\alpha))c_\alpha - \tfrac{1}{2} f^{\alpha\beta}_{\ \ \gamma} b^\gamma c_\alpha c_\beta. \tag{3.2.141}$$

and $[.,.]$ denotes the graded commutator (as it always will from now on)

$$[A, B] = AB - (-1)^{\deg(A).\deg(B)} BA \tag{3.2.142}$$

Note that $\deg(Q) = 1$.

This is the standard BRST complex associated to the first class constraints of the previous section. Of interest are the cohomology groups of this complex, $H^k(\Omega; d)$. The zeroth cohomology group is the quantization of the classical finite W algebra. Because the gauge group $\hat{H}$ in (39) acts properly on C, we expect the higher cohomologies of the BRST complex to vanish, as they are generically related to singularities in the quotient $C/\hat{H}$. In the mathematics literature the cohomology of the BRST complex is called the Hecke algebra $\mathcal{H}(g, g_+, \chi)$ associated to g, g_+, χ. Hecke algebras related to arbitrary sl_2 embeddings have not been computed, apart from those related to the principal sl_2 embeddings. In that case it was shown by Kostant [88] that the only

nonvanishing cohomology is $H^0(\Omega; d)$ and that it is isomorphic to the center of the universal enveloping algebra. Recall that the center of the $\mathcal{U}g$ is generated by the set of independent Casimirs of g. This set is closely related to the generators of standard infinite W_n-algebras; in that case there is one W field for each Casimir which form a highly nontrivial algebra [89]. We see that for finite W algebras the same generators survive, but that they form a trivial abelian algebra. For non principal sl_2 embeddings however quantum finite W algebras are non-trivial.

We now want to compute the cohomology of $(\Omega; d)$. Unfortunately this is a difficult problem to approach directly. We therefore use a well known trick of cohomology theory, namely we split the complex $(\Omega; d)$ into a double complex and calculate the cohomology via a spectral sequence argument. For this note that the operator d can be decomposed into two anti commuting pieces. Write $Q = Q_0 + Q_1$, with

$$\begin{aligned} Q_1 &= J^\alpha c_\alpha - \tfrac{1}{2} f^{\alpha\beta}_{\ \ \gamma} b^\gamma c_\alpha c_\beta, \\ Q_0 &= -\chi(J^\alpha) c_\alpha, \end{aligned} \tag{3.2.143}$$

and define $d_0(X) = [Q_0, X]$, $d_1(X) = [Q_1, X]$, then one can verify by explicit computation that $d_0^2 = d_0 d_1 = d_1 d_0 = d_1^2 = 0$. Associated to this decomposition is a bigrading of $\Omega = \oplus_{k,l} \Omega^{k,l}$ defined by

$$\begin{aligned} \deg(J^a) &= (-k, k), \qquad \text{if } t_a \in g_k \\ \deg(c_\alpha) &= (k, 1-k), \qquad \text{if } t_\alpha \in g_k \\ \deg(b^\alpha) &= (-k, k-1), \qquad \text{if } t_\alpha \in g_k, \end{aligned} \tag{3.2.144}$$

with respect to which d_0 has degree $(1, 0)$ and d_1 has degree $(0, 1)$. Thus $(\Omega^{k,l}; d_0; d_1)$ has the structure of a double complex. Explicitly, the action of d_0 and d_1 is given by

$$\begin{aligned} d_1(J^a) &= f^{\alpha a}_{\ \ b} J^b c_\alpha, \\ d_1(c_\alpha) &= -\tfrac{1}{2} f^{\beta\gamma}_{\ \ \alpha} c_\beta c_\gamma, \\ d_1(b^\alpha) &= J^\alpha + f^{\alpha\beta}_{\ \ \gamma} b^\gamma c_\beta, \\ d_0(J^a) = d_0(c_\alpha) &= 0, \\ d_0(b^\alpha) &= -\chi(J^\alpha). \end{aligned} \tag{3.2.145}$$

To simplify the algebra, it is advantageous to introduce

$$\hat{J}^a = J^a + f^{a\beta}_{\ \ \gamma} b^\gamma c_\beta. \tag{3.2.146}$$

Our motivation to introduce these new elements $\hat{J}^a$ is twofold: first, similar expressions were encountered in a study of the effective action for W_3 gravity [69], where it turned

out that the BRST cohomology for the infinite W_3 algebra case could conveniently be expressed in terms of $\hat{J}$'s; second, such expressions were introduced for the J^a's that live on the Cartan subalgebra of g in [20], and simplified their analysis considerably. In terms of $\hat{J}$ we have

$$\begin{aligned} d_1(\hat{J}^a) &= f^{\alpha a}_{\ \ \bar{\gamma}} \hat{J}^{\bar{\gamma}} c_\alpha, \\ d_1(c_\alpha) &= -\tfrac{1}{2} f^{\beta\gamma}_{\ \ \alpha} c_\beta c_\gamma, \\ d_1(b^\alpha) &= \hat{J}^\alpha, \\ d_0(\hat{J}^a) &= -f^{a\beta}_{\ \ \gamma} \chi(J^\gamma) c_\beta, \\ d_0(c_\alpha) &= 0, \\ d_0(b^\alpha) &= -\chi(J^\alpha). \end{aligned} \tag{3.2.147}$$

The advantage of having a double complex, is that we can apply the techniques of spectral sequences [90] to it, in order to compute the cohomology of $(\Omega; d)$. The results from the theory of spectral sequences that we need are gathered in the next section.

3.2.2 Spectral Sequences for Double Complexes

Let $(\Omega^{p,q}; d_0; d_1)$ denote a double complex, where d_0 has degree $(1,0)$ and d_1 has degree $(0,1)$. The standard spectral sequence for this double complex is a sequence of complexes $(E_r^{p,q}; D_r)_{r\geq 0}$, where D_r is a differential of degree $(1-r, r)$, and is defined as follows: $E_0^{p,q} = \Omega^{p,q}$, $D_0 = d_0$, $D_1 = d_1$, and for $r \geq 0$

$$E_{r+1}^{p,q} = H^{(p,q)}(E_r; D_r) = \frac{\ker\ D_r : E_r^{p,q} \to E_r^{p+1-r,q+r}}{\mathrm{im}\, D_r : E_r^{p-1+r,q-r} \to E_r^{p,q}}. \tag{3.2.148}$$

The differential D_{r+1} for $r > 0$ is given by $D_{r+1}(\alpha) = d_1(\beta)$, where β is chosen such that $d_0\beta = D_r\alpha$. Such a β always exists and D_{r+1} is uniquely defined in this way. The usefulness of this spectral sequence is provided by the following [91]

Theorem 3 *If* $\cap_q \oplus_{p,q\geq s} \Omega^{p,q} = \{0\}$, *then* $E_\infty^{p,q} = \cap_r E_r^{p,q}$ *exists, and*

$$E_\infty^{p,q} \simeq \frac{F^q H^{(p+q)}(\Omega; d)}{F^{q+1} H^{(p+q)}(\Omega; d)}, \tag{3.2.149}$$

where

$$F^q H^{(p+q)}(\Omega; d) = \frac{\ker\ d : \oplus_{s\geq q} \Omega^{p+q-s,s} \to \oplus_{s\geq q} \Omega^{p+q+1-s,s}}{\mathrm{im}\ d : \oplus_{s\geq q} \Omega^{p+q-1-s,s} \to \oplus_{s\geq q} \Omega^{p+q-s,s}} \tag{3.2.150}$$

This spectral sequence is especially useful if it collapses at the n^{th} term for some n, *i.e.* $D_r = 0$ for $r \geq n$, because then $E_n \simeq E_\infty$ and one needs only to compute the first n terms of the spectral sequence.

If the double complex is also an algebra, i.e. there is a multiplication operator $m : \Omega^{p,q} \otimes_{\mathbb{C}} \Omega^{p',q'} \to \Omega^{p+p',q+q'}$, and d satisfies the Leibniz rule with respect to this multiplication, then (3.2.149) is also an equality of algebras. It is in general nontrivial to reconstruct the full algebra structure of $H^*(\Omega; d)$ from $E_\infty^{p,q}$, due to the quotient in the right hand side of (3.2.149).

Another useful tool in the computation of cohomology is the following version of the Künneth theorem

Lemma 2 *Let $A = \oplus_k A^k$ be a graded differential algebra over $\mathbb{C}$ with a differential d of degree 1 that satisfies the Leibniz rule, and assume that A has two graded subalgebras $A_1 = \oplus_k A_1^k$ and $A_2 = \oplus_k A_2^k$ such that $d(A_1) \subset A_1$ and $d(A_2) \subset A_2$, that $A_2^k = \{0\}$ for k sufficiently large, and that $m : A_1 \otimes_{\mathbb{C}} A_2 \to A$ given by $m(a_1 \otimes a_2) = a_1 a_2$ is an isomorphism of vector spaces. Then*

$$H^*(A; d) \simeq \{a_1 a_2 | a_1 \in H^*(A_1; d), a_2 \in H^*(A_2; d)\}. \tag{3.2.151}$$

Proof: form the double complex $(\Omega^{p,q}; d_0; d_1)$ with $\Omega^{p,q} \simeq m(A_1^p \otimes A_2^q)$, $d_0(a_1 a_2) = d(a_1)a_2$, and $d_1(a_1 a_2) = (-1)^{\deg(a_1)} a_1 d(a_2)$. The spectral sequence for this double complex collapses at the E_2 term, and one finds $E_\infty^{p,q} = \{a_1 a_2 | a_1 \in H^p(A_1; d), a_2 \in H^q(A_2; d)\}$. The condition $A_2^k = 0$ for k sufficiently large guarantees that $F^q H^{p+q} = 0$ for q sufficiently large, and one can assemble $H^*(\Omega; d)$ from $E_\infty^{p,q}$ using (3.2.149). This leads to (3.2.151) on the level of vector spaces. Because $a_1 a_2$ is really a representative of an element of $H^*(A; d)$, it follows that (3.2.151) is also an isomorphism of algebras.

By induction, one can easily prove that the theorem still holds if instead of two subalgebras n subalgebras $A_1, \ldots A_n$ are given, with $A \simeq A_1 \otimes \ldots \otimes A_n$. The condition $A_2^k = \{0\}$ for sufficiently large k is replaced by $A_2^k = \ldots = A_n^k = \{0\}$ for sufficiently large k. Without such a condition, it may not be so easy to reassemble H^* from E_∞. To illustrate some of the difficulties that can arise, let us give an example where H^* cannot be recovered directly from E_∞. This example is not related to the above theorem, but it represents a situation we will encounter in the computation of the BRST cohomology.

Consider the algebra $\Omega = \mathbb{C}[x, y]/(y^2 = 0)$, where x is an even generator of bidegree $(1, -1)$, and y is an odd generator of bidegree $(0, -1)$. The differentials d_0, d_1 are given by $d_{0,1}(x) = 0$, $d_0(y) = x$ and $d_1(y) = -1$. One immediately computes $H^k(\Omega; d) = \mathbb{C}[x]/(x-1)\mathbb{C}[x] \simeq \mathbb{C}$ for $k = 0$, and $H^k = 0$ otherwise. The spectral sequence associated to the double complex collapses at the first term, and one finds $E_\infty^{p,q} = \mathbb{C}\delta_{p,0}\delta_{q,0}$. Because

$F^1H^* = 0$, one deduces that $H^k(\Omega; d) \simeq \mathbb{C}\delta_{k,0}$. On the other hand, we could also have started with the mirror double complex obtained by interchanging d_0 and d_1 and the bigrading. Thus, we assign bidegree $(-1, 1)$ to x and bidegree $(-1, 0)$ to y. The spectral sequence associated to the mirror double complex also collapses at the first term, but now one finds $E_\infty^{p,q} = 0$. This is not in conflict with the previous computation, because we cannot a priori find a q for which $F^qH^{p+q} = 0$, and we can only conclude that $F^qH^{p+q} \simeq F^{q+1}H^{p+q}$. If we compute explicitly with respect to this bigrading what F^qH^{p+q} is, we find that it is only nontrivial for $p + q = 0$, and then $F^qH^0 = x^q\mathbb{C}[x]/(x-1)x^q\mathbb{C}[x] \simeq \mathbb{C}$ for $q \geq 0$, and $F^qH^0 = \mathbb{C}[x]/(x-1)\mathbb{C}[x] \simeq \mathbb{C}$ for $q < 0$. This indeed yields $E_\infty^{p,q} = 0$. The lesson is that one should be careful in deriving $H^*(\Omega; d)$ from $E_\infty^{p,q}$.

Finally, let us present another fact that will be useful later.

Lemma 3 *Suppose A is a differential graded algebra, $A = \oplus_{n\geq 0}A^n$, with a differential of degree 1. Assume furthermore that A has a filtration*

$$\{0\} = F^0A \subset F^1A \subset F^2A \subset \cdots \subset A, \tag{3.2.152}$$

such that $F^pAF^qA \subset F^{p+q}A$, and that d preserves the filtration, $d(F^pA) \subset F^pA$. If $H^k(F^{p+1}A/F^pA; d) = 0$ unless $k = 0$, then we have the following isomorphism of vector spaces

$$H^0(A; d) \simeq \oplus_{p\geq 0}H^0(F^{p+1}A/F^pA; d). \tag{3.2.153}$$

Proof: One can assign a spectral sequence to such a filtered graded algebra [91], whose first term contains the cohomologies $H^k(F^{p+1}A/F^pA; d)$. If only $H^0 \neq 0$, then the spectral sequence collapses at the first term, and because the filtration is bounded from below ($\{0\} = F^0A$), one can collect the vector spaces that make up $E_\infty^{*,*}$, to get the isomorphism (3.2.153).

3.2.3 The BRST Cohomology

The first people to propose using the theory of spectral sequences in the setting of W algebras were Feigin and Frenkel [20]. In fact, in this way they computed the BRST cohomology in the infinite dimensional case but only for the special example of the principal sl_2 embedding (which are known to lead to the W_N algebras). Their calculation however has the drawback that it is very difficult to generalize to arbitrary embeddings and that it constructs the cohomology in an indirect way (via commutants of screening operators). Now, to any double complex there are canonically associated two spectral sequences which can be considered to be each others mirror image. In [97] it was first proposed to calculate the BRST cohomology using a spectral sequence that is the mirror

spectral sequence of the one used by Feigin and Frenkel. As it turns out this has drastic simplifying consequences for the calculation of the cohomology.

The following theorem [97] gives the BRST cohomology on the level of vectorspaces.

Theorem 4 *As before let $g_{lw} \subset g$ be the kernel of the map $\mathrm{ad}_{t_-} : g \to g$. Then the BRST cohomology is given by the following isomorphisms of vector spaces*

$$H^k(\Omega; d) \simeq (\mathcal{U}g_{lw})\delta_{k,0}. \tag{3.2.154}$$

Proof: The computation of the BRST cohomology is simplified considerably due to the introduction of the new set of generators $\hat{J}^a$. The simplification arises due to the fact that $H^*(\Omega; d) \simeq H^*(\Omega_{red}; d)$, where Ω_{red} is the subalgebra of Ω generated by $\hat{J}^{\bar{\alpha}}$ and c_α. In order to show this apply the Künneth theorem 2 to $\Omega_{red} \otimes (\otimes_\alpha \Omega_\alpha)$, where Ω_α is the algebra generated by $\hat{J}^\alpha$ and b^α. Note that $[\hat{J}^\alpha, b^\alpha] = 0$ and that the conditions of the Künneth theorem are satisfied. Therefore, $H^*(\Omega; d) \simeq H^*(\Omega_{red}; d) \otimes (\otimes_\alpha H^*(\Omega_\alpha; d))$. Now $(\Omega_\alpha; d)$ is essentially the same complex as the one we examined in the last part of the previous section, and one easily proves that $H^k(\Omega_\alpha; d) \simeq \mathbb{C}\delta_{k,0}$. This shows $H^*(\Omega; d) \simeq H^*(\Omega_{red}; d)$ The reduced complex $(\Omega_{red}; d)$ is described by the following set of relations:

$$\begin{aligned}
d_1(\hat{J}^{\bar{\alpha}}) &= f^{\alpha\bar{\alpha}}_{\ \ \bar{\gamma}} \hat{J}^{\bar{\gamma}} c_\alpha, \\
d_1(c_\alpha) &= -\tfrac{1}{2} f^{\beta\gamma}_{\ \ \alpha} c_\beta c_\gamma, \\
d_0(\hat{J}^{\bar{\alpha}}) &= -f^{\bar{\alpha}\beta}_{\ \ \gamma} \chi(J^\gamma) c_\beta, \\
d_0(c_\alpha) &= 0, \\
[\hat{J}^{\bar{\alpha}}, \hat{J}^{\bar{\beta}}] &= f^{\bar{\alpha}\bar{\beta}}_{\ \ \bar{\gamma}} \hat{J}^{\bar{\gamma}}, \\
[\hat{J}^{\bar{\alpha}}, c_\beta] &= -f^{\bar{\alpha}\gamma}_{\ \ \beta} c_\gamma, \\
[c_\alpha, c_\beta] &= 0.
\end{aligned} \tag{3.2.155}$$

The E_1 term of the spectral sequence is now given by the d_0 cohomology of Ω_{red}. To compute the cohomology we use lemma 3. Define the filtration on Ω_{red} by declaring $F^p\Omega_{red}$ to be spanned as a vector space by $\{\hat{J}^{\bar{\alpha}_1}\hat{J}^{\bar{\alpha}_2}\cdots\hat{J}^{\bar{\alpha}_r} c_{\beta_1} c_{\beta_2} \cdots c_{\beta_s} | r + s \leq p\}$. Thus $F^p\Omega_{red}/F^{p-1}\Omega_{red}$ is spanned by the products of precisely p $\hat{J}$'s and c's, and in this quotient $\hat{J}$ and c (anti)commute with each other. Now let us rewrite $d_0(\hat{J}^{\bar{\alpha}})$ as

$$\begin{aligned}
d_0(\hat{J}^{\bar{\alpha}}) &= -\mathrm{Tr}([\chi(J^\gamma)t_\gamma, t^{\bar{\alpha}}]t^\beta c_\beta) \\
&= -\mathrm{Tr}([t_+, t^{\bar{\alpha}}]t^\beta c_\beta).
\end{aligned} \tag{3.2.156}$$

From this it is clear that $d_0(\hat{J}^{\bar{\alpha}}) = 0$ for $t^{\bar{\alpha}} \in g_{hw}$. Furthermore, since $t_{\bar{\alpha}} \in g_0 \oplus g_-$ and $\dim(g_{lw}) = \dim(g_0)$, it follows that for each β there is a linear combination $a(\beta)_{\bar{\alpha}}\hat{J}^{\bar{\alpha}}$ with $d_0(a(\beta)_{\bar{\alpha}}\hat{J}^{\bar{\alpha}}) = c_\beta$. This proves that

$$\bigoplus_{p>0} \frac{F^p\Omega_{red}}{F^{p-1}\Omega_{red}} \simeq \bigotimes_{t_{\bar{\alpha}} \in g_{lw}} \mathbb{C}[\hat{J}^{\bar{\alpha}}] \bigotimes_{t_{\bar{\alpha}} \notin g_{lw}} (\mathbb{C}[\hat{J}^{\bar{\alpha}}] \oplus d_0(\hat{J}^{\bar{\alpha}})\mathbb{C}[\hat{J}^{\bar{\alpha}}]). \tag{3.2.157}$$

Using the Künneth theorem (lemma 2) for (3.2.157), we find that

$$H^k(\Omega_{red}; d_1) = \bigotimes_{t_{\bar{\alpha}} \in g_{lw}} \mathbb{C}[\hat{J}^{\bar{\alpha}}]\delta_{k,0} = (\mathcal{U}g_{lw})\delta_{k,0}. \tag{3.2.158}$$

Because there is only cohomology of degree 0, the spectral sequence collapses, and $E_\infty = E_1$. Because $\Omega^{k,l}_{red} = 0$ for $l > 0$, we can find $H^*(\Omega_{red}; d)$ from E_∞. This proves the theorem.

As expected, there is only cohomology of degree zero, and furthermore, the elements of g_{lw} are in one-to-one correspondence with the components of g that made up the lowest weight gauge in section 1. Therefore $H^*(\Omega; d)$ really is a quantization of the finite W algebra. What remains to be done is to compute the algebraic structure of $H^*(\Omega; d)$. The only thing that (3.2.154) tells us is that the product of two elements a and b of bidegree $(-p, p)$ and $(-q, q)$ is given by the product structure on $\mathcal{U}g_{lw}$, modulo terms of bidegree $(-r, r)$ with $r < p + q$. To find these lower terms we need explicit representatives of the generators of $H^0(\Omega; d)$ in Ω. Such representatives can be constructed using the so-called tic-tac-toe construction [90]: take some $\phi_0 \in g_{lw}$, of bidegree $(-p, p)$. Then $d_0(\phi_0)$ is of bidegree $(1 - p, p)$. Since $d_1 d_0(\phi_0) = -d_0 d_1(\phi_0) = 0$, and there is no d_1 cohomology of bidegree $(1-p, p)$, $d_0(\phi_0) = d_1(\phi_1)$ for some ϕ_1 of bidegree $(1-p, p-1)$. Now repeat the same steps for ϕ_1, giving a ϕ_2 of bidegree $(2-p, p-2)$, such that $d_0(\phi_1) = d_1(\phi_2)$. Note that $d_1 d_0(\phi_1) = -d_0 d_1(\phi_1) = -d_0^2(\phi) = 0$. In this way we find a sequence of elements ϕ_l of bidegree $(l - p, p - l)$. The process stops at $l = p$. Let

$$W(\phi) = \sum_{l=0}^{p} (-1)^l \phi_l \tag{3.2.159}$$

Then $dW(\phi) = 0$, and $W(\phi)$ is a representative of ϕ_0 in $H^0(\Omega; d)$. The algebraic structure of $H^0(\Omega; d)$ is then determined by calculating the commutation relations of $W(\phi)$ in Ω, where ϕ_0 runs over a basis of g_{lw}. This is the quantum finite W algebra.

Let us now give an example of the construction described above.

3.2.4 Example

Consider again the nonprincipal sl_2 embedding into sl_3 associated to the following partition of the number 3: $3 = 2 + 1$. We constructed the classical W associated to this embedding earlier. We shall now quantize this Poisson algebra by the methods developed above. Take the following basis of sl_3:

$$r_a t_a = \begin{pmatrix} \frac{r_4}{6} - \frac{r_5}{2} & r_2 & r_1 \\ r_6 & -\frac{r_4}{3} & r_3 \\ r_8 & r_7 & \frac{r_4}{6} + \frac{r_5}{2} \end{pmatrix}. \tag{3.2.160}$$

Remember that (in the present notation) the sl_2 embedding is given by $t_+ = t_1$, $t_0 = -t_5$ and $t_- = t_8$. The nilpotent subalgebra g_+ is spanned by $\{t_1, t_3\}$, g_0 by $\{t_2, t_4, t_5, t_6\}$ and g_- by $\{t_7, t_8\}$. The d_1 cohomology of Ω_{red} is generated by $\{\hat{J}^4, \hat{J}^7, \hat{J}^6, \hat{J}^8\}$, and using the tic-tac-toe construction one finds representatives for these generators in $H^0(\Omega_{red}; d)$:

$$\begin{aligned} W(\hat{J}^4) &= \hat{J}^4, \\ W(\hat{J}^6) &= \hat{J}^6, \\ W(\hat{J}^7) &= \hat{J}^7 - \tfrac{1}{2}\hat{J}^2\hat{J}^5 - \tfrac{1}{2}\hat{J}^4\hat{J}^2 + \tfrac{1}{2}\hat{J}^2, \\ W(\hat{J}^8) &= \hat{J}^8 + \tfrac{1}{4}\hat{J}^5\hat{J}^5 + \hat{J}^2\hat{J}^6 - \hat{J}^5. \end{aligned} \tag{3.2.161}$$

Let us introduce another set of generators

$$\begin{aligned} C &= -\tfrac{4}{3}W(\hat{J}^8) - \tfrac{1}{9}W(\hat{J}^4)W(\hat{J}^4) - 1, \\ H &= -\tfrac{2}{3}W(\hat{J}^4) - 1, \\ E &= W(\hat{J}^7), \\ F &= \tfrac{4}{3}W(\hat{J}^6). \end{aligned} \tag{3.2.162}$$

The commutation relations between these generators are given by

$$\begin{aligned} [H, E] &= 2E, \\ [H, F] &= -2F, \\ [E, F] &= H^2 + C, \\ [C, E] = [C, F] = [C, H] &= 0. \end{aligned} \tag{3.2.163}$$

These are precisely the same as the relations for the finite $W_3^{(2)}$ algebra given in [74]. Notice that in this case the quantum relations are identical to the classical ones. The explicit $\hbar$ dependence can be recovered simply by multiplying the right hand sides of (3.2.163) by $\hbar$.

In [72] Rocek studied nonlinear deformations of $su(2)$ by imposing Jacobi identities on a general deformation. He also gave a general discussion of the existence of unitary representations and some other features of their representation theory. As a solution of the Jacobi identities he found $\bar{W}_3^{(2)}$. In [73] an attempt was made to formulate a gauge theory based on this nonlinear algebra by using the Noether method. However due to the nonlinearity of $\bar{W}_3^{(2)}$ the action they obtained has infinitely many terms. It is therefore an interesting fact that this algebra is a reduction of sl_3 as we have shown above. It might be possible to construct a manageable action for $\bar{W}_3^{(2)}$ using this knowledge.

In the appendix we also discuss the explicit quantization of all the finite W algebras that can be obtained from sl_4. There one does encounter certain quantum effects, i.e the quantum relations will contain terms of order $\hbar^2$ or higher.

In this section we have seen that it is possible to quantize the classical finite W algebras introduced in the first section of this chapter using the BRST formalism. What we have obtained are nonabelian and in general nonlinear associative algebras. These algebras are in principle still abstract. For applications to physics we are interested in the representation theory of quantum finite W algebras. This is the subject of the final section of this chapter.

3.3 The Representation Theory of Finite W algebras

In physics symmetry groups are always represented in terms of unitary linear operators. It follows that knowledge of the representation theory of groups and algebras is as important to physics as knowledge of their structure. For any physical application of finite W algebras it is therefore crucial to develop their representation theory.

Even if the representation theory of finite W algebras turns out to be of no particular interest to physics on its own it certainly will be important to the theory of infinite W algebras. As we said in the introduction of this chapter the structure and representation theory of affine Lie algebras is determined for a large part by the structure and representation theory of simple Lie algebras. For example, the space of singular or highest weight vectors of an affine Lie algebra representation will carry a representation of the simple Lie algebra underlying the affine algebra. We can therefore sweep out the set of singular vectors by acting with this underlying finite algebra on the vacuum state of the theory (which is of course also a singular vector).

Answering questions like 'what are the W singular vectors' have turned out to be rather difficult even though some progress has been made in particular cases. Here finite W algebras may provide an important now tool for the same reasons as described above. So also seen from this point of view it is very important to develop the representation theory of finite W algebras.

Let us therefore begin with the simplest nontrivial case, the finite W algebra $\bar{W}_3^{(2)}$. The finite dimensional representation theory of this algebra is the subject of the next section.

3.3.1 The representation theory of $\bar{W}_3^{(2)}$

Consider again the associative algebra generated by $\{H, E, F, C\}$ subject to the relations (3.2.163). We shall now construct the finite dimensional representation theory of this algebra. For this we are going to need the following

Lemma 4 *The following identities are true*

$$\begin{aligned} [H, E^k] &= 2kE^k \\ [H, F^k] &= -2kF^k \\ [E, F^k] &= F^{k-1}(kH^2 + kC - 2k(k-1)H + \frac{2}{3}k(k-1)(2k-1)) \end{aligned} \quad (3.3.164)$$

Proof: One can easily prove these identities by induction using the defining relations of $\bar{W}_3^{(2)}$ and the formula

$$\sum_{i=1}^{p-1} i^2 = \frac{1}{6}p(p-1)(2p-1) \quad (3.3.165)$$

Consider the real span of H and C to be the Cartan subalgebra. The next theorem then identifies the highest weight representations of $\bar{W}_3^{(2)}$.

Theorem 5 *Let p be a positive integer and x a real number.*

1. *For every pair (p, x) the algebra $\bar{W}_3^{(2)}$ has a unique highest weight representation $W(p, x)$ of dimension p.*

2. *$W(p, x)$ is spanned by eigenvectors (weight vectors) of H. The weights are given by*

$$\{j(p, x) - 2k\}_{k=0}^{p-1} \quad (3.3.166)$$

where the highest weight $j(p, x)$ is equal to

$$j(p, x) = p + x - 1 \quad (3.3.167)$$

3. The value of the central element C on $W(p,x)$ is given by

$$c(p,x) = \frac{1}{3}(1-p^2) - x^2 \tag{3.3.168}$$

Proof: Define the p-dimensional representation W as follows: Let v be a (formal) element and define

$$\begin{aligned} H \cdot v &= jv \\ E \cdot v &= 0 \\ C \cdot v &= cv \end{aligned}$$

Define the vectorspace W as the span of the set

$$\{v_k \equiv F^k \cdot v\}_{k=0,1,2,\ldots} \tag{3.3.169}$$

W is then a representation of $W_3^{(2)}$ and by the previous lemma we find

$$\begin{aligned} H \cdot v_k &= (j-2k)v_k \\ E \cdot v_k &= (k(j^2+c) - 2k(k-1)j + \frac{2}{3}k(k-1)(2k-1))v_{k-1} \\ C \cdot v_k &= cv_k \end{aligned}$$

Since we are looking for finite dimensional representations there must be a positive integer p such that $F^p v = 0$. From

$$\begin{aligned} 0 &= EF^p \cdot v \\ &= (pj^2 - 2p(p-1)j + pc + \frac{2}{3}p(p-1)(2p-1))v_{p-1} \end{aligned}$$

follows that

$$pj^2 - 2p(p-1)j + pc + \frac{2}{3}p(p-1)(2p-1) = 0 \tag{3.3.170}$$

Solving this equation for j and demanding that it be real we find that

$$\begin{aligned} c(p,x) &= \frac{1}{3}(1-p^2) - x^2 \\ j(p,x) &= p + x - 1 \end{aligned}$$

where x is an arbitrary real number. This proves the theorem .

So we get infinitely many highest weight representations of a given dimension. Not all these representations are irreducible however. In fact there are $(p-1)$ reducible representations of dimension p. This is the subject of the next theorem.

Theorem 6 *Let $k \in \{1, ..., p-1\}$ then*

1. *$W(p; \frac{2}{3}k - \frac{1}{3}p)$ has a $p-k$ dimensional sub-representation isomorphic to $W(p-k; -\frac{1}{3}(k+p))$.*

2. *The quotient representation is isomorphic to $W(k; \frac{2}{3}p - \frac{1}{3}k)$, i.e.*

$$W(p; \frac{2}{3}k - \frac{1}{3}p)/W(p-k; -\frac{1}{3}(k+p)) \simeq W(k; \frac{2}{3}p - \frac{1}{3}k) \tag{3.3.171}$$

Proof: Consider the representation $W(p, x)$. It might happen that $E.v_k = 0$ for some $k < p$. The subspace of $W(p, x)$ spanned by $F^l.v_k$ is then an invariant subspace. Let us now consider when this happens. Again we have to solve the equation $EF^k.v = 0$. This leads (as in the proof of the previous theorem) to the equation

$$k(j^2 + c - 2(k-1)j + \frac{2}{3}(k-1)(2k-1)) = 0 \tag{3.3.172}$$

This equation has three solutions for k, namely two we already know

$$\begin{aligned} k &= 0 \\ k &= p \end{aligned}$$

and a new one

$$k = \frac{1}{2}p + \frac{3}{2}x \tag{3.3.173}$$

This means that if p and x are such that $\frac{1}{2}p + \frac{3}{2}x$ is an integer between 1 and $p-1$ then

$$E.v_{\frac{1}{2}p+\frac{3}{2}x} = 0 \tag{3.3.174}$$

The sub-representation that can be built out of $v_{\frac{1}{2}p+\frac{3}{2}x}$ is obviously a $p-k$ dimensional highest weight representation of dimension $p-k$. It is therefore equal to $W(p-k; \tilde{x})$ for some $\tilde{x}$. Since C acts on the sub-representation $W(p-k; \tilde{x})$ as the same multiple of unity as on $W(p; \frac{2}{3}k - \frac{1}{3}p)$ and since $H(F^k v) = j(p; \frac{2}{3}k - \frac{1}{3}p) - 2k$ we see that in order to determine $\tilde{x}$ we have to solve the equations

$$\begin{aligned} c(p; \frac{2}{3}k - \frac{1}{3}p) &= c(p-k; \tilde{x}) \\ j(p; \frac{2}{3}k - \frac{1}{3}p) - 2k &= j(p-k; \tilde{x}) \end{aligned}$$

Inserting the expression for $c(p,x)$ and $j(p,x)$ given in the previous theorem we find $\tilde{x} = -\frac{1}{3}(k+p)$. For the second part a similar argument shows that we have to solve $\tilde{x}$ from the equations

$$\begin{aligned} c(p; \frac{2}{3}k - \frac{1}{3}p) &= c(k;\tilde{x}) \\ j(p; \frac{2}{3}k - \frac{1}{3}p) &= j(k;\tilde{x}) \end{aligned}$$

Again using the explicit expressions for c and j we find $\tilde{x} = \frac{2}{3}p - \frac{1}{3}k$. This proves the theorem .

The most important representations from a physical point of view are the unitary ones. The existence of these unitary representations is our next subject.

Define an anti-involution ω on $W_3^{(2)}$ by

$$\begin{aligned} \omega(E) &= F \\ \omega(F) &= E \\ \omega(H) &= H \\ \omega(C) &= C \end{aligned} \tag{3.3.175}$$

It is easy to check that ω is an algebra endomorphism, i.e. that it preserves the relations of $W_3^{(2)}$.

Lemma 5 *For all positive integers p and real numbers x the representation $W(p;x)$ carries a unique bilinear symmetric form $\langle .,. \rangle$ such that*

1. $\langle v, v \rangle = 1$
2. $\langle .,. \rangle$ *is contravariant w.r.t. ω, i.e.* $\langle A \cdot v_l, v_k \rangle = \langle v_l, \omega(A) \cdot v_k \rangle$ *for all* $A \in \bar{W}_3^{(2)}$.

Proof: Define the bilinear symmetric form $\langle .,. \rangle$ by putting $\langle v, v \rangle = 1$ and

$$\langle v_l, v_k \rangle = \delta_{lk} \prod_{i=1}^{k} A(i) \tag{3.3.176}$$

for $l, k \in \{0, 1, ..., p-1\}$, where

$$A(k) = k(j^2 + c) - 2k(k-1)j + \frac{2}{3}k(k-1)2k-1) \tag{3.3.177}$$

We have to check contravariance. Obviously it suffices to check this for the generators. For example take $A = F$, then

$$\langle F \cdot v_l, v_k \rangle = \delta_{l+1,k} \prod_{i=1}^{k} A(i) \tag{3.3.178}$$

On the other hand $\langle v_l, \omega(F) \cdot v_k \rangle = \langle v_l, E \cdot v_k \rangle$ which by lemma 1 is equal to

$$\begin{aligned} \langle v_l, E \cdot v_k \rangle &= \langle v_l, v_{k-1} \rangle A(k) \\ &= \delta_{l+1,k} \prod_{i=1}^{k-1} A(i) A(k) \\ &= \delta_{l+1,k} \prod_{i=1}^{k} A(i) \end{aligned}$$

which proves contravariance for F. The proof for the other generators is similar. Uniqueness follows by induction on k .

So the representations $W(p;x)$ all carry bilinear symmetric forms such that the basis vectors v_k are orthogonal. This is not enough for unitarity however for we still need to check if the bilinear form is positive definite. This is the subject of the next theorem.

Theorem 7 *The representation $W(p,x)$ is unitary if and only if*

$$x > \frac{1}{3}p - \frac{2}{3} \tag{3.3.179}$$

Proof: We have to find out which $W(p,x)$ have the property that the vectors v_l all have positive norm $\langle v_l, v_l \rangle > 0$ for $l = 1, ..., p-1$. Now, by definition

$$\langle v_l, v_l \rangle = \prod_{k=1}^{l} (k(j^2 + c) - 2k(k-1)j + \frac{2}{3}k(k-1)(2k-1)) \tag{3.3.180}$$

For unitarity we have to demand that this is > 0 for all $l = 1, 2, ..., p-1$. This can only be the case if $A(k) > 0$ for all $k = 1, ..., p-1$ separately. Inserting the expressions for $j(p;x)$ and $c(p;x)$ into this equation we find that it reduces to

$$x > \frac{2}{3}k - \frac{1}{3}p \tag{3.3.181}$$

This must be true for all $k = 1, ..., p-1$ which is only the case if

$$x > \frac{1}{3}p - \frac{2}{3} \tag{3.3.182}$$

This proves the theorem .



Figure 3.1: The location of the reducible (depicted by dots) and unitary representations.

Note that from theorems 2 and 3 follows that the quotient representation of a reducible rep w.r.t. its invariant subspace is a unitary irreducible representation. This concludes our presentation of the representation theory of $\bar{W}_3^{(2)}$.

3.3.2 Induced represention

In the previous section we constructed the representation theory of $\bar{W}_3^{(2)}$ in a way that is in principle completely analogous to the way in which one constructs the representation theory of sl_2. This of course was possible due to the special structure of this algebra. However, one look at the finite W algebras listed in the appendix of this chapter reveals that such an analysis would be extremely cumbersome for other finite W algebras because of their very complicated (nonlinear) structure. Also one would hardly be able to make any general statements which means that one would have to construct the representation theory case by case.

In the previous section however we did not use one crucial piece of information, namely that finite W algebras are reductions of simple Lie algebras. It might by possible to let representations of these Lie algebras 'descend' to finite W representations. Let us explain how this works in general. Let g again be a simple Lie algebra. We have seen that finite W algebras are the zeroth cohomology of the complex $(\Omega; d)$, where

$\Omega = \mathcal{U}g \otimes Cl(g_+ \oplus g_+^*)$, $d(X) = [Q, X]$ and Q is given by (3.2.141). The Clifford algebra $Cl(g_+ \oplus g_+^*)$ is finite dimensional and has a unique irreducible module given by

$$L_{gh} = Cl(g_+ \oplus g_+^*)/Cl(g_+ \oplus g_+^*).g_+ \tag{3.3.183}$$

Essentially we can describe L_{gh} as follows: it is spanned by

$$\{[c_{\alpha_{i_1}} \dots c_{\alpha_{i_k}}]\}_{\alpha_{i_1} \leq \dots \leq \alpha_{i_k}} \tag{3.3.184}$$

where we have defined some ordening on the basis $\{t_\alpha\}$ of g_+. Obviously the dimension of L_{gh} is $2^{|g_+|}$, where $|g_+| = \dim(g_+)$. We can introduce a grading on L_{gh} by putting

$$\deg([c_{\alpha_{i_1}} \dots c_{\alpha_{i_k}}]) = k \tag{3.3.185}$$

so we get

$$L_{gh} = \bigoplus_{k \geq 0} L_{gh}^{(k)} \tag{3.3.186}$$

We find therefore that any representation L_Λ of g induces a unique representation $L_\Lambda(\Omega) = L_\Lambda \otimes L_{gh}$ of Ω and furthermore that this representation is graded

$$L_\Lambda(\Omega) = \bigoplus_{k \geq 0} L_\Lambda(\Omega)^{(k)} \tag{3.3.187}$$

where $L_\Lambda(\Omega)^{(k)} = L_\Lambda \otimes L_{gh}^{(k)}$.

Now the point is that Q turns $L_\Lambda(\Omega)$ into a complex

$$Q : L_\Lambda(\Omega)^{(n)} \longrightarrow L_\Lambda(\Omega)^{(n+1)} \tag{3.3.188}$$

and that the zeroth cohomology $H^0(\Omega; d)$ of the complex $(\Omega; d)$ has a natural action on the cohomology spaces $H^n(L_\Lambda(\Omega); Q)$ that turns them into $H^0(\Omega; d)$ representations. We can describe this action as follows: Let $[x] \in H^0(\Omega; d)$ and $[v] \in H^n(L_\Lambda(\Omega); Q)$ then $[x].[v] \equiv [x.v]$. It is a standard excercise to verify that this definition does not depend on the representatives x and v of $[x]$ and $[v]$ respectively, and also that $x.v \in \ker(Q) \cap L_\Lambda(\Omega)^{(n)}$ if $v \in \ker(Q) \cap L_\Lambda(\Omega)^{(n)}$. Now, since $H^0(\Omega; d)$ is nothing but the finite W algebra in question we conclude that the spaces $H^n(L_\Lambda(\Omega); Q)$ are finite W representations. This provides us with a functor from the category of g representations to the category of representations of the finite W algebra in question.

Even though this procedure is quite general and elegant it is not the one that we shall persue. The reason for this is that it is rather nontrivial to find g representations L_Λ that reduce to interesting finite W representations. Let us illustrate this in the example of $W_3^{(2)}$. Again choosing the basis $\{t_a\}$ as specified by (3.2.160) Q reads

$$Q = J^3 c_3 + (J^1 - 1)c_1 \tag{3.3.189}$$

Also $L_{gh}^{(0)} = \mathbb{C}$, $L_{gh}^{(1)} = \mathbb{C}[c_1] \oplus \mathbb{C}[c_3]$ and $L_{gh}^{(2)} = \mathbb{C}[c_1 c_3]$. Let us take L_Λ to be the fundamental representation of sl_3, then one can easily verify that $H^*(L_\Lambda(\Omega); Q) = 0$ which means that the fundamental representation of sl_3 reduces to the trivial representation of $W_3^{(2)}$. This in fact holds true for all irreducible representation L_Λ of sl_3 which is rather disappointing. In view of this we shall take a slightly different approach towards the construction of the representation theory of finite W algebras. In essence what we will show is that there exists an injective algebra homomorphism from any finite W algebra to the universal enveloping algebra of the grade 0 piece g_0 of g (of course this is a quantum version of the Miura map presented earlier). This means that any g_0 representation induces a representation of the finite W algebra. Since g_0 is just a semisimple Lie algebra its representation theory is well known.

Let us now make all this more precise. Denote by $X^{0,0}$ the component of an element $X \in \Omega_{red}$ of bidegree $(0,0)$, so that $W^{0,0}(\phi) = (-1)^p \phi_p$, then

$$[W^{0,0}(\phi), W^{0,0}(\phi')] = [W(\phi), W(\phi')]^{0,0} \tag{3.3.190}$$

and therefore $W(\phi) \to W(\phi)^{0,0}$ is a homomorphism of algebras. We now have the following important theorem [97] which is a quantum version of the generalized Miura map.

Theorem 8 *(Quantum Miura Transformation) The map $W(\phi) \to W(\phi)^{0,0}$, or, equivalently, the map $H^0(\Omega; d) \to H^0(\Omega; d)^{0,0}$, is an isomorphism of algebras.*

Proof: Now that we know the cohomology of $(\Omega; d)$, let us go back to the original double complex $(\Omega_{red}; d_0; d_1)$. Because there is only cohomology of degree 0, we know that the $E_\infty^{p,q}$ term of the spectral sequence associated to $(\Omega_{red}; d_0; d_1)$ must vanish unless $p + q = 0$. If we look at $d_0(\hat{J}^{\bar{\alpha}}) = f^{\beta\bar{\alpha}}_{\bar{\gamma}} \hat{J}^{\bar{\gamma}} c_\beta$, we see that $d_0(\hat{J}^{\bar{\alpha}}) = 0$ if and only if $\bar{\alpha} \in g_0^*$. From this it is not difficult, repeating arguments similar as those in the proof of theorem 4, to prove that the only nonvanishing piece of $\oplus_r E_1^{r,-r}$ is in $E_1^{0,0}$. This implies that $E_\infty^{p,q}$ is only nonzero for $p = q = 0$. Because $H^0(\Omega; d) = E_\infty^{0,0} = H^0(\Omega; d)^{0,0}$ as vector spaces, it follows that the map $H^0(\Omega; d) \to H^0(\Omega; d)^{0,0}$ can have no kernel, and is an isomorphism.

The quantum Miura map is an injective algebra homomorphism into the subalgebra Ω^0_{red} of Ω_{red} generated by $\{J^{\bar{\alpha}}\}_{t_{\bar{\alpha}} \in g_0}$. Since

$$[\hat{J}^{\bar{\alpha}}, \hat{J}^{\bar{\beta}}] = f^{\bar{\alpha}\bar{\beta}}_{\bar{\gamma}} \hat{J}^{\bar{\gamma}} \tag{3.3.191}$$

we find that Ω^0_{red} is isomorphic as an algebra to $\mathcal{U}g_0$. We have therefore indeed found the result that any finite W algebra can be embedded into $\mathcal{U}g_0$.

Since $\mathcal{U}(g_0)$ is abelian for the principal sl_2 embeddings, this implies that in those cases the quantum finite W algebras are also abelian, something which was already proven by Kostant [88]. To get some interesting novel structure, one should therefore consider nonprincipal sl_2 embeddings.

Again let us consider the example of $3 = 2 + 1$. The expressions for the quantum Miura transformation of this algebra are obtained from (3.2.161) by restricting these expressions to the bidegree $(0, 0)$ part. If we introduce $s = (\hat{J}^4 + 3\hat{J}^5)/4$, $h = (\hat{J}^5 - \hat{J}^4)/4$, $f = 2\hat{J}^6$ and $e = \hat{J}^2/2$, then h, e, f form an sl_2 Lie algebra, $[h, e] = e$ and $[h, f] = -f$ and $[e, f] = 2h$ while s commutes with everything. In terms of s and h, e, f, the quantum Miura transformation reads

$$\begin{aligned} C &= -\tfrac{4}{3}(h^2 + \frac{1}{2}ef + \frac{1}{2}fe) - \tfrac{4}{9}s^2 + \tfrac{4}{3}s - 1, \\ H &= 2h - \tfrac{2}{3}s + 1, \\ E &= -2(s - h - 1)e, \\ F &= \tfrac{2}{3}f. \end{aligned} \tag{3.3.192}$$

Notice that C contains the second Casimir of sl_2, which is what one would expect because it commutes with everything. Every $g_0 = sl_2 \oplus u(1)$ module gives, using the expressions (3.3.192), a module for the finite quantum algebra $W_3^{(2)}$. So if we have a representation of $sl_2 \oplus u(1)$ in terms of $n \times n$ matrices, we immediately get a representation of $\bar{W}_3^{(2)}$ in terms of $n \times n$ matrices.

An irreducible representation of $sl_2 \oplus u(1)$ is characterized by two numbers: the 'spin' j of the sl_2 representation and a $u(1)$ 'charge' s. It is then easy to verify that the (j, s) representation of $sl_2 \oplus u(1)$ gives rise to the representation $W(2j + 1; 1 - \frac{2}{3}s)$ of $W_3^{(2)}$. Note also that the map from $sl_2 \oplus u(1)$ representations to $W_3^{(2)}$ representations is surjective.

In the appendix we give explicit formulas for the quantum Miura transformations associated to the finite W algebras that can be obtained from sl_4.

3.3.3 Fock realizations

Using the quantum Miura transformation we can turn any Fock realization of g_0 into a Fock realization of the corresponding finite W algebra. In this section we briefly discuss how we can obtain Fock realizations for simple Lie algebras. We will then show in the example of $W_3^{(2)}$ that its finite dimensional representations can be realized as submodules of certain Fock modules.

As usual define an oscillator algebra O to be an algebra generated by operators $\{a_i, a_i^\dagger\}_{i=1}^n$ with the following commutation relations

$$[a_i, a_j] = [a_i^\dagger, a_j^\dagger] = 0 \ ; \ \ [a_i, a_j^\dagger] = \delta_{ij} \tag{3.3.193}$$

In physics terminology this algebra describes a collection of n uncoupled harmonic oscillators. The Hilbert space of such a system is isomorphic to the space $\mathbb{C}[z_1, \ldots, z_n]$ of polynomials in n variables, and (the quantum operators) a_i and $a_j^\dagger$ can be realized as

$$a_i = \frac{\partial}{\partial z_i} \quad \text{and} \quad a_i^\dagger = z_i \tag{3.3.194}$$

This is called the 'Fock realization' of the (quantum) harmonic oscillator algebra. Since any polynomial P can be reduced to a multiple of 1 by successive application of $\{a_i\}$ we can construct any polynomial Q out of P by the action of the universal enveloping algebra of the oscillator algebra. This means that the Fock realization is irreducible. A Fock realization of a Lie algebra g is an injective homomorphism from g to the universal enveloping algebra of the oscillator algebra.

As we shall need this later we now discuss how to obtain a Fock realization for an arbitrary simple Lie algebra g (see for details [92]).

Let G be a simple Lie group with Lie algebra g. Furthermore let B_- be its lower Borel subgroup and N_+ the group associated to the positive roots. Consider the coset space

$$B_-\backslash G = \{B_- g \mid g \in G\} \tag{3.3.195}$$

The triple $(G, B_-\backslash G, B_-)$ is a principal bundle with total space G, basespace $B_-\backslash G$ and structure group B_-. Let now $\chi : B_- \to \mathbb{C}$ be a 1-dimensional representation (a character) of the Lie group B_-, then we can construct the associated complex line bundle $\mathcal{L}_\chi$ over $B_-\backslash G$

$$\mathcal{L}_\chi = G \times \mathbb{C}/\sim \tag{3.3.196}$$

where $(b.g, \chi(b^{-1})z) \sim (g, z)$ for all $b \in B_-$. Denote the set of holomorphic global sections of this bundle by Γ_χ. The group G acts from the right on $\mathcal{L}_\chi$ as follows

$$[(g, z)] \cdot g' = [(gg', z)] \tag{3.3.197}$$

and this induces of course an action of G (and therefore of g) on Γ_χ.

Now let H and b_- be the Cartan subalgebra and negative Borel subalgebra of g respectively. The derivative of χ in the unit element

$$\phi_\chi(b) = \frac{d}{dt}\chi(e^{tx})|_{t=0} \qquad x \in g \tag{3.3.198}$$

is a one dimensional representation of b_-. However this implies that

$$0 = [\phi_\chi(h), \phi_\chi(e_{-\alpha})] = -\langle \alpha, h\rangle \phi_\chi(e_\alpha) \tag{3.3.199}$$

where $h \in H$ and $e_{-\alpha}$ is a rootvector of g, which means that $\phi_\chi(e_{-\alpha}) = 0$ for all $-\alpha \in \Delta_+$. Therefore ϕ_χ determines really an element Λ_χ of H^*. It is also clear that all elements of H^* can be obtained in this way.

It was already said above that Γ_χ is a representation space of g. In fact the Borel-Weil theorem states that for Λ_χ an dominant weight of g the space of holomorphic sections Γ_χ of $\mathcal{L}_\chi$ is isomorphic to the finite dimensional irreducible representation of g with highest weight Λ_χ. If Λ_χ is not a dominant weight then the complex line bundle $\mathcal{L}_\chi$ has no non-trivial holomorphic sections.

The construction that leads to the Fock realization of the Lie algebra g is very similar to the Borel-Weil construction. First one makes use of the fact that $B_-\backslash G$ possesses a canonical decomposition, called the Bruhat decomposition

$$B_-\backslash G = \bigcup_{w\in W} C_w \tag{3.3.200}$$

where W is the Weyl group of g. The spaces C_w are called 'Schubert cells' and are isomorphic to $\mathbb{C}^n$. Denote from now on the Schubert cel corresponding to the unit element of W by Y. Since $Y \subset B_-\backslash G$ the group G acts on Y by right multiplication. It can also be shown that Y is diffeomorphic to N_+. This diffeomorphism is defined by assigning to $[g] \in Y$ the unique element $x \in N_+$ such that $[x] = [g]$ (i.e. such that $x = bg$ for some $b \in B_-$). The right G-action on Y induces a right G-action on N_+ via this diffeomorphism. Let's from now on identify Y and N_+.

The set B_-N_+ is a submanifold of G and the triple (B_-N_+, Y, B_-) is again a principle bundle (in fact it is nothing but the restriction of the bundle $(G, B_-\backslash G, B_-)$ to the cell $Y \subset B_-\backslash G$. Let again χ be a character of B_- and let $\bar{\mathcal{L}}_\chi$ be the associated complex line bundle over Y. Denote the space of sections of this line bundle by $\bar{\Gamma}_\chi$. It is a standard result (and easy to prove) that there exists a 1-1 correspondence between $\bar{\Gamma}_\chi$ and the set $\bar{R}_\chi$ of holomorphic functions on B_-N_+ such that

$$f(b.g) = \chi^{-1}(b) f(g) \tag{3.3.201}$$

where $b \in B_-$ and $g \in N_+$. Since obviously the space of holomorphic functions on N_+ and $\bar{R}_\chi$ are isomorphic (a holomorphic function f on N_+ uniquely determines a holomorphic function $\hat{f}$ on B_-N_+ with the property (3.3.201) if we set $\hat{f}(b.g) = \chi^{-1}(b)f(g)$) we find that

$$\bar{\Gamma}_\chi \simeq Hol(Y) \tag{3.3.202}$$

In [92] Kostant showed that there exists a representation of the Lie algebra g of G in terms of first order smooth differential operators on the manifold Y. Since the space Y is isomorphic to $\mathbb{C}^{|\Delta_+|}$ where $|\Delta_+|$ is the number of positive roots of g we can choose a set of global coordinates $z \equiv \{z_\alpha\}_{\alpha\in\Delta_+}$ on Y. The representation

$$\sigma_\Lambda : g \to \text{Diff}(Y) \tag{3.3.203}$$

of g on the set of differential operators $Diff(Y)$ on Y is then given by

$$\sigma_\Lambda(x) = \xi(x) + h_\Lambda(x) \quad x \in g \tag{3.3.204}$$

where $\xi(x)$ is a Killing vector of the right G-action on Y

$$\left(\xi(x)f\right)(z) = \frac{d}{dt} f(z.e^{tx})|_{t=0} \quad z \in Y \tag{3.3.205}$$

and

$$\left(h_\Lambda(x)f\right)(z) = \langle \Lambda_*, Ad_z x\rangle f(z) \tag{3.3.206}$$

Here Λ_* is the trivial extension of $\Lambda \in H^*$ to g and f is some holomorphic function of Y. Note that in (3.3.206) we made use of the fact we have identified Y and N_+ or else Ad_z would not make sense.

The subspace $\mathbb{C}[\{z_\alpha\}]$ of $Hol(Y)$ is obviously invariant under $\sigma_\Lambda(g)$ which means that we have obtained Fock realizations of the Lie algebra g (one for each $\Lambda \in H^*$). Let us denote by F_Λ the pair $(\mathbb{C}[\{z_\alpha\}], \sigma_\lambda)$.

For sl_2 the Fock realizations σ_Λ have the form

$$\begin{aligned}
\sigma_\Lambda(e) &= \frac{d}{dz} \\
\sigma_\Lambda(f) &= (\Lambda,\alpha)z - z^2\frac{d}{dz} \\
\sigma_\Lambda(h) &= \frac{1}{2}(\Lambda,\alpha) - z\frac{d}{dz}
\end{aligned} \tag{3.3.207}$$

where α is the root of sl_2. The expressions for arbitrary sl_n can be found in [75] and will not be given here.

Using the results described above one is always able to immediately write down a Fock realization of the algebra g_0 (since it is essentially a direct sum of sl_k algebras). Then using the quantum Miura transformation one thus arrives at a Fock realization of the finite W algebra in question. Let us now explicitly do this in the example of $\bar{W}_3^{(2)}$. Inserting the expressions (3.3.207) into the Miura map (3.3.192) one finds

$$\begin{aligned}
\sigma_{\Lambda,s}(H) &= (\Lambda,\alpha) - \frac{2}{3}s + 1 - 2z\frac{d}{dz} \\
\sigma_{\Lambda,s}(E) &= 2(1 - s + \frac{1}{2}(\Lambda,\alpha))\frac{d}{dz} - 2z\frac{d^2}{dz^2} \\
\sigma_{\Lambda,s}(F) &= \frac{2}{3}(\Lambda,\alpha)z - \frac{2}{3}z^2\frac{d}{dz} \\
\sigma_{\Lambda,s}(C) &= -\frac{1}{3}(\Lambda,\alpha)^2 - \frac{2}{3}(\Lambda,\alpha) - \frac{4}{9}s^2 + \frac{4}{3}s - 1
\end{aligned} \tag{3.3.208}$$

(where we consider s to be a number). This realization is equal to the zero mode structure of the free field realization of the infinite $W_3^{(2)}$ algebra constructed in [93]. Note however that the derivation is completely different since in [93] (using standard methods) the expressions in terms of free fields were obtained by constructing the generators of the commutant of certain screening charges. Constructing this commutant is in general however rather cumbersome. The method we presented above is more direct and works for arbitrary embeddings (and realizations).

The Fock realizations of g are not all irreducible. In fact all finite dimensional irreducible representations of simple Lie algebras can be realized as submodules fo certain Fock realizations. Let us briefly describe how this works before we show that the same statement is true for finite W algebras.

Suppose $D \in Diff(Y)$ is a differential operator such that $[\sigma_\Lambda(x), D] = 0$ for all $x \in g$ then the subspace $V = \{v \in F_\Lambda \mid Dv = 0\}$ of F_Λ is an invariant subspace, i.e. $\sigma_\Lambda(g)V \subset V$. A 'complete' set of such operators can be constructed as follows: Let x be an element of n_+, then define the differential operator $\rho(x)$ by

$$(\rho(x)f)(z) = \frac{d}{dt}f(e^{-tx}z)|_{t=0} \tag{3.3.209}$$

where $f \in C^\infty(Y)$. Here we again used the isomorphism $Y \simeq N_+$ in order to define the left action of N_+ on Y. Now let $\{e_i\}_{i=1}^l$ be the simple rootvectors of g corresponding to the simple roots $\{\alpha_i\}_{i-1}^l$, then the differential operators

$$\rho(e_i)^{(\Lambda,\alpha_i)} \tag{3.3.210}$$

(called intertwiners) commute with $\sigma_\Lambda(x)$ for all $x \in g$ if Λ is an integral dominant

weight (i.e. if (Λ, α_i) is a positive integer). Furthermore the submodule

$$V = \{v \in F_\Lambda \mid \rho(e_i)^{(\Lambda,\alpha_i)+1} v = 0, \text{ for all } i\} \tag{3.3.211}$$

is isomorphic to the (unique) finite dimensional irreducible g representation L_Λ with highest weight Λ. The operators $s_i \equiv \rho(e_i)$ are usually called 'sreening charges' in physics literature.

Let us illustrate all this with an example. For sl_2 we have $\rho(e) = -\frac{d}{dz}$. It is easy to check that for $(\Lambda, \alpha) = m \in \mathbf{Z}_+$ the space $V = \{v \in \mathbb{C}[z] \mid (\frac{d}{dz})^{m+1} v = 0\}$ carries the $m+1$ dimensional irreducible multiplet of sl_2.

Since the intertwiners commute with all elements of $\mathcal{U}g_0$ they also commute with the Fock representants of the finite W algebra in question. Therefore the finite dimensional spaces (3.3.211) are also finite dimensional representation spaces of the finite W algebra. This provides us with Fock realizations of the finite dimensional representations of W algebras.

Let us again consider the case of $W_3^{(2)}$. Let $m = (\Lambda, \alpha)$ then the subspace $V = \{v \in \mathbb{C}[z] \mid (\frac{d}{dz})^{m+1} v = 0\}$ of the Fock representation $F_{\Lambda,s} = (\mathbb{C}[z], \sigma_{\Lambda,s})$ of $W_3^{(2)}$ then it is easy to see that $(V, \sigma_{\Lambda,s})$ is isomorphic as a $W_3^{(2)}$ representation to $W(m+1; 1 - \frac{2}{3}s)$.

3.4 Relation with Lie algebra cohomology

According to the general construction in this chapter one associates a finite W algebra to any triple (g, g_+, χ) where g is a simple Lie algebra, g_+ is the positive grade subspace w.r.t. some grading element δ and χ is a certain one dimensional representation (character) of g_+ (note that in principle one could consider the case where g_+ and χ are *any* nilpotent subalgebra and character respectively, not necessarily related to an sl_2 embedding. The analysis given in the previous sections still remains valid without alterations). We shall now briefly indicate the relation of these constructions with Lie algebra cohomology.

Recall the definition of Lie algebra cohomology: Let h be a Lie algebra and let (V, ϕ) be a representation of h (i.e. V is a vectorspace and ϕ is a Lie algebra homomorphism from h to the set of linear transformations on V). Consider the spaces

$$C^p = \mathrm{Hom}(\bigwedge^p h, V) \tag{3.4.212}$$

(where of course $C^0 = V$) and define

$$C = \bigoplus_{p=0}^{\infty} C^p \tag{3.4.213}$$

The Lie algebra h has a natural action on C^p defined as follows: Let $x \in h$ and $\alpha \in C^p$, then $x.\alpha \equiv \phi(x) \circ \alpha \in C^p$. One then defines a coboundary operator $d_p : C^p \to C^{p+1}$ by putting

$$\begin{aligned}(d_p\alpha)(x_1 \wedge \ldots \wedge x_{p+1}) &= \sum_{j=1}^{p+1}(-1)^{j+1}(x_j.\alpha)(x_1 \wedge \ldots \wedge \hat{x}_j \wedge \ldots \wedge x_{p+1}) \\ &\quad + \sum_{1\le i<jp+1}(-1)^{i+j}\alpha([x_i, x_j] \wedge \ldots \wedge \hat{x}_i \wedge \ldots \wedge \hat{x}_j \wedge \ldots \wedge x_{p+1})\end{aligned}$$

where $\alpha \in C^p$ and $x_i \in h$ for all $i = 1, \ldots, p+1$. It is now easy to check that $d_p \circ d_{p-1} = 0$ for all p and the cohomology derived from this complex

$$H^p(h; V) = \frac{\mathrm{Ker}(d_p : C^p \to C^{p+1})}{\mathrm{Im}(d_{p-1} : C^{p-1} \to C^p)} \tag{3.4.214}$$

is called the p^{th} Lie algebra cohomology of h with coefficients in V.

We will now slightly reformulate the above to make contact with BRST cohomology. For this note that

$$\mathrm{Hom}(\textstyle\bigwedge^p h, V) \simeq \bigwedge^p h^* \otimes V \tag{3.4.215}$$

where the isomorphism can be described as follows: Let $\alpha \in \mathrm{Hom}(\bigwedge^p h, V)$ then

$$\alpha \mapsto \sum_{i_1<\ldots i_p} < (e^*_{i_1} \wedge \ldots \wedge e^*_{i_p}) \otimes \alpha(e_{i_1} \wedge \ldots \wedge e_{i_p}) \tag{3.4.216}$$

where $\{e_i\}$ and $\{e^*_i\}$ are dual bases of h and h^* respectively, i.e. $e^*_i(e_j) = \delta_{ij}$. It is clear that the map (3.4.216) is independent of the chosen basis and that it defines an isomorphism.

It is now straightforward to calculate the action of d_p on the element $e^*_{i_1} \wedge \ldots \wedge e^*_{i_p}) \otimes v \in \bigwedge^p h \otimes V$. It reads

$$\begin{aligned}d((e^*_{i_1} \wedge \ldots \wedge e^*_{i_p}) \otimes v) &= \hat{d}(e^*_{i_1} \wedge \ldots \wedge e^*_{i_p}) \otimes v + \\ &\quad + \sum_i (e^*_i \wedge e^*_{i_1} \wedge \ldots \wedge e^*_{i_p}) \otimes \phi(e_i)v\end{aligned} \tag{3.4.217}$$

where the derivation $\hat{d} : \bigwedge^p h^* \to \bigwedge^{p+1} h^*$ is determined by

$$\hat{d}(e_i^*) = -\frac{1}{2} \sum_{jk} C_{jk}^i e_j^* \wedge e_k^* \tag{3.4.218}$$

and C_{jk}^i are the structure constants of h in the basis $\{e_i\}$.

Now, having chosen a certain basis $\{e_i\}$ in h we can interpret elements of $\bigwedge^p h$ as (formal) polynomials of anticommuting variables $\{\theta_i\}$ with coefficients in V, i.e. if

$$\alpha = \sum_{i_1 < \ldots < i_p} e_{i_1}^* \wedge \ldots \wedge e_{i_p}^* \otimes v^{i_1 \ldots i_p} \tag{3.4.219}$$

(where $v^{i_1 \ldots i_p} \in V$) then the polynomial in terms of the Grassman variables θ_i associated to it is given by

$$\alpha = \sum_{i_1 < \ldots < i_p} v^{i_1 \ldots i_p} \theta_{i_1} \ldots \theta_{i_p} \tag{3.4.220}$$

The advantage of the description of Lie algebra cohomology in terms of Grassman variables is that the operator

$$d = \bigoplus_p d_p : C \to C \tag{3.4.221}$$

takes on the following simple form

$$d = \sum_i \phi(e_i)\theta_i - \frac{1}{2} \sum_{ijk} C_{jk}^i \theta_j \theta_k \frac{\partial}{\partial \theta} \tag{3.4.222}$$

as can easy be checked using formula (3.4.217). Note that θ_i and $\frac{\partial}{\partial \theta_j}$ satisfy canonical anti commutation relations. The relation between Lie algebra cohomology and BRST cohomology is now clear: Letting $h = g_+$ and denoting

$$c_i = \theta_i \quad \text{and} \quad b^i = \frac{\partial}{\partial \theta_i} \tag{3.4.223}$$

we find that the Lie algebra cohomology of g_+ with coefficients in the universal enveloping algebra of g is equal to the cohomology of the operator d_1 (in the notation of the previous sections). Finite W algebras therefore do not correspond precisely to Lie algebra cohomology due to the fact that the BRST operator involves another term, i.e. $d = d_0 + d_1$. At this point we can only speculate as to the relevance of finite W algebras to the theory of Lie algebras but it may be that this role is an important one. In order to see this recall that Lie algebra cohomology arises in the theory of Lie groups as de

Rham cohomology of left invariant vectorfields and also through the theory of group extensions [87]. Finite W algebras may therefore correspond globally (i.e. on the group level) with a 'deformation' of de Rham theory. Whether this is true and what this new geometric theory is is not clear at this time but we will come back to this in future publications.

Conclusion

We have seen in this chapter that the theory of W algebras has a natural finite counterpart. The classical theory, which is a theory of Poisson reductions of the very important Kirillov Poisson structure, lead to a set of Poisson algebras (eqn. (3.1.72)) which are in general nonlinear but finitely generated. In the second part of the chapter we have seen that all these algebras can be quantized by the BRST formalism. A completely algorithmic method has been presented to construct the generators of the quantum finite W algebra and to calculate their relations. In the last section we have developed the representation theory of quantum finite W algebras. It was shown that this can be done easily by the so called 'quantum Miura transformation'. In this way it was also possible to obtain realizations of all finite W algebras in terms a finite collection of harmonic oscillators.

Appendix

In this appendix we discuss the finite W algebras that can be obtained from sl_4. Using the standard constructions developed in this chapter we calculate their relations and the quantum Miura maps.

$4 = 2 + 1 + 1$

The basis of sl_4 we use to study this quantum algebra is

$$r_a t_a = \begin{pmatrix} \frac{1}{2}r_{10} - \frac{1}{8}r_8 - \frac{1}{8}r_9 & r_{11} & r_{12} & r_{15} \\ r_5 & \frac{3}{8}r_8 - \frac{1}{8}r_9 & r_7 & r_{14} \\ r_4 & r_6 & -\frac{1}{8}r_8 + \frac{3}{8}r_9 & r_{13} \\ r_1 & r_2 & r_3 & -\frac{1}{2}r_{10} - \frac{1}{8}r_8 - \frac{1}{8}r_9 \end{pmatrix}. \quad (3.4.224)$$

The sl_2 embedding is given by $t_+ = t_{15}$, $t_0 = t_{10}$ and $t_- = \frac{1}{2}t_1$. The nilpotent algebra g_+ is spanned by $\{t_{13}, t_{14}, t_{15}\}$, g_0 by $\{t_4, \ldots, t_{12}\}$ and g_- by $\{t_1, t_2, t_3\}$. The d_1 cohomology

of Ω_{red} is generated by $\hat{J}^a$ for $a = 1, \ldots, 9$. Representatives that are exactly d-closed, are given by $W(\hat{J}^a) = \hat{J}^a$ for $a = 4, \ldots, 9$, and by

$$\begin{aligned} W(\hat{J}^1) &= \hat{J}^1 + \hat{J}^4\hat{J}^{12} + \hat{J}^5\hat{J}^{11} + \tfrac{1}{4}\hat{J}^{10}\hat{J}^{10} + \tfrac{\hbar}{2}\hat{J}^{10}, \\ W(\hat{J}^2) &= \hat{J}^2 + \hat{J}^6\hat{J}^{12} + \tfrac{1}{2}\hat{J}^8\hat{J}^{11} + \tfrac{1}{2}\hat{J}^{10}\hat{J}^{11} + \hbar\hat{J}^{11}, \\ W(\hat{J}^3) &= \hat{J}^3 + \hat{J}^7\hat{J}^{11} + \tfrac{1}{2}\hat{J}^9\hat{J}^{12} + \tfrac{1}{2}\hat{J}^{12}\hat{J}^{10} + \tfrac{\hbar}{2}\hat{J}^{12}. \end{aligned} \tag{3.4.225}$$

Introduce a new basis of fields as follows

$$\begin{aligned} U &= \tfrac{1}{4}(W(\hat{J}^8) + W(\hat{J}^9)), \\ H &= \tfrac{1}{4}(W(\hat{J}^8) - W(\hat{J}^9)), \\ F &= -W(\hat{J}^7), \\ E &= -W(\hat{J}^6), \\ G_1^- &= -W(\hat{J}^3), \\ G_1^+ &= W(\hat{J}^2), \\ G_2^- &= W(\hat{J}^5), \\ G_2^+ &= W(\hat{J}^4), \\ C &= W(\hat{J}^1) + \tfrac{1}{2}EF + \tfrac{1}{2}FE + H^2 + \tfrac{1}{2}U^2 + 2\hbar U. \end{aligned} \tag{3.4.226}$$

If we compute the commutators of these expressions, we find that C commutes with everything, $\{E, F, H\}$ form a sl_2 subalgebra and $G_i^\pm$ are spin $\frac{1}{2}$ representations for this sl_2 subalgebra. U represents an extra $u(1)$ charge. The nonvanishing commutators, with $\hbar$ dependence, are

$$\begin{aligned} [E, F] &= 2\hbar H, \\ [H, E] &= \hbar E, \\ [H, F] &= -\hbar F, \\ [U, G_1^\pm] &= \hbar G_1^\pm, \\ [U, G_2^\pm] &= -\hbar G_2^\pm, \\ [H, G_i^\pm] &= \pm\tfrac{\hbar}{2} G_i^\pm, \\ [E, G_i^-] &= \hbar G_i^+, \\ [F, G_i^+] &= \hbar G_i^-, \\ [G_1^+, G_2^+] &= -2\hbar E(U + \hbar), \\ [G_1^-, G_2^-] &= 2\hbar F(U + \hbar), \\ [G_1^+, G_2^-] &= \hbar(-C + EF + FE + 2H^2 + \tfrac{3}{2}U^2 + 2HU) + \hbar^2(2H + 3U), \\ [G_1^-, G_2^+] &= \hbar(C - EF - FE - 2H^2 - \tfrac{3}{2}U^2 + 2HU) + \hbar^2(2H - 3U). \end{aligned} \tag{3.4.227}$$

Let us also present the quantum Miura transformation for this algebra. In this case, $g_0 = sl_3 \oplus u(1)$. Standard generators of g_0 can be easily identified. A generator of the $u(1)$ is $s = \frac{1}{2}\hat{J}^8 + \frac{1}{2}\hat{J}^9 + 2\hat{J}^{10}$, and the sl_3 generators are $e_1 = \hat{J}^5$, $e_2 = \hat{J}^6$, $e_3 = \hat{J}^4$, $f_1 = \hat{J}^{11}$, $f_2 = \hat{J}^7$, $f_3 = \hat{J}^{12}$, $h_1 = -\frac{1}{2}\hat{J}^8 + \frac{1}{2}\hat{J}^{10}$ and $h_2 = \frac{1}{2}\hat{J}^8 - \frac{1}{2}\hat{J}^9$. The convention is such that the commutation relations between $\{e_i, f_i, h_i\}$ are the same as those of corresponding matrices defined by

$$a_i e_i + b_i f_i + c_i h_i = \hbar \begin{pmatrix} c_1 & a_1 & a_3 \\ b_1 & c_2 - c_1 & a_2 \\ b_3 & b_2 & -c_2 \end{pmatrix}. \tag{3.4.228}$$

The quantum Miura transformation reads

$$\begin{aligned} U &= \tfrac{1}{6}(s - 2h_2 - 4h_1), \\ H &= \tfrac{1}{2}h_2, \\ F &= -f_2, \\ E &= -e_2, \\ G_1^- &= -f_2 f_1 - \tfrac{1}{3}(s - 2h_2 - h_1 + 3\hbar) f_3, \\ G_1^+ &= e_2 f_3 + \tfrac{1}{3}(s + h_2 - h_1 + 3\hbar) f_1, \\ G_2^- &= e_1, \\ G_2^+ &= e_3, \\ C &= (\tfrac{1}{24}s^2 + \tfrac{\hbar}{2}s) + \tfrac{1}{2}(e_1 f_1 + f_1 e_1 + e_2 f_2 + f_2 e_2 + e_3 f_3 + f_3 e_3) \\ &\quad + \tfrac{1}{3}(h_1^2 + h_1 h_2 + h_2^2). \end{aligned} \tag{3.4.229}$$

In C we again recognize the second casimir of sl_3. It is a general feature of finite W algebras that they contain a central element C, whose Miura transform contains the second casimir of g_0. C is the finite counterpart of the energy momentum tensor that every infinite W algebra possesses.

$4 = 2 + 2$

A convenient basis to study this case is

$$r_a t_a = \begin{pmatrix} \frac{r_6}{4} + \frac{r_{10}}{2} & \frac{r_5}{2} + \frac{r_8}{2} & r_{12} & r_{14} \\ \frac{r_7}{2} + \frac{r_9}{2} & -\frac{r_6}{4} - \frac{r_{11}}{2} & r_{13} & r_{15} \\ r_1 & r_2 & \frac{r_6}{4} - \frac{r_{10}}{2} & \frac{r_5}{2} - \frac{r_8}{2} \\ r_3 & r_4 & \frac{r_7}{2} - \frac{r_9}{2} & -\frac{r_6}{4} + \frac{r_{11}}{2} \end{pmatrix}. \tag{3.4.230}$$

The sl_2 embedding is given by $t_+ = t_{12} + t_{15}$, $t_0 = t_{10} - t_{11}$ and $t_- = \frac{1}{2}(t_1 + t_4)$. The subalgebra g_+ is spanned by $\{t_{12}, \ldots, t_{15}\}$, g_0 by $\{t_5, \ldots, t_{11}\}$ and g_- by $\{t_1, \ldots, t_4\}$.

The d_1 cohomology of Ω_{red} is generated by $\hat{J}^1, \ldots, \hat{J}^7$. The d-closed representatives are $W(\hat{J}^a) = \hat{J}^a$ for $a = 5, 6, 7$, and

$$\begin{aligned}
W(\hat{J}^1) &= \hat{J}^1 - \tfrac{1}{4}\hat{J}^5\hat{J}^9 + \tfrac{1}{4}\hat{J}^7\hat{J}^8 + \tfrac{1}{4}\hat{J}^8\hat{J}^9 + \tfrac{1}{4}\hat{J}^{10}\hat{J}^{10} + \tfrac{3\hbar}{4}\hat{J}^{10} - \tfrac{\hbar}{4}\hat{J}^{11}, \\
W(\hat{J}^2) &= \hat{J}^2 + \tfrac{1}{4}\hat{J}^5\hat{J}^{10} + \tfrac{1}{4}\hat{J}^5\hat{J}^{11} - \tfrac{1}{4}\hat{J}^6\hat{J}^8 + \tfrac{1}{4}\hat{J}^8\hat{J}^{10} - \tfrac{1}{4}\hat{J}^{10}\hat{J}^{11} + \tfrac{\hbar}{2}\hat{J}^8, \\
W(\hat{J}^3) &= \hat{J}^3 + \tfrac{1}{4}\hat{J}^6\hat{J}^9 - \tfrac{1}{4}\hat{J}^7\hat{J}^{10} - \tfrac{1}{4}\hat{J}^7\hat{J}^{11} + \tfrac{1}{4}\hat{J}^9\hat{J}^{10} - \tfrac{1}{4}\hat{J}^9\hat{J}^{11} + \tfrac{\hbar}{2}\hat{J}^9, \\
W(\hat{J}^4) &= \hat{J}^4 + \tfrac{1}{4}\hat{J}^5\hat{J}^9 - \tfrac{1}{4}\hat{J}^7\hat{J}^8 + \tfrac{1}{4}\hat{J}^8\hat{J}^9 + \tfrac{1}{4}\hat{J}^{11}\hat{J}^{11} + \tfrac{\hbar}{4}\hat{J}^{10} - \tfrac{3\hbar}{4}\hat{J}^{11}. \qquad (3.4.231)
\end{aligned}$$

To display the properties of the algebra as clearly as possible, we introduce a new basis of fields

$$\begin{aligned}
H &= \tfrac{1}{2}W(\hat{J}^6), \\
E &= -W(\hat{J}^7), \\
F &= -W(\hat{J}^5), \\
G^+ &= W(\hat{J}^3), \\
G^0 &= W(\hat{J}^1) - W(\hat{J}^4), \\
G^- &= -W(\hat{J}^2), \\
C &= W(\hat{J}^1) + W(\hat{J}^4) + \tfrac{1}{8}W(\hat{J}^6)W(\hat{J}^6) + \tfrac{1}{4}W(\hat{J}^5)W(\hat{J}^7) \\
&\quad + \tfrac{1}{4}W(\hat{J}^7)W(\hat{J}^5) + \tfrac{\hbar}{4}W(\hat{J}^6). \qquad (3.4.232)
\end{aligned}$$

Here, C is the by now familiar central element, $\{E, H, F\}$ form an sl_2 algebra and $\{G^+, G^0, G^-\}$ form a spin 1 representation with respect to this sl_2 algebra. The nonvanishing commutators are

$$\begin{aligned}
[E, F] &= 2\hbar H, \\
[H, E] &= \hbar E, \\
[H, F] &= -\hbar F, \\
[E, G^0] &= 2\hbar G^+, \\
[E, G^-] &= \hbar G^0, \\
[F, G^+] &= \hbar G^0, \\
[F, G^0] &= 2\hbar G^-, \\
[H, G^+] &= \hbar G^+, \\
[H, G^-] &= -\hbar G^-, \\
[G^0, G^+] &= \hbar(-CE + EH^2 + \tfrac{1}{2}EEF + \tfrac{1}{2}EFE) - 2\hbar^3 E, \\
[G^0, G^-] &= \hbar(-CF + FH^2 + \tfrac{1}{2}FFE + \tfrac{1}{2}FEF) - 2\hbar^3 F, \\
[G^+, G^-] &= \hbar(-CH + H^3 + \tfrac{1}{2}HEF + \tfrac{1}{2}HFE) - 2\hbar^3 H. \qquad (3.4.233)
\end{aligned}$$

Since $g_0 = sl_2 \oplus sl_2 \oplus u(1)$, the quantum Miura transformation expresses this algebra in term of generators $\{e_1, h_1, f_1\}$, $\{e_2, h_2, f_2\}$, s of g_0. The relation between these generators and the $\hat{J}^a$ are: $s = \hat{J}^{10} - \hat{J}^{11}$, $h_1 = \frac{1}{2}(\hat{J}^6 + \hat{J}^{10} + \hat{J}^{11})$, $h_2 = \frac{1}{2}(\hat{J}^6 - \hat{J}^{10} - \hat{J}^{11})$, $e_1 = \frac{1}{2}(\hat{J}^7 + \hat{J}^9)$, $e_2 = \frac{1}{2}(\hat{J}^7 - \hat{J}^9)$, $f_1 = \frac{1}{2}(\hat{J}^5 + \hat{J}^8)$ and $f_2 = \frac{1}{2}(\hat{J}^5 - \hat{J}^8)$. The commutation relations for these are $[e_1, f_1] = \hbar h_1$, $[h_1, e_1] = 2\hbar e_1$, $[h_1, f_1] = -2\hbar f_1$, and similar for $\{e_2, h_2, f_2\}$. For the quantum Miura transformation one then finds

$$\begin{aligned}
H &= \tfrac{1}{2}(h_1 + h_2), \\
E &= -e_1 - e_2, \\
F &= -f_1 - f_2, \\
G^+ &= \tfrac{1}{2}e_1h_2 - \tfrac{1}{2}e_2h_1 + \tfrac{1}{4}s(e_1 - e_2) + \hbar(e_1 - e_2), \\
G^0 &= f_1e_2 - f_2e_1 + \tfrac{1}{4}s(h_1 - h_2) + \hbar(h_1 - h_2), \\
G^- &= \tfrac{1}{2}f_1h_2 - \tfrac{1}{2}f_2h_1 - \tfrac{1}{4}s(f_1 - f_2) - \hbar(f_1 - f_2), \\
C &= (\tfrac{1}{8}s^2 + \hbar s) + \tfrac{1}{2}(e_1f_1 + f_1e_1 + e_2f_2 + f_2e_2) + \tfrac{1}{4}(h_1^2 + h_2^2). \qquad (3.4.234)
\end{aligned}$$

The infinite dimensional version of this algebra is one of the 'covariantly coupled' algebras that have been studied in [96]. The finite algebra (3.4.233) is almost a Lie algebra. If we assign particular values to C and to the second casimir $C_2 = (H^2 + \frac{1}{2}EF + \frac{1}{2}FE)$ of the sl_2 subalgebra spanned by $\{E, H, F\}$, then (3.4.233) reduces to a Lie algebra. For a generic choice of the values of C and C_2 this Lie algebra is isomorphic to $sl_2 \oplus sl_2$. An interesting question is, whether a similar phenomena occurs for different covariantly coupled algebras.

$4 = 3 + 1$

The last nontrivial nonprincipal sl_2 embedding we consider is $\underline{4}_4 \simeq \underline{3}_2 \oplus \underline{1}_2$. We choose yet another basis

$$r_a t_a = \begin{pmatrix} \frac{r_5}{12} - \frac{r_6}{3} & r_8 & r_{11} & r_{12} \\ r_4 & -\frac{r_5}{4} & r_{13} & r_{14} \\ \frac{r_3}{2} - \frac{r_9}{2} & r_{10} & \frac{r_5}{12} + \frac{r_6}{6} + \frac{r_7}{2} & r_{15} \\ r_1 & r_2 & \frac{r_3}{2} + \frac{r_9}{2} & \frac{r_5}{12} + \frac{r_6}{6} - \frac{r_7}{2} \end{pmatrix}, \qquad (3.4.235)$$

in terms of which the sl_2 embedding is $t_+ = t_{11} + t_{15}$, $t^0 = -3t_6 + t_7$ and $t_- = 2t_3$. The subalgebra g_+ is generated by $\{t_{11}, \ldots, t_{15}\}$, g_- is generated by $\{t_1, t_2, t_3, t_9, t_{10}\}$, and g_0 is generated by $\{t_4, \ldots, t_8\}$. The d_1 cohomology is generated by $\hat{J}^1, \ldots, \hat{J}^5$, and d-closed representatives are given by $W(\hat{J}^4) = \hat{J}^4$, $W(\hat{J}^5) = \hat{J}^5$, and

$$W(\hat{J}^1) = \hat{J}^1 + \tfrac{1}{6}\hat{J}^3\hat{J}^6 - \tfrac{1}{12}\hat{J}^6\hat{J}^3 + \tfrac{1}{4}\hat{J}^7\hat{J}^3 + \hat{J}^4\hat{J}^{10} + \tfrac{1}{4}\hat{J}^6\hat{J}^9 - \tfrac{1}{4}\hat{J}^7\hat{J}^9 - \tfrac{1}{3}\hat{J}^4\hat{J}^5\hat{J}^8$$

$$
\begin{aligned}
& -\tfrac{1}{3}\hat{J}^4\hat{J}^6\hat{J}^8 - \tfrac{1}{108}\hat{J}^6\hat{J}^6\hat{J}^6 + \tfrac{1}{12}\hat{J}^6\hat{J}^7\hat{J}^7 + \tfrac{31\hbar}{12}\hat{J}^4\hat{J}^8 - \tfrac{3\hbar}{4}\hat{J}^9 + \tfrac{5\hbar}{48}\hat{J}^6\hat{J}^6 \\
& +\tfrac{\hbar}{6}\hat{J}^7\hat{J}^6 - \tfrac{3\hbar}{16}\hat{J}^7\hat{J}^7 - \tfrac{3\hbar^2}{8}\hat{J}^7 - \tfrac{7\hbar^2}{24}\hat{J}^6, \\
W(\hat{J}^2) &= \hat{J}^2 - \tfrac{1}{2}\hat{J}^3\hat{J}^8 - \tfrac{1}{3}\hat{J}^5\hat{J}^{10} - \tfrac{1}{6}\hat{J}^6\hat{J}^{10} + \tfrac{1}{2}\hat{J}^7\hat{J}^{10} - \tfrac{1}{2}\hat{J}^8\hat{J}^9 + \tfrac{3\hbar}{2}\hat{J}^{10} + \tfrac{1}{9}\hat{J}^5\hat{J}^5\hat{J}^8 \\
& +\tfrac{1}{9}\hat{J}^5\hat{J}^6\hat{J}^8 + \tfrac{1}{36}\hat{J}^6\hat{J}^6\hat{J}^8 - \tfrac{1}{4}\hat{J}^7\hat{J}^7\hat{J}^8 - \hbar\hat{J}^5\hat{J}^8 - \tfrac{\hbar}{2}\hat{J}^6\hat{J}^8 - \tfrac{\hbar}{2}\hat{J}^7\hat{J}^8 + 2\hbar^2\hat{J}^8, \\
W(\hat{J}^3) &= \hat{J}^3 + \hat{J}^4\hat{J}^8 + \tfrac{1}{12}\hat{J}^6\hat{J}^6 + \tfrac{1}{4}\hat{J}^7\hat{J}^7 + \tfrac{\hbar}{2}\hat{J}^7 - \tfrac{\hbar}{2}\hat{J}^6. \qquad (3.4.236)
\end{aligned}
$$

We introduce a new basis

$$
\begin{aligned}
U &= \tfrac{1}{4}W(\hat{J}^5), \\
G^+ &= W(\hat{J}^4), \\
G^- &= W(\hat{J}^2), \\
S &= W(\hat{J}^1), \\
C &= W(\hat{J}^3) + \tfrac{1}{24}W(\hat{J}^5)W(\hat{J}^5) - \tfrac{\hbar}{2}W(\hat{J}^5). \qquad (3.4.237)
\end{aligned}
$$

In this case, the fields are not organized according to sl_2 representations, because the centralizer of this sl_2 embedding in sl_4 does not contain an sl_2. Again C is a central element, and the nonvanishing commutators are

$$
\begin{aligned}
[U, G^+] &= \hbar G^+, \\
[U, G^-] &= -\hbar G^-, \\
[S, G^+] &= \hbar G^+(-\tfrac{2}{3}C + \tfrac{20}{9}U^2 - \tfrac{43\hbar}{9}U + \tfrac{29\hbar^2}{27}), \\
[S, G^-] &= \hbar(\tfrac{2}{3}C - \tfrac{20}{9}U^2 + \tfrac{43\hbar}{9}U - \tfrac{29\hbar^2}{27})G^-, \\
[G^+, G^-] &= \hbar S - \tfrac{4\hbar}{3}CU + \tfrac{3\hbar^2}{4}C + \tfrac{88\hbar}{27}U^3 - \tfrac{17\hbar^2}{2}U^2 + \tfrac{25\hbar^2}{6}U. \qquad (3.4.238)
\end{aligned}
$$

This is the first example where the brackets are no longer quadratic, but contain third order terms. For the sake of completeness, let us also give the quantum Miura transformation for this algebra. We identify generators $\{e, f, h\}$,s_1,s_2 of $sl_2 \oplus u(1) \oplus u(1) = g_0$ via $f = \hat{J}^8$, $e = \hat{J}^4$, $h = \frac{1}{3}(\hat{J}^5 - \hat{J}^6)$, $s_1 = \frac{1}{3}(2\hat{J}^6 + \hat{J}^5)$ and $s_2 = \hat{J}^7$. The only nontrivial commutators between these five generators are $[e, f] = \hbar h$, $[h, e] = 2\hbar e$ and $[h, f] = -2\hbar f$. The quantum Miura transformation now reads

$$
\begin{aligned}
U &= \tfrac{1}{4}s_1 + \tfrac{1}{2}h, \\
G^+ &= e, \\
G^- &= (\tfrac{1}{4}s_1^2 + \tfrac{1}{2}s_1 h + \tfrac{1}{4}h^2 - \tfrac{1}{4}s_2^2 - \tfrac{\hbar}{2}(3s_1 + 3h + s_2) + 2\hbar^2)f, \\
S &= -\tfrac{1}{12}e(8s_1 + 4h - 31\hbar)f - \tfrac{1}{108}(s_1 - h)^3 + \tfrac{1}{12}(s_1 - h)s_2^2 + \tfrac{5\hbar}{48}(s_1 - h)^2 \\
& +\tfrac{\hbar}{6}s_2(s_1 - h) - \tfrac{3\hbar}{16}s_2^2 - \tfrac{3\hbar^2}{8}s_2 - \tfrac{7\hbar^2}{24}(s_1 - h), \\
C &= (\tfrac{1}{2}ef + \tfrac{1}{2}fe + \tfrac{1}{4}h^2) + (\tfrac{1}{4}s_2^2 + \tfrac{\hbar}{2}s_2) + (\tfrac{1}{8}s_1^2 - \hbar s_1). \qquad (3.4.239)
\end{aligned}
$$

This completes our list of finite quantum W algebras from sl_4.

Chapter 4

W algebras from affine Lie algebras

In this chapter we develop, both classically and quantum mechanically, the point of view that CFT's with W symmetry are reductions of theories with affine symmetry. Examples of such theories are the well known Wess-Zumino-Witten models describing strings moving on group manifolds [25]. Due to this correspondence theories with affine symmetry obtain a universal status and many properties of quantum field theories with W symmetry can be derived by pulling down the properties of the theory with affine symmetry. A classic example of this was the discovery by Polyakov [50] that the 2 dimensional induced gravity action which appears in noncritical string theory has, in the chiral (or light cone) gauge, a 'hidden' affine $sl(2)$ symmetry. Motivated by this it was shown [68] that the induced gravity action, sometimes also called 'Liouville gravity', is nothing but a gauged version of the WZW model associated to $sl(2;\mathbf{R})$. As we have seen in chapter 2 this corresponds on an algebraic level to a realization of the Virasoro algebra (which is the symmetry algebra of the Liouville theory) as a reduction of the affine $sl(2;\mathbf{R})$ algebra. The consequences of this correspondence between the WZW model and the Liouville action are quite far reaching. For example it relates the correlation functions in chiral 2D gravity to those in an $sl(2;\mathbf{R})$ WZW model and it also gives a handle on the study of renormalization effects in 2D gravity [43].

Recently there has also been a great deal of interest in theories which are called 'higher spin generalizations of 2D gravity'. These higher spin gravities, which are generalizations of ordinary gravity in 2 dimensions, arise when one reduces an $sl(N;\mathbf{R})$ WZW model instead of an $sl(2;\mathbf{R})$ model. The theories which one obtains after reduction turn out to be Toda field theories which have been studied extensively within the context of integrable systems. Motivated by the $sl(2;\mathbf{R})$ case one interprets Toda field theory as a generalization of 2D induced gravity in the chiral gauge.

The classical versions of the covariant gravity theories that reduce in the chiral gauge to Toda theories have been constructed recently in [69]. The physical and geometrical interpretation of these generalized 2D gravity theories remains a mistery however.

On the algebraic level the reduction from $sl(N;\mathbf{R})$ WZW to Toda field theory is a reduction from an affine $sl(N;\mathbf{R})$ algebra to W_N. This lead to the remarkable insight that Toda field theories, which have been around for a long time, have as their symmetry algebras the nonlinear W_N algebras (this was realized independently in [81]). of course it was well known that Toda field theories are conformally invariant for this follows immediately from their description as dimensional reductions of selfdual Yang-Mills theories [17]. In this description one shows that the SDYM equations together with the constraint of spherical symmetry are equivalent to the Toda field equations of motion. However, it was not known that $sl(N;\mathbf{R})$ Toda field theory was in fact invariant under the higher spin extension of conformal symmetry known as W_N. We see that W algebras may also have applications in Yang-Mills theory.

In principle the generators of rotations with respect to which one defines sperical symmetry are not uniquely defined. In fact every $SU(2)$ embedding into the gauge group of the Yang-Mills system can be seen as generating rotations. The standard Toda field theories with W_N symmetry are related to the so called 'principal $SU(2)$ embeddings'. In [3] the problem of constructing the field theories associated to the other $SU(2)$ embeddings was addressed. This lead to a whole new class of generalized Toda field theories all of which are conformally invariant.

In view of the fact that ordinary Toda field theories are invariant under W extensions of conformal symmetry it is natural to ask whether this is also true for the generalized Toda theories. The answer to this question is affirmative and in this chapter we shall construct these W algebras. Again they will be reductions of affine Lie algebras and in analogy with the reductions that lead generalized Toda theories the different reductions are associated to the different $sl(2)$ embeddings into the gauge group. The class of W algebras we thus obtain is very large and contains many known examples.

This chapter consists of 2 parts. The first part deals with the classical theory. The classical W algebras thus obtained have, apart from the applications described above, also great relevance to the theory of integrable systems and solitary wave physics. In fact they are nothing but the second Hamiltonian structures of new and highly complicated but integrable hierarchies of evolution equations that generalize the KdV hierarchy. We shall be taking a look at some of these in the next chapter.

The second part of this chapter deals with the quantization of the classical W algebras constructed in the first part. For this we again use the BRST formalism which is ideally suited for jobs like this. In the entire chapter we continually emphasize the formal similarity between the theory of finite W algebras presented in the previous chapter and the theory of infinite W algebras. As we mentioned at the beginning of the previous chapter this is completely analogous to the theory of loop groups and algebras over finite dimensional simple Lie algebras.

4.1 Classical W algebras

In this section we develop the classical theory of W algebras. Within the context of integrable systems classical W algebras have a longer history than in conformal field theory. There they were known as Gelfand-Dickii Poisson brackets on spaces of pseudodifferential operators [26]. An example of a Gelfand-Dickii bracket is the well known second Hamiltonian structure of the KdV hierarchy. This bracket, which is the GD bracket associated to sl_2 is known to be nothing but the Virasoro algebra. The classical W_N algebras are just the GD bracket algebras associated to sl_N.

It was shown by Drinfeld and Sokolov [66] that the GD Poisson structure associated to a Lie algebra g is a reduction of the Kirillov Poisson algebra associated to the affine Lie algebra over g. This reduction, which is just Poisson reduction in infinitely many dimensions, is nowadays called Drinfeld-Sokolov reduction. It was the first evidence supporting the claim that many W algebras (perhaps all) can be viewed as reductions of some sort of Lie algebras. In this first section we generalize the (classical) construction by Drinfeld and Sokolov to include a large number of W algebras other than W_N. We also generalize the well known Miura map and present an elegant algebraic description of classical W algebras. In doing this we shall benefit greatly from the theory developed in the previous chapter since many of the statements that we have to prove here have already been proved in the finite case. We shall therefore refer heavily to the previous chapter.

4.1.1 Generalized Drinfeld-Sokolov reduction

Let $\{t_a\}$ be a basis of the simple Lie algebra g. Furthermore let $\mathcal{F}$ be the algebra of (classical) fields generated by $\{J^a(z)\}$ (where the parameter z is thought of as parametrizing S^1, or equivalently $U(1)$), i.e. $\mathcal{F}$ is defined inductively as follows:

1. $J^a(z)$ is an element of $\mathcal{F}$ for all a.
2. If $A(z), B(z) \in \mathcal{F}$ then $\alpha A(z) + \beta B(z) \in \mathcal{F}$.
3. If $A(z) \in \mathcal{F}$ then $\frac{dA}{dz}(z) \in \mathcal{F}$.
4. If $A(z), B(z) \in \mathcal{F}$ then $(AB)(z) = A(z)B(z) \in \mathcal{F}$.

Note that since we are still dealing with classical fields the multiplication is abelian, i.e. $(AB)(z) = (BA)(z)$. The fields $J^a(z)$ are to be considered as functionals on the affine Lie algebra $\bar{g} = Map(S^1; g) \oplus \mathbb{C}c$ over g: Any element $y \in Map(S^1; g)$ can be written as $y = y^a t_a$ where y^a are smooth maps from S^1 to the real numbers. The functional $J^a(z)$

is then defined by

$$J^a(z)(y) = y^a(z) \tag{4.1.1}$$

Define $Fun(\bar{g})$ to be the set of functionals on $\bar{g}$ of the form

$$F = \int_{S^1} dz\ A(z) \tag{4.1.2}$$

where $A(z) \in \mathcal{F}$. Obviously $Fun(\bar{g})$ is a subset of the space of smooth functions on $\bar{g}$.

$Fun(\bar{g})$ is a Poisson algebra w.r.t. the Kirillov Poisson structure on $\bar{g}$ which is given by

$$\{F, G\}(y) = (([\partial + y, \mathrm{grad}_y F], \mathrm{grad}_y G)) \tag{4.1.3}$$

where for $y_1, y_2 \in \bar{g}$

$$((y_1, y_2)) = \int_{S^1} dz\ (y_1(z), y_2(z)) \tag{4.1.4}$$

is the Killing form on $\bar{g}$ and $\mathrm{grad}_y F \in \bar{g}$ is defined by

$$\frac{d}{d\epsilon} F(y + \epsilon y_2)|_{\epsilon=0} = ((\mathrm{grad}_y F, y_2)) \tag{4.1.5}$$

for all $y_2 \in \bar{g}$. The commutator bracket $[.,.]$ in (4.1.3) denotes the Lie bracket of g, *not* of $\bar{g}$ which involves an extra central term. This is why there appears a $\partial \equiv \frac{d}{dz}$ in (4.1.41). Had we taken $[.,.]$ to be the Lie bracket in $\bar{g}$ then the right hand side of (4.1.3) would have been the usual expression for the Kirillov bracket (i.e. without the $\frac{d}{dz}$). In terms of the functionals $J^a(z)$ the bracket (4.1.3) reads

$$\{J^a(z), J^b(w)\} = f_c^{ab} J^c(w)\delta(z-w) + k g^{ab}\delta'(z-w) \tag{4.1.6}$$

where g^{ab} is the inverse of $g_{ab} = (t_a, t_b)$ (as in the previous chapter $(.,.)$ denotes the Cartan-Killing form on g), $[t_a, t_b] = f_{ab}^c t_c$ and k is the 'central element'. Note that strictly speaking the right hand side of this equation is no longer an element of $\mathcal{F}$ due to the delta functions, i.e. the Poisson bracket of two elements of $\mathcal{F}$ takes one outside of $\mathcal{F}$. This however is a well known phenomenon and the usual trick to avoid it is by considering so called 'smeared fields'. This however will nowhere be essential to us and we shall therefore ignore this subtlety from now on.

The algebra (4.1.6) is well known to arise as the current algebra of the so called WZW model. It should be compared to the Poisson relation

$$\{J^a, J^b\} = f_c^{ab} J^c \tag{4.1.7}$$

which we used as the starting point for the theory of finite W algebras in the previous chapter.

The algebra $\mathcal{F}$ contains fields which form Virasoro algebras. An example of such a Virasoro field is the so called (classical) 'affine Sugawara stress energy tensor' given by

$$T^s(z) = \frac{1}{2k} g_{ab} J^a(z) J^b(z) \tag{4.1.8}$$

It is easy to verify using the relations (4.1.6) that the Poisson bracket of this field with itself reads

$$\{T^s(z), T^s(w)\} = 2T^s(w)\delta'(z-w) - \partial T^s(w)\delta(z-w) \tag{4.1.9}$$

Its Fourier modes therefore satisfy a Virasoro algebra with central charge 0. In fact, in the WZW realization of the algebra (4.1.6) the affine Sugawara stress energy tensor is really the true stress energy tensor of the system.

There are however many other elements of $\mathcal{F}$ whose modes form a Virasoro algebra. Let λ^a be a set of real numbers then the fields

$$T^I(z) = T^s(z) + \lambda_a \partial J^a(z) \tag{4.1.10}$$

form Virasoro algebras with nonzero central extensions (even classically)

$$\{T^I(z), T^I(w)\} = 2T^I(w)\delta' - \partial T^I(w)\delta - k^2 \lambda_a \lambda^a \delta''' \tag{4.1.11}$$

One could think that all these Virasoro fields have the same status. This is however not the case since the fields $\{J^a(z)\}$ are only *all* primary fields w.r.t. the affine Sugawara stress energy tensor. This follows from

$$\{T^I(z), J^a(w)\} = (J^a(w) - f^{ab}_c \lambda_b J^c(w))\delta' - \partial J^a(w)\delta + k\lambda^a \delta'' \tag{4.1.12}$$

One can see from this equation that if we require all fields $J^a(z)$ to be primary then all λ^b must be zero. In that case $T^I(z)$ reduces to $T^s(z)$ and all fields $J^a(z)$ become primary fields of conformal weight 1.

Suppose we want to apply the Poisson reduction scheme that we used in the previous chapter to the case at hand. We have seen that this can lead to nonlinear algebras which, in this case, will be generated by a finite number of fields. However, we do want to make the restriction that the algebra that we obtain after reduction is a W algebra. Other reductions may be interesting from a mathematical point of view, but in this thesis we are only interested in reductions that lead to W algebras since these appear to be the

most interesting from a physical point of view. Let us therefore heuristically try to find a set of constraints that may lead to interesting reductions (of course these constraints will turn out to be just the constraints that we used in the previous chapter to construct finite W algebras). We shall then show that these constraints indeed lead to W algebras.

As we mentioned above all fields $J^a(z)$ are primary fields of conformal weight 1 w.r.t. the affine Sugawara stress energy tensor. Therefore a constraint of the form

$$J^a(z) = \text{constant} \tag{4.1.13}$$

is not in agreement with conformal invariance since constants have conformal weight 0. To put it differently, since the affine Sugawara stress energy tensor does not Poisson commute, even on the constrained manifold, with the constraint (i.e. it is not a gauge invariant quantity) it does not descend to a generator of the reduced algebra. At this moment we recall that the affine Sugawara stress energy tensor was by far not the only Virasoro generator. In fact, from equation (4.1.12) follows that, even though not all currents $J^a(z)$ are primary, most of them are and some may have conformal weight 0 w.r.t. an improved stress energy tensor. This opens up the possibility that if we impose a set of constraints like (4.1.13) that perhaps one or more improved stress energy tensors descend to become generators in the reduced algebra. of course we can turn this around, fix a certain improved stress energy tensor and then impose a set of constraints compatible to it, i.e. one constrains only currents that are primary and have conformal dimension 0. The choice we make for the improvement term will be determined by an sl_2 embedding. Given the improved stress energy tensor we shall postulate a set of constraints and then motivate these afterwards by showing that they lead to a W algebra. The role of the $sl(2)$ subalgebra will become clear later.

Let $\{t_0, t_+, t_-\}$ be an sl_2 subalgebra of g, then we can again choose a basis $\{t^{(\mu)}_{j,m}\}$ compatible with the sl_2 multiplet structure of g (see previous chapter). An arbitrary smooth map from S^1 into g can then be written as

$$J(z) = \sum_{j,m;\mu} J^{j,m}_{(\mu)}(z) t^{(\mu)}_{j,m} \tag{4.1.14}$$

Here again we take $t^{(1)}_{1,\pm 1} \equiv t_\pm$ and $t^{(1)}_{1,0} \equiv t_0$. We now take the improved affine Sugawara stress energy tensor to be

$$T^I(z) = T^s(z) + \partial J^{1,0}_{(\mu)}(z) \tag{4.1.15}$$

i.e. we let $\lambda^a t_a = t_0$. The nice thing about this improved stress energy tensor is that all currents $\{J^{j,1}_{(\mu)}\}_{j;\mu}$ are primary of conformal dimension 0. They can therefore in principle be constrained to nonzero constants without breaking conformal invariance. Not any

choice will lead to consistent reductions however (we will come to this later). A choice that leads to consistent and physically interesting reductions is

$$J^{j,1}_{(\mu)}(z) = \delta^j_1 \delta^1_\mu \tag{4.1.16}$$

i.e. one constrains the current associated to t_+ to 1 and all other currents of grade 1 are constrained to 0. The currents $\{J^{j,m}_{(\mu)}(z)\}$ for $m > 1$ are primary fields of negative conformal dimension. This we should not allow since negative conformal dimensions lead to correlations in the quantum field theory that increase with distance. Therefore these fields have to be eliminated from the theory and the only way to do this consistently with conformal invariance is by constraining them to be zero.

When the gradation of g by t_0 is half integer there will also be currents $\{J^{j,\frac{1}{2}}\}_{j;\mu}$ which are primary fields of conformal weight $\frac{1}{2}$ w.r.t. the improved stress energy tensor. In principle there is no reason to constrain these currents and in fact the constraints specified above do lead to consistent reductions. However, the algebras that one thus obtains are *not* W algebras (we shall comment further on this in a moment). We therefore also constrain these currents to zero.

Putting it all together we arrive at the following set of constraints

$$\{\phi^{j,m}_{(\mu)}(z) \equiv J^{j,m}_{(\mu)} - \delta^j_1 \delta^m_1 \delta^1_\mu\}_{\mu;\, j\in\frac{1}{2}\mathbf{N};\, m\geq 0} \tag{4.1.17}$$

Note that the constraints (3.1.28) are the direct finite dimensional analogue of the constraints (4.1.17) thus finally explaining the term 'finite W algebra'.

Now we have the following

Lemma 6 *The constraint $\phi^{j,m}_{(\mu)}(z)$ is first class if $m \geq 1$.*

The proof of this lemma in the finite case has been given in the previous chapter. In the infinite dimensional case the proof is completely analogous.

As in the finite dimensional case we can determine the gauge group generated by these first class constraints. Following exactly the same arguments we find that it is given by

$$\bar{H} \equiv Map(S^1, H) \tag{4.1.18}$$

where H is the group associated to the nilpotent Lie subalgebra

$$h = \bigoplus_{k\geq 1} g^{(-k)} \tag{4.1.19}$$

of g. Here, as before, $g^{(l)} = \mathrm{Span}(\{t^{(\mu)}_{j,l}\}_{j;\mu}$. The action of an element $g(z) \in Map(S^1, H)$ on $y(z) \in Map(S^1, g)$ is given by

$$y \mapsto gyg^{-1} + \partial g g^{-1} \tag{4.1.20}$$

Note that all these formulas are the natural affine versions of the formulas encountered in the theory of finite W algebras.

The constrained surface in $\bar{g}$ will be denoted by $\bar{g}_c$, i.e. $\bar{g}_c$ consists of elements of the form

$$y(z) = t_+ + \sum_j \sum_{m\leq 0} \sum_\mu y^{j,m}_{(\mu)}(z) t^{(\mu)}_{j,m} \tag{4.1.21}$$

One would again like to eliminate the gauge invariance by fixing it. of course there are many ways to do this. On the level of the reduced algebra a change in gauge means just an invertible change of generators. Note that a relatively simple algebra can be made to look very complicated when written in a nontrivial basis. Choosing the right gauge is therefore very important. For W algebras there is a specific type of basis which is very natural. In such a basis the algebra is generated by the Virasoro field (the stress energy tensor of the conformal field theory) and a number of W fields that are *primary* w.r.t. this stress energy tensor. As we will show there exists a natural gauge choice that automatically puts the W algebra in this preferred basis. In fact it is precisely the 'lowest weight' gauge that we used in the previous chapter. In order to see this introduce the space $\bar{g}_{fix}$ consisting of currents of the form

$$y(z) = t_+ + \sum_{j;\mu} y^{j,-j}_{(\mu)}(z) t^{(\mu)}_{j,-j} \tag{4.1.22}$$

It can be shown that

$$\bar{H} \times \bar{g}_{fix} \simeq \bar{g}_c \tag{4.1.23}$$

The proof of this statement uses precisely the same 'counting by grade' argument that was used in the finite case. From (4.1.23) follows immediately that

$$\bar{g}_c / \bar{H} \simeq \bar{g}_{fix} \tag{4.1.24}$$

Note that since the improved stress energy tensor $T^I(z)$ weakly commutes with the first class constraints (i.e. it commutes after imposing the constraints) it is a gauge invariant quantity and descends to a generator of the reduced algebra. In fact it will be the stress energy tensor of the reduced theory.

The Poisson bracket (4.1.6) induces a Dirac bracket on $\bar{g}_{fix}$. This Dirac bracket reads in the infinite dimensional case

$$\{\bar{F},\bar{G}\}^* = \overline{\{F,G\} - \int dzdw\{F,\phi^{j,m}_{(\mu)}(z)\}\Delta^{(\mu\mu')}_{j,m;j',m'}(z,w)\{\phi^{j',m'}_{(\mu')}(w),G\}} \tag{4.1.25}$$

where

$$\int dz\,\Delta^{(\mu\mu')}_{j,m;j'm'}(z,w)\Delta^{j',m';j'',m''}_{(\mu'\mu'')}(w,z') = \delta^{j''}_{j}\delta^{m''}_{m}\delta^{\mu}_{\mu''}\delta(z-z') \tag{4.1.26}$$

and

$$\Delta^{j,m;j'm'}_{(\mu\mu')}(z-w) = \{\phi^{j,m}_{(\mu)}(z),\phi^{j',m'}_{(\mu')}(w)\} \tag{4.1.27}$$

In these formulas the bar denotes restriction to $\bar{g}_{fix}$. The algebra that we are interested in is then the subalgebra $\mathcal{F}_{lw}$ of $\mathcal{F}$ generated by the fields $\{J^{j,-j}_{(\mu)}(z)\}_{j;\mu}$ and with Poisson relations

$$\{J^{j,-j}_{(\mu)}(z),J^{j',-j'}_{(\mu')}(w)\}^* \tag{4.1.28}$$

From now on we denote this algebra by $W(\bar{g};t_0)$. We now have the following important theorem

Theorem 9 *The algebra $W(\bar{g};t_0)$ is a W algebra.*

Proof: we will show that the fields $J^{j,-j}_{(\mu)}(z)$ are all primary fields w.r.t. the stress energy tensor $T^I(z)$. Consider the generator Q^I_ϵ of conformal transformations defined by

$$Q^I = \int dz\,\epsilon(z)T^I(z) \tag{4.1.29}$$

The action of this element on an arbitrary current $J^a(z)$ can easily be calculated. Denoting $J(z) = J^a(z)t_a$ we find

$$\delta_\epsilon J(z) \equiv \{Q^I_\epsilon, J(z)\} = (\epsilon J)' - k\epsilon'' t_0 - \epsilon'[t_0,J] \tag{4.1.30}$$

Now, note that $\bar{g}_c$ has been so constructed that it is stable under the transformations generated by Q^I_ϵ. This follows immediately from the fact that Q^I_ϵ commutes weakly with the constraints. A current of the form (4.1.21) will therefore be form invariant. All this does not mean however that the submanifold $\bar{g}_{fix}$ of $\bar{g}_c$ will be stable. In fact let $J(z)$ be a current of the form (4.1.22) then from (4.1.30) one finds that

$$\delta_\epsilon J = \sum_{j;\mu}\left(\epsilon\,\partial J^{j,-j}_{(\mu)} + (j+1)\epsilon' J^{j,-j}_{(\mu)}\right)t^{(\mu)}_{j,-j} - k\epsilon'' t_0 \tag{4.1.31}$$

Due to the last term in this equation it follows that if $J \in \bar{g}_{fix}$ then $J + \delta_\epsilon J$ is *not* an element of $\bar{g}_{fix}$. Note however that $J + \delta_\epsilon J$ *is* still an element of $\bar{g}_c$. From eq.(4.1.23) it therefore immediately follows that there exists a unique gauge transformation that brings us back onto $\bar{g}_{fix}$. The combined action of Q^I_ϵ and $\bar{H}$ therefore induces an action δ^*_ϵ on $\bar{g}_{fix}$. It is well known that this action is still generated by Q^I_ϵ but now by means of the Dirac bracket instead of the ordinary bracket. Let $h(z) = k\epsilon(z)''t_- \in \bar{h}$, where $\bar{h}$ is the Lie algebra of the gauge group $\bar{H}$, then the infinitesimal gauge transformation generated by this element is given by

$$\delta_h J = [h(z), J(z)] + kh'(z) \tag{4.1.32}$$

and it can easily be verified that

$$\begin{aligned} \delta^*_\epsilon J &\equiv \delta_\epsilon J + \delta_h J \\ &= \sum_{j;\mu} \left(\epsilon \partial J^{j,-j}_{(\mu)} + (j+1)\epsilon' J^{j,-j}_{(\mu)} - k^2 \epsilon''' \delta^j_1 \delta^\mu_1 \right) t^{(\mu)}_{j,-j} \qquad (4.1.33) \\ &\equiv \{Q^I_\epsilon, J(z)\}^* \qquad (4.1.34) \end{aligned}$$

From this equation it follows immediately that the current $J^{j,-j}_{(\mu)}(z)$ is a primary field of conformal dimension $j+1$ w.r.t. the stress energy tensor $T^I(z)$, i.e.

$$\delta^*_\epsilon J^{j,-j}_{(\mu)} = \epsilon \partial J^{j,-j}_{(\mu)} + (j+1)\epsilon' J^{j,-j}_{(\mu)} \tag{4.1.35}$$

This proves the theorem .

In the calculations leading up to equation (4.1.33) it is crucial that $[t_-, t^{(\mu)}_{j,-j}] = 0$. If we had not constrained the elements $t^{(\mu)}_{j,\frac{1}{2}}$ we could still have performed all the calculations, i.e. we could have found a compensating gauge transformation h and we would have found an equation similar to equation (4.1.33). However, this equation would not have had this specific form, i.e. the currents would not have been primary fields and therefore the algebra would not have been a W algebra.

Note that from the proof of the theorem follows that we can predict the field content of the W algebra $W(\bar{g}; t_0)$ without having to actually calculate any Poisson brackets. Namely, the number of fields that generate the W algebra is equal to the number of sl_2 multiplets $|i|$ into which g decomposes under the adjoint action of sl_2. Also from eq.(4.1.35) follows that the conformal dimensions are simply given by $\{\Delta = j+1\}$ where j are the spins of the sl_2 multiplets.

Having proved that the algebras that one obtains after reduction w.r.t. the constraints (4.1.17) are W algebras we now want to calculate the Dirac brackets between

the fields $J^{j,-j}_{(\mu)}(z)$. These are the brackets that will involve the nonlinearities. However, one look at the formula (4.1.25) for the Dirac bracket makes it clear that a direct calculation of these relations would be extremely cumbersome. Fortunately the method used in the previous chapter to summarise all Dirac brackets within one formula involving a matrix R generalizes to the infinite dimensional case. Again the proofs of the statements that we will make below are exact analogues of the proofs given in the previous chapter and will therefore be omitted.

Let $\mathcal{S}$ be the set of smooth test functions on S^1 and denote by $\bar{\mathcal{R}}$ the set of smooth maps

$$R : \mathcal{S}^{|i|} \times \bar{g}_{fix} \to \bar{g} \tag{4.1.36}$$

(where $|i|$ again denotes the number of sl_2 multiplets in the branching of the adjoint representation of g) of the form

$$R[\vec{a}; y](z) = \sum_{j;\mu} \sum_{k} (\partial^k a^{(\mu)}_j)(z) R^j_{(\mu;k)}[y](z) \tag{4.1.37}$$

where $\vec{a} \in \mathcal{S}^{|i|}$, $y \in \bar{g}_{fix}$ and $R^j_{(\mu;k)}[y] \in \bar{g}$. The crucial thing about elements of $\bar{\mathcal{R}}$ is that they are linear in the test fuctions $a_i(z)$.

From now on denote by $J_{fix}(z)$ the $\bar{g}_{fix}$ valued function on $\bar{g}$ given by

$$J_{fix}(z) = t_+ + \sum_{j;\mu} J^{j,-j}_{(\mu)}(z) t^{(\mu)}_{j,-j} \tag{4.1.38}$$

we then have the following theorem which is the direct analogue of theorems 2 and 3 of the previous chapter.

Theorem 10 *There exists a unique $R \in \bar{\mathcal{R}}$ such that*

$$t_+ + [R[\vec{a}; y], y] + k\partial R[\vec{a}; y] \quad \in \bar{g}_{fix} \tag{4.1.39}$$

and

$$(J_{fix}(z), R[\vec{a}; y](z)) = \sum_{j;\mu} a^{(\mu)}_j(z) J^{j,-j}_{(\mu)}(z) + (R[\vec{a}; y](z), t_+) \tag{4.1.40}$$

for all $\vec{a} \in \mathcal{S}^{|i|}$ and $y \in \bar{g}_{fix}$. Furthermore this R has the property that

$$\int dz\, a^{(\mu)}_j(z)\, \{J^{j,-j}_{(\mu)}(z), J^{j',-j'}_{(\mu')}(w)\}^*(y)\, t^{(\mu')}_{j',-j'} = [R[\vec{a}; y], y] + k\partial R[\vec{a}; y] \tag{4.1.41}$$

Eq.(4.1.41) makes it possible to calculate all the Dirac brackets. All one has to do is solve R from equations (4.1.39) and (4.1.40) after which one can read off the desired Poisson relations from eq.(4.1.41).

Again it is possible to derive a general formula for R [69]. It reads

$$R[\vec{a};y] = \frac{1}{1+L(\partial+ad_w)}F[\vec{a}] \tag{4.1.42}$$

where $w(z) = y(z) - t_+$ for $y \in \bar{g}_{fix}$ and

$$F[\vec{a}] = a^{(\mu)}_{j,-j}t^{j,-j}_{(\mu)} \tag{4.1.43}$$

Let Q be a smooth function on $\bar{g}_{lw} = Map(S^1, g_{lw})$ then define $\mathrm{grad}_w Q \in \bar{g}_{hw}$ by

$$\frac{d}{d\epsilon}Q(w+\epsilon w')|_{\epsilon=0} = ((\mathrm{grad}_w Q, w')) \tag{4.1.44}$$

for all $w' \in \bar{g}_{lw}$. From eq.(4.1.41) follows that the Dirac bracket between two arbitrary smooth functions F and G on $\bar{g}_{lw}$ is then given by

$$\{F,G\}^*(w) = (([\partial + w, \frac{1}{1+L(\partial+ad_w)}\mathrm{grad}_w F], \mathrm{grad}_w G)) \tag{4.1.45}$$

Note that for the trivial embedding (L=0) this formula reduces to the ordinary Kirillov bracket.

Let us now give some examples of the formalism developed above. In order to illustrate the rather formal construction developed above we consider the simplest example in some detail. Let $g = sl_2$, then the only nontrivial sl_2 embedding into g is the identity map. An arbitrary element $y \in Map(S^1, g)$ can then be written as

$$y = \begin{pmatrix} y^{1,0} & y^{1,1} \\ y^{1,-1} & -y^{1,0} \end{pmatrix} \tag{4.1.46}$$

while $\bar{g}_{fix}$ consists of currents of the form

$$y_{fix} = \begin{pmatrix} 0 & 1 \\ y^{1,-1} & 0 \end{pmatrix} \tag{4.1.47}$$

Let us now try to find the map $R \equiv R_{1,-1}t^{1,-1} + R_{1,0}t^{1,0} + R_{1,1}t^{1,1}$. From eq.(4.1.40) follows immediately that $R[a;y]_{1,-1} = a$. The matrix elements $R_{1,0}$ and $R_{1,1}$ can be derived from eq.(4.1.39). Doing this one easily finds that

$$\begin{aligned} R[a;y]_{1,-1} &= a \\ R[a;y]_{1,0} &= -k\partial a \\ R[a;y]_{1,1} &= ay^{1,-1} + \frac{1}{2}k^2\partial^2 a \end{aligned}$$

From eq.(4.1.41) now follows that

$$\int dz\, a(z)\{J^{1,-1}(z), J^{1,-1}(w)\}^*(y)\, t_{1,-1} = 2k\partial(ay^{1,-1}) - ka\partial y^{1,-1} + \frac{1}{2}k^3\partial^3 a \quad (4.1.48)$$

which means that if we define $T(z) = -\frac{1}{k}J^{1,-1}(z)$ then

$$\{T(z), T(w)\}^* = 2T(w)\delta' - \partial T(w)\delta - k\delta''' \quad (4.1.49)$$

which is the Virasoro algebra. We conclude that the Virasoro algebra, which is the simplest W algebra, is a reduction of the affine sl_2 algebra. This now explains our choice of constraints. Since an affine sl_2 algebra leads to a Virasoro algebra and since any W algebra *must* contain the Virasoro algebra it seems natural to reduce an affine Lie algebra such that some affine sl_2 subalgebra reduces to the required Virasoro algebra. This is precisely what we did.

It was shown in [7] that the reduction of the affine sl_N algebra w.r.t. the principal sl_2 embedding leads to W_N. Let us therefore illustrate our formalism by considering a case in which the sl_2 embedding is nonprincipal. As in the previous chapter we take for this the nonprincipal sl_2 embedding into $g = sl_3$. An arbitrary element $y \in Map(S^1, g)$ can then be written as (in the basis compatible with the sl_2 embedding)

$$y = \begin{pmatrix} \frac{1}{2}y^{1,0} + y^{0,0} & y_{(1)}^{\frac{1}{2},\frac{1}{2}} & y^{1,1} \\ y_{(2)}^{\frac{1}{2},-\frac{1}{2}} & -2y^{0,0} & y_{(2)}^{\frac{1}{2},\frac{1}{2}} \\ y^{1,-1} & y_{(1)}^{\frac{1}{2},-\frac{1}{2}} & y^{0,0} - \frac{1}{2}y^{1,0} \end{pmatrix} \quad (4.1.50)$$

while $\bar{g}_{fix}$ consists of currents of the form

$$y_{fix} = \begin{pmatrix} y^{0,0} & 0 & 1 \\ y_{(2)}^{\frac{1}{2},-\frac{1}{2}} & -2y^{0,0} & 0 \\ y^{1,-1} & y_{(1)}^{\frac{1}{2},-\frac{1}{2}} & y^{0,0} \end{pmatrix} \quad (4.1.51)$$

Again it is easy to calculate R using equations (4.1.39) and (4.1.40). Applying equation (4.1.41) oncemore we find that in the Dirac brackets between the generators

$$\begin{aligned} T(z) &= -\frac{4}{3}(J^{1,-1} + 3(J^{0,0})^2) \\ G^+(z) &= J^{\frac{1}{2},-\frac{1}{2}}_{(1)} \\ G^-(z) &= \frac{4}{3}J^{\frac{1}{2},-\frac{1}{2}}_{(2)} \\ H(z) &= 4J^{0,0} \end{aligned}$$

read

$$\begin{aligned} \{T(z),T(w)\}^* &= 2T(w)\delta' - \partial T(w)\delta - \frac{1}{2}\delta''' \\ \{T(z),G^\pm(w)\}^* &= \frac{3}{2}G^\pm(w)\delta' - \partial G^\pm(w)\delta \\ \{T(z),H(w)\}^* &= H(w)\delta' - \partial H(w)\delta \\ \{H(z),G^\pm(w)\}^* &= \pm 2G^\pm\delta \\ \{H(z),H(w)\}^* &= \delta' \\ \{G^+(z),G^-(w)\}^* &= (H^2(w) + T(w) + \partial H(w))\delta - 2H(z)\delta' + \delta''' \end{aligned}$$

This algebra is known in the literature as $W_3^{(2)}$. The quadratic version of sl_2 studied in the previous chapter is the finite W algebra underlying this algebra.

Using the formalism developed above it is easy to construct many other examples.

4.1.2 An algebraic description of W algebras

It is also possible to give an elegant algebraic description of infinite W algebras. For this remember that we can replace the mixed set of first and second class constraints (3.1.28) by an equivalent (in the sense that they lead to the same W algebra) set of constraints that is entirely first class. The point was roughly to split into half the space $g^{(\frac{1}{2})}$, associated to the second class constraints, and to add one half of it to $g^{(0)}$ and the other half to $g^{(1)}$. All this was accomplished by grading the algebra by the element δ of the Cartan subalgebra instead of t_0. Consider again the triangular decomposition

$$g = g_- \oplus g_0 \oplus g_+ \tag{4.1.52}$$

of the algebra g into a negative, zero and positive grade piece w.r.t. the operator ad_δ. As before let $\{t_a\} \equiv \{t^{(\mu)}_{j,m}\}$ be a basis of g and let $\{t_\alpha\}$ and $\{t_{\bar{\alpha}}\}$ be the subbases of g_+

and $g_0 \oplus g_-$ respectively. Remember that the alternative set of constraints is then given by

$$\phi^\alpha(z) \equiv J^\alpha(z) - \chi(J^\alpha(z)) \tag{4.1.53}$$

where $\chi(J^\alpha(z)) = 1$ if $t_\alpha = t_+$ and zero otherwise. Let $\mathcal{C}$ be the algebra of fields generated by the constraints $\{\phi^\alpha(z)\}$, i.e. as $\mathcal{C}$ contains all derivatives products and their linear combinations of the constraints. Denote by I the subalgebra

$$I \equiv \mathcal{F}.\mathcal{C} \tag{4.1.54}$$

of $\mathcal{F}$. Obviously I is an ideal in $\mathcal{F}$ and one can easily verify that it also is a Poisson subalgebra w.r.t. the Kirillov Poisson structure. Consider the quotient algebra

$$\mathcal{F}/I \equiv \mathcal{F}_{red} \tag{4.1.55}$$

It is an easy exercise to show that $\mathcal{F}_{red}$ is isomorphic to the field algebra generated by $\{J^{\bar\alpha}(z)\}$. Now, define $\mathcal{F}_{red}^{inv}$ by

$$\mathcal{F}_{red}^{inv} = \{[A(z)] \in \mathcal{F}_{red} \mid [\{\phi^\alpha, A(w)\}] = 0, \text{ for all } \alpha\} \tag{4.1.56}$$

then the Kirillov bracket on $\mathcal{F}$ naturally defines a Kirillov bracket on $\mathcal{F}_{red}^{inv}$ as follows

$$\{[A(z)], [B(w)]\}^* = [\{A(z), B(w)\}] \tag{4.1.57}$$

By construction this algebra is isomorphic to the W algebra $W(\bar g; t_0)$.

Let us now explicitly construct $\mathcal{F}_{red}^{inv}$. Its elements are fields $A(z)$ of $\mathcal{F}_{red}$ such that

$$[\{\phi^\alpha(z), A(w)\}] == 0 \tag{4.1.58}$$

As in the finite case we can construct a set of generators of $\mathcal{F}_{red}^{inv}$ as follows: Let

$$J_c(z) = t_+ + \sum_{\bar\alpha} J^{\bar\alpha}(z) t_{\bar\alpha} \tag{4.1.59}$$

then there exists a unique element $a : \bar g \to \bar{\hat H}$, where $\hat H$ is the Lie group associated to g_+, such that

$$a(z) J_c(z) a(z)^{-1} + \partial a(z) a(z)^{-1} = t_+ + \sum_{t_{\bar\alpha} \in g_{lw}} W^{\bar\alpha}(z) t_{\bar\alpha} \tag{4.1.60}$$

The fields $W^{\bar{\alpha}}(z)$ are differential polynomials in the fields $J^{\bar{\alpha}}(z)$ of the form

$$W^{\bar{\alpha}}(z) = \sum_{l=0}^{p} W_l^{\bar{\alpha}}(z) \tag{4.1.61}$$

where the degree of $W^{\bar{\alpha}}(z)$ is l, $W_p^{\bar{\alpha}}(z) = J^{\bar{\alpha}}(z)$ and $p = \deg(J^{\bar{\alpha}}(z))$. Obviously these fields are gauge invariant. The Poisson relations in $W(\bar{g}; t_0)$ are now given by

$$\{W^{\bar{\alpha}}(z), W^{\bar{\beta}}(w)\}^* = \{W^{\bar{\alpha}}(z), W^{\bar{\beta}}(w)\} \tag{4.1.62}$$

Let us give two examples of the construction described above. Obviously we have

$$J_c(z) = \begin{pmatrix} J^{1,0}(z) & 1 \\ J^{1,-1}(z) & -J^{1,0}(z) \end{pmatrix} \tag{4.1.63}$$

From eq.(4.1.60) we then find that

$$W(z) = J^{1,-1}(z) + (J^{1,0})^2(z) + \partial J^{1,0}(z) \tag{4.1.64}$$

Using eq.(4.1.62) one can now easily calculate the relation of this field with itself and one finds that it is a Virasoro algebra as expected.

It is now easy to also do the $W_3^{(2)}$ case. One finds

$$\begin{aligned} W^{0,0}(z) &= J^{0,0}(z) \\ W_{(2)}^{\frac{1}{2},-\frac{1}{2}}(z) &= J_{(2)}^{\frac{1}{2},-\frac{1}{2}} \\ W_{(1)}^{\frac{1}{2},-\frac{1}{2}}(z) &= J_{(1)}^{\frac{1}{2},-\frac{1}{2}}(z) + \frac{1}{2} J^{1,0} J_{(1)}^{\frac{1}{2},\frac{1}{2}} - 3 J_{(1)}^{\frac{1}{2},\frac{1}{2}} J^{0,0} + \partial J_{(1)}^{\frac{1}{2},\frac{1}{2}} \\ W^{1,-1}(z) &= J^{1,-1} + \frac{1}{4}(J^{1,0})^2 + J_{(1)}^{\frac{1}{2},\frac{1}{2}} J_{(2)}^{\frac{1}{2},-\frac{1}{2}} + \frac{1}{2}\partial J^{1,0} \end{aligned} \tag{4.1.65}$$

Introducing the generators

$$\begin{aligned} T(z) &= \frac{4}{3}(W^{1,-1} + 3W^{0,0}W^{0,0}) \\ G^+(z) &= W_{(1)}^{\frac{1}{2},-\frac{1}{2}} \\ G^-(z) &= \frac{4}{3} W_{(2)}^{\frac{1}{2},-\frac{1}{2}} \\ H(z) &= 4W^{0,0} \end{aligned}$$

we find, using formula (4.1.62) that these form the algebra $W_3^{(2)}$.

As in the finite case the W algebra $W(\bar{g};t_0)$ has in general a Kirillov subalgebra. Here, in the infinite dimensional case, this subalgebra is isomorphic to the Kirillov algebra associated to the affine Lie algebra over the centralizer of the sl_2 embedding. In short we have

Theorem 11 *The W algebra $W(g;t_0)$ has a Poisson subalgebra which is isomorphic to $K(\bar{C})$ where $\bar{C}$ is the affinization of the centralizer C of the sl_2 subalgebra $\{t_0,t_\pm\}$ of g.*

Proof: This follows immediately from the fact that

$$W^{0,0}_{(\mu)}(z) = J^{0,0}_{(\mu)} \tag{4.1.66}$$

for all μ .

We now describe a generalization of the Miura transformation. For this note that the Poisson bracket of a grade l_1 field and a grade l_2 field is a field of grade l_1+l_2. Therefore the Poisson brackets between the fields $W_0^{\bar{\alpha}}(z)$ and $W_0^{\bar{\beta}}(z)$ must be of precisely the same form as the Poisson brackets between the fields $W^{\bar{\alpha}}(z)$ and $W^{\bar{\beta}}(z)$. This means that the map

$$W^{\bar{\alpha}}(z) \mapsto W_0^{\bar{\alpha}}(z) \tag{4.1.67}$$

is a Poisson algebra homomorphism from the algebra generated by the fields $\{W^{\bar{\alpha}}(z)\}$ to the algebra generated by the fields $\{W_0^{\bar{\alpha}}(z)\}$. Since these fields are themselves elements of the algebra generated by the fields $\{J^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}}\in g_0}$ we find that the map (4.1.67) is an embedding of the algebra $W(\bar{g};t_0)$ into the Kirillov algebra associated to $\bar{g}_0$. This is the Miura map. We postpone the discussion of some examples of this construction until we have quantized the W algebras $W(\bar{g};t_0)$.

In this section we have generalized the classical Drinfeld-Sokolov reduction scheme. What we have obtained is a large class of W algebras. It is a notable fact that the development of the theory of infinite W algebras was completely analogous to the development of the theory of finite W algebras. This is in fact very much the same situation as with the theories of finite Lie algebras and loop algebras. The way we have presented the material was to emphasize the similarity between the finite and the infinite theory.

In the next section we are going to quantize the DS reduction scheme for arbitrary sl_2 embeddings. Apart from some formal changes related to the fact that we will be dealing with algebras of field operators the quantization will again follow closely the lines which were presented in the previous chapter.

4.2 Quantum W algebras

Having developed the theory of generalized DS reductions in the previous section we now turn to their quantization. The first attempt to quantize the standard DS reduction leading to W_N was made in [19]. First they represented the classical W_N algebras in terms of $N-1$ free bosonic fields and then quantized them by quantizing the free fields and normal ordening the expressions of the W generators. In general this procedure does not lead to a quantum algebra that closes, i.e. one will not be able to represent the commutator of two W generators in terms of W generators. It is easy to see this even if one has only a finite number of degrees of freedom (like with finite W algebras). However, the procedure *does* work, more or less by accident, in the case of sl_N (but not for other algebras) and leads to exactly the quantum relations originally given by Zamolodchikov.

A proper way to quantize DS reductions is by the BRST procedure. This was poineered in [6, 20] where the BRST cohomology for W_N was calculated by a spectral sequence argument. Unfortunately the arguments used in [20] are very difficult to generalize to DS reductions associated to arbitrary sl_2 embeddings. Also it makes use of a free field realization of the original affine Lie algebra which means that in the end one obtains the W algebra in its free field form. If the W algebra is representable in terms of affine currents, as we will show to be the case, then this will be obscured by the too explicit free field form of the W generators.

Another disadvantage of the methods developed in [6, 20] is that the W algebra is in the end described as the commutant of certain screening operators. This is very inconvenient since calculating such a commutant is in general very complicated.

In this section we shall quantize the generalized DS reductions by a method that resembles the methods of [20] but which works for arbitrary embeddings and which also lead to a completely algorithmic procedure for constructing the W algebra generators and their relations. This will also lead to a quantization of the generalized Miura transformation.

4.2.1 Quantization of generalized DS reductions

Let $\{t_a\}$ be a basis of the Lie algebra $g \equiv sl_n$. The affine Lie algebra $\bar{g}$ associated to g is the span of $\{J^a_n\}$ and the central element K. The commutation relations are given by

$$\begin{aligned} [J^a_n, J^b_m] &= f^{ab}_c J^c_{n+m} + n g^{ab} K \delta_{n+m,0} \\ [K, J^a_n] &= 0 \end{aligned} \tag{4.2.68}$$

where g^{ab} is the inverse of $g_{ab} = \mathrm{Tr}(t_a t_b)$ and $[t_a, t_b] = f^c_{ab} t_c$. As usual we use g_{ab} to raise and lower indices. Let $\mathcal{U}_k\bar{g}$ ($k \in \mathbb{C}$) be the universal enveloping algebra of g quotiented

by the ideal generated by $K - k$. The algebra $\mathcal{U}_k\bar{g}$ is the natural quantization of the Kirillov Poisson algebra associated to $\bar{g}$.

Let there again be given an sl_2 subalgebra $\{t_0, t_\pm\}$ of g and consider the grading defined by δ

$$g = \bigoplus_{k\in\mathbf{Z}} g_k \tag{4.2.69}$$

where $g_k = \{x \in g \mid [\delta, x] = kx\}$. Remember that the basis elements $\{t_a\}$ of g are chosen such that they all are basis vectors of some sl_2 multiplet. Since the sl_2 subalgebra $\{t_0, t_\pm\}$ is a triplet under its own adjoint action there must be some α_+ such that $t_+ = A\, t_{\alpha_+}$. Define the character χ of g_+ (where $\bar{g}_+$ is the affinization of g_+) by putting $\chi(J_n^\alpha) = A\delta^{\alpha,\alpha_+}\delta_{n+1,0}$ The constraints are then $J_n^\alpha = \chi(J_n^\alpha)$. As we have seen these constraints are first class which means that we can use the BRST formalism. Thereto introduce the fermionic ghost variables c_α^n, b_n^α with ghost numbers 1 and -1 respectively and relations

$$c_\alpha^n b_m^\beta + b_m^\beta c_\alpha^n = \delta_{\alpha\beta}\delta_{n+m,0} \tag{4.2.70}$$

The algebra generated by these ghost variables is the Clifford algebra $Cl(g_+ \oplus g_+^*)$. As usual one then considers the algebra

$$\Omega_k = \mathcal{U}_k g \otimes Cl(g_+ \oplus g_+^*) \tag{4.2.71}$$

For calculational purposes it is convenient (as is standard practice in conformal field theory) to introduce the following 'basic fields'

$$\begin{aligned} J^a(z) &= \sum_n J_n^a\, z^{-n-1} \\ c_\alpha(z) &= \sum_n c_\alpha^n\, z^{-n} \\ b^\alpha(z) &= \sum_n b_n^\alpha\, z^{-n-1} \end{aligned} \tag{4.2.72}$$

It is well known that the commutation relations in Ω_k can then be expressed in terms of the operator product expansions (OPE)

$$J^a(z)J^b(w) = \frac{kg^{ab}}{(z-w)^2} + \frac{f_c^{ab}}{z-w}J^c(w) + \ldots \tag{4.2.73}$$

$$c_\alpha(z)b^\beta(w) = \frac{\delta_\alpha^\beta}{z-w} \tag{4.2.74}$$

Now inductively define the *algebra of fields* $F(\Omega_k)$ as follows:

1. $J^a(z), c_\alpha(z), b^\alpha(z)$ are elements of $F(\Omega_k)$ with 'conformal dimensions' $\Delta = 1, 0, 1$ respectively.

2. if $A(z), B(z) \in F(\Omega_k)$ then $\alpha A(z) + \beta B(z) \in F(\Omega_k)$

3. if $A(z)$ is an element of $F(\Omega_k)$ of conformal dimension Δ then $\frac{dA}{dz}(z)$ is also an element of $F(\Omega_k)$ and has conformal dimension $\Delta + 1$

4. if $A(z), B(z)$ are elements of $F(\Omega_k)$ of conformal dimensions Δ_A and Δ_B respectively then the normal ordened product $(AB)(z) \equiv A_-(z)B(z) \pm B(z)A_+(z)$ (where one has the minus sign if A and B are fermionic) is also an element of $F(\Omega_k)$ and has conformal dimension $\Delta_A + \Delta_B$. Here $A_-(z) = \sum_{n \leq -\Delta_A} A_n \, z^{-n-\Delta_A}$ and $A_+(z) = A(z) - A_-(z)$.

We say that $F(\Omega_k)$ is 'generated' by the basic fields. Note that $F(\Omega_k) \subset \mathcal{U}_k g[[z, z^{-1}]]$. The algebra $F(\Omega_k)$ is graded by ghost number, i.e. $J^a(z)$, $c_\alpha(z)$ and $b^\alpha(z)$ have degrees 0,1 and -1 respectively and we have the decomposition

$$F(\Omega_k) = \bigoplus_n F(\Omega_k)^{(n)} \tag{4.2.75}$$

The algebra of fields $F(\Omega_k)$ is not simply the set of 'words' in the fields that can be made using the rules given above, there are also relations. If we denote the operator product expansion of A and B by $A(z)B(w) \sim \sum_r \frac{\{AB\}_r}{(z-w)^r}$, then the relations valid in $F(\Omega_k)$ are [89]

$$\begin{aligned} (AB)(z) - (BA)(z) \equiv [A, B](z) &= \sum_{r>0} (-1)^{r+1} \frac{\partial^r}{r!} \{AB\}_r \\ (A(BC))(z) - (B(AC))(z) &= ([A, B]C)(z) \\ \partial(AB)(z) &= (\partial AB)(z) + (A\partial B)(z). \end{aligned} \tag{4.2.76}$$

The BRST operator is then [86] $d(.) = [Q, .]$ where $Q = \oint \frac{dz}{2\pi i} Q(z)$ and

$$Q(z) = (J^\alpha(z) - \chi(J^\alpha(z)))c_\alpha(z) - \frac{1}{2} f_\gamma^{\alpha\beta} (b^\gamma (c_\alpha c_\beta))(z) \tag{4.2.77}$$

As one immediately sees from eq. (4.2.77) we have

$$d(F(\Omega_k)^{(l)}) \subset F(\Omega_k)^{(l+1)}) \tag{4.2.78}$$

which means that d is of degree 1. Furthermore, by explicit calculation one finds that $d^2 = 0$ and the space $F(\Omega_k)$ is turned into a complex

$$\ldots \to F(\Omega_k)^{(n-1)} \to F(\Omega_k)^{(n)} \to F(\Omega_k)^{(n+1)} \to \ldots \tag{4.2.79}$$

We want to calculate the zeroth cohomology of this complex. This calculation is completely analogous to the one we did for finite W algebras [98] and it will therefore not be necessary to go into it in detail. Again one splits the BRST operator into two anticommuting pieces d_0 and d_1 (3.2.143) and turns the complex (4.2.79) into a double complex by the double grading (3.2.144). The action of d_0 and d_1 on the generators $\hat{J}^a(z) = J^a(z) + f^{a\beta}_{\gamma}(b^\gamma c_\beta)(z)$, $c_\alpha(z)$ and $b^\alpha(z)$ of $F(\Omega_k)$ is the same as in the finite case (3.2.147) except for

$$D_1(\hat{J}^a(z)) = f^{\alpha a}_{\bar{\beta}}\hat{J}^{\bar{\beta}}(z)c_\alpha(z) + (kg^{a\alpha} - f^{\alpha e}_{\beta}f^{\beta a}_{e})\partial c_\alpha(z) \tag{4.2.80}$$

Obviously the last two terms, which involve derivatives, were not present earlier. These terms do not spoil however the fact that the subalgebra $F^\alpha(\Omega_k)$ of $F(\Omega_k)$ generated by $J^\alpha(z)$ and $b^\alpha(z)$ is a subcomplex and that the cohomology of this complex is equal to $\mathbb{C}$ as in the finite case. From here on the calculation is completely analogous, i.e. again one shows that $H^*(F(\Omega_k); d) \simeq H^*(F_{red}(\Omega_k); d)$ where $F_{red}(\Omega_k)$ is the subcomplex of $F(\Omega_k)$ generated by $\hat{J}^{\bar{\alpha}}(z)$ and $c_\alpha(z)$, and then one calculates $H^*(F_{red}(\Omega_k); d)$ by a spectral sequence argument. It will be no surprise that the final answer reads

$$H^n(F(\Omega_k); D) \simeq F_{lw}(\Omega_k)\delta_{n,0} \tag{4.2.81}$$

where $F_{lw}(\Omega_k)$ is the subspace of $F(\Omega_k)$ generated by the fields $\{J^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}} \in g_{lw}}$. Equation (4.2.81) should be understood to hold purely on the level of vectorspaces. In order to explicitly construct the generators of cohomology we again need to apply the tic-tac-toe construction. It is however possible to immediately write down one generator, namely the one that will correspond with the stress energy tensor of the reduced theory. For this consider the field

$$T = T_I + T_{gh} \tag{4.2.82}$$

where T_I is the improved affine Sugawara stress energy tensor

$$T_I = \frac{1}{2(k+h)} g_{ab}(J^a J^b) - s_a \partial J^a, \tag{4.2.83}$$

s_a are determined by $t_0 = s^a t_a$ and T_{gh} is the 'ghost stress energy tensor'

$$T_{gh} = (\delta_\alpha - 1)b^\alpha \partial c_\alpha + \delta_\alpha \partial b^\alpha \tag{4.2.84}$$

The coefficients δ_α are determined by $[t_0, t_\alpha] = \delta_\alpha t_\alpha$ while h is the dual Coxeter number of the Lie algebra g. Obviously $T \in F(\Omega_k)$ and the operator product expansion of T

with itself reveals that it satisfies a Virasoro algebra with central charge

$$c(k;t_0) = \dim(g^{(0)}) - \frac{1}{2}\dim(g^{(\frac{1}{2})}) - 12\mathrm{Tr}\left(\frac{\rho}{\sqrt{k+h}} - t_0\sqrt{k+h}\right)^2 \tag{4.2.85}$$

where ρ is the half sum of the positive roots. (Remember that the grade n subspace of g w.r.t t_0 is denoted by $g^{(n)}$ while the grade n subspace w.r.t. δ is denoted by g_n.) The point is that one can show by explicit calculation that $T(z)$ is a nontrivially BRST closed field, i.e.

$$d(T(z)) = 0 \tag{4.2.86}$$

and that it therefore is a generator of cohomology. Since the product of two cohomology classes is simply the cohomology class of the product of two representatives we find that in cohomology the field T also generates a Virasoro algebra with central charge (4.2.85).

To obtain the other generators of $H^*(F(\Omega_k);d)$ one again applies the tic-tac-toe construction in the same way as we did in the finite case. For this consider a generator $\hat{J}^{\bar{\alpha}}(z)$ of degree $(p,-p)$ of the field algebra $F_{lw}(\Omega_k)$ (i.e. $t_{\bar{\alpha}} \in g_{lw}$) then the generator of cohomology associated to this element is given by

$$W^{\bar{\alpha}}(z) = \sum_{l=0}^{p}(-1)^l W_l^{\bar{\alpha}}(z) \tag{4.2.87}$$

where $W_0^{\bar{\alpha}}(z) \equiv J^{\bar{\alpha}}(z)$ and $W_l^{\bar{\alpha}}(z)$ can be determined inductively by

$$d_1(W_l^{\bar{\alpha}}(z)) = d_0(W_{l+1}^{\bar{\alpha}}(z)) \tag{4.2.88}$$

It is easy to check, using the fact that $d_0(J^{\bar{\alpha}}(z)) = 0$ for $t_{\bar{\alpha}} \in g_{lw}$ that indeed $d(W^{\bar{\alpha}}(z)) = 0$. These then are the generators of the W algebra. The relations in the W algebra can be found by calculating the OPE's between the fields $\{W^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}}\in g_{lw}}$ using the OPEs in $F(\Omega_k)$. In principle this algebra closes only modulo d-exact terms. But since we computed the d cohomology on a reduced comlex generated by $\hat{J}^{\bar{\alpha}}(z)$ and $c_\alpha(z)$, and this reduced complex is zero at negative ghost number, there simply aren't any D exact terms at ghost number zero. Thus the algebra generated by $\{W^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}}\in g_{lw}}$ closes in itself.

As was shown in the previous chapter for finite W algebras, the operator product algebra generated by the fields $W^{\bar{\alpha}}(z)$ is isomorphic to the operator product algebra generated by their (bi)grade (0,0) components $W_p^{\bar{\alpha}}(z)$ (the proof in the infinite dimensional case is completely analogous and will not be repeated here). The fields $W_p^{\bar{\alpha}}(z)$ are of course elements of the field algebra generated by the currents $\{\hat{J}^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}}\in g_0}$. The

relations (i.e. the OPEs) satisfied by these currents are almost identical to the relations satisfied by the unhatted currents

$$\hat{J}^{\bar{\alpha}}(z)\hat{J}^{\bar{\beta}}(w) = \frac{kg^{\bar{\alpha}\bar{\beta}} + k^{\bar{\alpha}\bar{\beta}}}{(z-w)^2} + \frac{f_{\bar{\gamma}}^{\bar{\alpha}\bar{\beta}}\hat{J}^{\bar{\gamma}}(w)}{z-w} + \dots \tag{4.2.89}$$

where $k^{\bar{\alpha}\bar{\beta}} = f_{\gamma}^{\bar{\alpha}\lambda} f_{\lambda}^{\bar{\beta}\gamma}$. Now, it is easy to see that g_0 is just a direct sum of sl_{p_j} and $u(1)$ algebras, i.e. forgetting for a moment about the $u(1)$ algebras one can write

$$g_0 \sim \bigoplus_j sl_{p_j} \tag{4.2.90}$$

Within the sl_{p_j} component of g_0 we have the identity

$$k^{\bar{\alpha}\bar{\beta}} = g^{\bar{\alpha}\bar{\beta}}(h - h_j) \tag{4.2.91}$$

where h is the dual coxeter number of g and h_j is the dual coxeter number of sl_{p_j}. We therefore find that the field algebra generated by the currents $\{\hat{J}^{\bar{\alpha}}(z)\}_{t_{\bar{\alpha}} \in g_0}$, denoted from now on by $\hat{F}_0$, is nothing but the field algebra associated to a semisimple affine Lie algebra the components of which are affine sl_{p_j} and $u(1)$ Lie algebras. This semisimple affine Lie algebra is not simply $\bar{g}_0$ (whose field algebra is generated by the unhatted currents) however because in $\bar{g}_0$ all components have the same level while in $\hat{F}_0$ the level varies from component to component as follows from equation (4.2.91). This is just a result of the ghost contributions $k^{\bar{\alpha}\bar{\beta}}$ in the OPEs of the hatted currents.

From the above we find that the map

$$W^{\bar{\alpha}}(z) \mapsto (-1)^p W_p^{\bar{\alpha}}(z) \tag{4.2.92}$$

is an embedding of the W algebra into $\hat{F}_0$. This provides a quantization and generalization to arbitrary sl_2 embeddings of the well known Miura map. In [19] the standard Miura map for W_N algebras was naively quantised by simply normal ordening the classical expressions. This is known to work only for certain algebras [63]. Our construction gives a rigorous derivation of the quantum Miura transformations for arbitrary affine Lie algebras and sl_2 embeddings (the generalized Miura transformations for a certain special class of sl_2 embeddings were also recently given in [15]).

As a result of the generalized quantum Miura transformation *any* representation or realization of $\hat{F}_0$ gives rise to a representation or realization of the W algebra. In particular one obtains a free field realization of the W algebra by choosing the Wakimoto free field realization of $\hat{F}_0$. Given our formalism it is therefore straightforward to construct free field realizations for any W algebra that can be obtained by Drinfeld-Sokolov reduction.

4.2.2 Examples

In this section we consider some representative examples of the construction described above. For notational convenience we shall omit the explicit z dependence of the fields where possible.

The Virasoro algebra

The Virasoro algebra is the simplest W algebra and it is well known to arise from the affine sl_2 algebra by quantum Drinfeld-Sokolov reduction [6]. It is the W algebra associated to the only nontrivial embedding of sl_2 into itself, namely the identity map. We consider this example here to contrast our methods to the ones used by Bershadsky and Ooguri.

Choose the following basis of sl_2

$$J^a t_a = \begin{pmatrix} J^2/2 & J^3 \\ J^1 & -J^2/2 \end{pmatrix}. \tag{4.2.93}$$

where $t_0 = -t_2,\ t_+ = t_1$ and $t_- = t_3$. The positive grade piece of the Lie algebra $g = sl_2$ is generated by t_1, and the constraint to be imposed is $J^1 = 1$. The BRST current $d(z)$ is given simply by

$$d(z) = (J^1 c_1) - c_1. \tag{4.2.94}$$

The 'hatted' currents are $\hat{J}^1 = J^1, \hat{J}^2 = J^2 + 2(b^1 c_1)$ and $\hat{J}^3 = J^3$. The actions of d_0 and d_1 are given by

$$\begin{array}{lcllcl} d_0(\hat{J}^2) & = & -2c_1 & d_1(\hat{J}^3) & = & (\hat{J}^2 c_2) + (k+2)\partial c_1 \\ d_0(b^1) & = & -1 & d_1(b^1) & = & \hat{J}^1. \end{array} \tag{4.2.95}$$

On the other fields d_0 and d_1 vanish. From (4.2.95) it is immediately clear that $H(F_{red}(\Omega_k); d_0)$ is generated by $W_0^3 \equiv \hat{J}^3$, in accordance with the general arguments in section 2. To find the generator of the d-cohomology, we apply the tic-tac-toe construction. We are looking for an element $W_1^3(z) \in F_{red}(\Omega_k)$ such that $d_0(W_1^3(z)) = d_1(W_0^3)$. The strategy is to write down the most general form of $W_1^3(z)$, and then to fix the coefficients. In general, $W_l^{\bar{\alpha}}$ must satisfy the following two requirements

1. if $W_l^{\bar{\alpha}}$ has bidegree $(-k, k)$, then $W_{l+1}^{\bar{\alpha}}$ must have bidegree $(-k-1, k+1)$
2. if we define inductively the *weight* h of a monomial in the $J^{\bar{\alpha}}$ by $h(J^{\bar{\alpha}}) = 1 - k$ if $t_{\bar{\alpha}} \in g_k$, $h((AB)) = h(A) + h(B)$ and $h(\partial A) = h(A) + 1$*, then $h(W_l^{\bar{\alpha}}) = h(W_{l+1}^{\bar{\alpha}})$

* The weight h is similar to the conformal weight, but not always equal to it. It is independent of the way in which the $\hat{J}$ are ordered.

These two conditions guarantee that the most general form of $W_l^{\bar{\alpha}}$ will contain only a finite number of parameters, so that in a sense the tic-tac-toe construction is a finite algorithm. In the case under hand, the most general form of W_1^3 is $a_1(\hat{J}^2\hat{J}^2)+a_2\partial\hat{J}^2$, and the D_0 of this equals $-4a_1(\hat{J}^2c_1)-(4a_1+2a_2)\partial c_1$. Thus, $a_1=-1/4$ and $a_2=-(k+1)/2$. Since $d_1(W_1^3)=0$, the tic-tac-toeing stops here and the generator of the D cohomology reads

$$W^3 = W_0^3 - W_1^3 = \hat{J}^3 + \frac{1}{4}(\hat{J}^2\hat{J}^2) + \frac{(k+1)}{2}\partial\hat{J}^2. \tag{4.2.96}$$

As one can easily verify, $T = W^3/(k+2)$ generates a Virasoro algebra with central charge

$$c(k) = 13 - 6(k+2) - \frac{6}{k+2} \tag{4.2.97}$$

a result first found by Bershadsky and Ooguri [6].

Let's now consider the quantum Miura transformation. In the case at hand $\bar{g}_0$ is the Cartan subalgebra spanned by t_2 and $\hat{F}_0$ is an affine $u(1)$ field algebra at level $k+2$ generated by $\hat{J}^2$. Indeed defining the field $\partial\phi \equiv \nu\hat{J}^2$ where $\nu = \sqrt{2(k+2)}$, it is easy to check that

$$\partial\phi(z)\,\partial\phi(w) = \frac{1}{(z-w)^2} + \dots \tag{4.2.98}$$

In terms of the field ϕ the (bi)grade (0,0) piece of T is given by

$$T^{(0,0)} = \frac{1}{2}(\partial\phi\partial\phi) + \alpha_0\partial^2\phi \tag{4.2.99}$$

where $\alpha_0 = \frac{1}{2}\nu - \frac{1}{\nu}$. This is the usual expression for the Virasoro algebra in terms of a free field ϕ.

The Zamolodchikov W_3 algebra

Having illustrated the construction in some detail for the Virasoro algebra, we will now briefly discuss two other examples. We start with the Zamolodchikov W_3 algebra [62]. This algebra is associated to the principal sl_2 embedding into sl_3. In terms of the following basis of sl_3

$$J^a t_a = \begin{pmatrix} \frac{J^4}{6} + \frac{J^5}{2} & \frac{1}{2}(J^6+J^7) & J^8 \\ \frac{1}{2}(J^2+J^3) & -\frac{J^4}{3} & \frac{1}{2}(J^7-J^6) \\ J^1 & \frac{1}{2}(J^2-J^3) & \frac{J^4}{6} - \frac{J^5}{2} \end{pmatrix} \tag{4.2.100}$$

The sl_2 subalgebra is given in this case by $t_+ = 4t_2$, $t_0 = -2t_5$ and $t_- = 2t_7$. The constraints are therefore $J^1 = J^3 = 0$ and $J^2 = 1$ according to the general prescription.

The BRST current associated to these (first class) constraints read

$$Q(z) = J^1 c_1 + (J^2 - 1)c_2 + J^3 c_3 + 2(b^1(c_2 c_3)). \tag{4.2.101}$$

The BRST cohomology is generated by $\hat{J}^7$ and $\hat{J}^8$ since t_7 and t_8 are the lowest weight vectors w.r.t. the sl_2 subalgebra. The tic-tac-toe construction gives as generators of $H^*(F(\Omega_k); d)$ the fields $W^7 = W_0^7 - W_1^7$ and $W^8 = W_0^8 - W_1^8 + W_2^8$ where

$$\begin{aligned}
W_0^7 &= \hat{J}^7 \\
W_1^7 &= -\frac{1}{6}(\hat{J}^4\hat{J}^4) - \frac{1}{2}(\hat{J}^4\hat{J}^5) - 2(k+2)\partial\hat{J}^5 \\
W_0^8 &= \hat{J}^8 \\
W_1^8 &= -(\hat{J}^5\hat{J}^6) + \frac{1}{3}(\hat{J}^4\hat{J}^7) - (k+2)\partial\hat{J}^6 \\
W_2^8 &= -\frac{1}{27}(\hat{J}^4\hat{J}^4\hat{J}^4) + \frac{1}{3}(\hat{J}^4(\hat{J}^5\hat{J}^5)) + \frac{(k+2)}{3}(\hat{J}^4\partial\hat{J}^4) + (k+2)(\hat{J}^5\partial\hat{J}^4) \\
&\quad + \frac{2(k+2)^2}{3}\partial^2\hat{J}^4
\end{aligned} \tag{4.2.102}$$

with some work, for instance by using the program for computing OPE's by Thielemans [57], one finds that

$$\begin{aligned}
T &= \frac{1}{2(k+3)}W^7 \\
W &= \left(\frac{3}{(5c+22)(k+3)^3}\right)^3 W^8
\end{aligned} \tag{4.2.103}$$

generate the W_3 algebra with central charge

$$c(k) = 50 - 24(k+3) - \frac{24}{k+3} \tag{4.2.104}$$

For any principal embedding the grade zero subalgebra $\bar{g}_0$ is just the Cartan subalgebra. In the case at hand $\hat{F}_0$ is therefore a direct sum of two affine $u(1)$ field algebras, both of level $k+3$, generated by $\hat{J}^4$ and $\hat{J}^5$. Defining $\partial\phi_1 \equiv \nu_1\hat{J}^4$ and $\partial\phi_2 \equiv \nu_2\hat{J}^5$, where $\nu_1 = \sqrt{6(k+3)}$ and $\nu_2 = \sqrt{2(k+3)}$ it is easy to check that

$$\partial\phi_i(z)\partial\phi_j(w) = \frac{\delta_{ij}}{(z-w)^2} + \dots \tag{4.2.105}$$

According to the general prescription the Miura transformation reads in this case $W^7 \mapsto -W_1^7$ and $W^8 \mapsto W_2^8$, and the fields

$$\begin{aligned} T^{(0,0)} &= -\nu_2^{-2} W_1^7 \\ W^{(0,0)} &= \left(\frac{3}{(5c+22)(k+3)^3}\right)^{\frac{1}{2}} W_2^8 \end{aligned} \tag{4.2.106}$$

also generate a W_3 algebra with central charge (4.2.104). Note that according to (4.2.102) $T^{(0,0)}$ and $W^{(0,0)}$ only depend on ϕ_1 and ϕ_2. This is the well known free field realization of W_3.

The two examples given above are associated to principal embeddings. The construction presented above also works for nonprincipal embeddings however as we shall illustrate in the next example.

The $W_3^{(2)}$ algebra

To describe the $W_3^{(2)}$ algebra, we pick the following basis of sl_3

$$J^a t_a = \begin{pmatrix} \frac{J^4}{6} - \frac{J^5}{2} & J^6 & J^8 \\ J^2 & -\frac{J^4}{3} & J^7 \\ J^1 & J^3 & \frac{J^4}{6} + \frac{J^5}{2} \end{pmatrix}. \tag{4.2.107}$$

The sl_2 embedding reads $t_+ = t_1$, $t_0 = t_5$ and $t_- = t_8$. The gradation of $\bar{g}$ with respect to ad_{t_0} is half-integer which means that there will be second class constraints [67] corresponding to the fields with grade -1/2. If one wants to use the BRST formalism all constraints should be first class. One way to get around this problem is to introduce auxiliary fields [93]. This is not necessary however as was shown in [80, 97] since it is always possible to replace the half integer grading and the constraints associated to it by an integer grading and a set of first class constraints that nevertheless lead to the same Drinfeld-Sokolov reduction. As mentioned earlier we have to replace the grading by t_0 by a grading w.r.t. an element δ. [†] In this specific case $\delta = \frac{1}{3}\mathrm{diag}(1, 1, -2)$. With respect to ad_δ, t_1 and t_3 have degree 1 and span g_+. The BRST current is

$$d(z) = ((J^1 - 1)c_1) + J^3 c_3. \tag{4.2.108}$$

Notice that there is no need for auxiliary fields, since the constraints $J^1 = 1$ and $J^3 = 0$ are first class. The cohomology $H(F_{red}(\Omega_k); D_0)$ is generated by $\{\hat{J}^4, \hat{J}^6, \hat{J}^7, \hat{J}^8\}$. Again

[†]Essentially what one does is split the set of second class constraints in two halves. The constraints in the first half, corresponding to positive grades w.r.t. δ, are still imposed but have now become first class. The other half can then be obtained as gauge fixing conditions of the gauge invariance generated by the first half.

using the tic-tac-toe construction one finds that $H(F(\Omega_k);D)$ is generated by $W^4 = W_0^4$; $W^6 = W_0^6$; $W^7 = W_0^7 - W_1^7$ and $W^8 = W_0^8 - W_1^8$, where

$$\begin{aligned}
W_0^4 &= \hat{J}^4, \\
W_0^6 &= \hat{J}^6, \\
W_0^7 &= \hat{J}^7, \\
W_1^7 &= \frac{1}{2}(\hat{J}^2\hat{J}^5) + \frac{1}{2}(\hat{J}^2\hat{J}^4) - (k+1)\partial\hat{J}^2, \\
W_0^8 &= \hat{J}^8, \\
W_1^8 &= -\frac{1}{4}(\hat{J}^5\hat{J}^5) - (\hat{J}^2\hat{J}^6) + \frac{(k+1)}{2}\partial\hat{J}^5.
\end{aligned} \tag{4.2.109}$$

The OPEs of the hatted currents involving shifts in level are in this case

$$\begin{aligned}
\hat{J}^4(z)\hat{J}^4(w) &\sim \frac{6(k+\frac{3}{2})}{(z-w)^2} + \cdots \\
\hat{J}^5(z)\hat{J}^5(w) &\sim \frac{2(k+\frac{5}{2})}{(z-w)^2} + \cdots \\
\hat{J}^4(z)\hat{J}^5(w) &\sim \frac{3}{(z-w)^2} + \cdots \\
\hat{J}^2(z)\hat{J}^6(w) &\sim \frac{(k+1)}{(z-w)^2} + \frac{\frac{1}{2}(\hat{J}^4 - \hat{J}^5)}{(z-w)} + \cdots.
\end{aligned} \tag{4.2.110}$$

If we now define the following generators

$$\begin{aligned}
H &= -W^4/3, \\
G^+ &= W^6, \\
G^- &= W^7, \\
T &= \frac{1}{k+3}(W^8 + \frac{1}{12}(W^4W^4)),
\end{aligned} \tag{4.2.111}$$

we find that these generate the $W_3^{(2)}$ algebra [93]

$$\begin{aligned}
T(z)T(w) &= \frac{c(k)}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w}\cdots \\
T(z)G^{\pm}(w) &= \frac{\frac{3}{2}G^{\pm}(w)}{(z-w)^2} + \frac{\partial G^{\pm}(w)}{z-w}\cdots \\
T(z)H(w) &= \frac{H(w)}{(z-w)^2} + \frac{\partial H(w)}{z-w}\cdots \\
G^+(z)G^-(w) &= \frac{(k+1)(2k+3)}{(z-w)^3} + \frac{3(k+1)H(w)}{(z-w)^2} +
\end{aligned}$$

$$+\frac{1}{z-w}(3H^2(w)-(k+3)T(w)+\frac{3(k+1)}{2}\partial H(w))+\dots$$

$$H(z)H(w) = \frac{\frac{1}{3}(2k+3)}{(z-w)^2} \tag{4.2.112}$$

with

$$c(k)=25-6(k+3)-\frac{24}{k+3} \tag{4.2.113}$$

a formula that was found in [93] by a counting argument.

In the case at hand the subalgebra g_0 is spanned by the elements $t_4,\ t_5,\ t_6$ and t_2. Obviously $g_0 \simeq sl_2 \oplus u(1)$. Therefore $\hat{F}_0$ is the direct sum of an affine sl_2 and an affine $u(1)$ field algebra, and using equation (4.2.91) the levels of these can be calculated to be $k+1$ and $k+3$ respectively. Indeed if we introduce the currents

$$\begin{aligned}
\partial\phi &= \frac{1}{4}(\hat{J}^4+3\hat{J}^5)\\
J^0 &= \frac{1}{4}(\hat{J}^5-\hat{J}^4)\\
J^- &= \frac{1}{2}\,\hat{J}^2\\
J^+ &= 2\,\hat{J}^6
\end{aligned}$$

then these satisfy the following OPEs

$$\begin{aligned}
J^0(z)J^\pm(w) &= \frac{\pm J^\pm(w)}{z-w}+\dots\\
J^+(z)J^-(w) &= \frac{(k+1)}{(z-w)^2}+\frac{2J^0(w)}{z-w}+\dots\\
J^0(z)J^0(w) &= \frac{\frac{1}{2}(k+1)}{(z-w)^2}+\dots\\
\partial\phi(z)\,\partial\phi(w) &= \frac{\frac{3}{2}(k+3)}{(z-w)^2}+\dots
\end{aligned}$$

and all other OPEs are regular. As stated before the shifts in the levels that one can see in these OPEs are a result of the ghost contributions.

The quantum Miura transformation in this case reads: $W^4 \mapsto W_0^4,\ W^6 \mapsto W_0^6,$ $W^7 \mapsto -W_1^7,\ W^8 \mapsto -W_1^8$. This means that in terms of the currents $J^\pm,\ J^0$ and ϕ the grade (0,0) components of the W generators read (let's for notational convenience denote the (0,0) components of H, G^+, G^- and T again simply by the same letters since they generate an isomorphic algebra anyway)

$$H = J^0-\frac{1}{3}\partial\phi$$

$$\begin{aligned} G^+ &= \frac{1}{2}J^+ \\ G^- &= (J^-J^0) + (J^0J^-) + 2(k+3)\partial J^- - 2\partial\phi J^- \\ T &= \frac{1}{2(k+3)}\Big(2(J^0J^0) + (J^-J^+) + (J^+J^-) + (k+3)\partial J^0 + \frac{2}{3}(\partial\phi\partial\phi) + Q_0\partial^2\phi\Big) \end{aligned}$$

where $Q_0 = -(k+1)$. We recognise in the expression for T the improved sl_2 affine Sugawara stress energy tensor and the free boson in a background charge. Note that these formulas provide an embedding of $W_3^{(2)}$ into $\hat{F}_0$. In [67] a realization of this type was called a 'hybrid field realization' since it represents the W algebra partly in terms of affine currents and partly in terms of free fields.

It is now easy to obtain a realization of $W_3^{(2)}$ completely in terms of free fields by inserting for the affine sl_2 currents $J^{\pm}$, J^0 their Wakimoto free field form

$$\begin{aligned} J^- &= \beta \\ J^+ &= -(\gamma^2\beta) - k\partial\gamma - \sqrt{2k+6}\,\gamma\, i\partial\varphi \\ J^0 &= -2(\gamma\beta) - \sqrt{2k+6}\, i\,\partial\varphi \end{aligned} \tag{4.2.114}$$

where as usual β, γ and φ are bosonic fields with the following OPEs

$$\begin{aligned} \gamma(z)\beta(w) &= \frac{1}{z-w} + \dots \\ i\partial\varphi(z)\, i\partial\varphi(w) &= \frac{1}{(z-w)^2} + \dots \end{aligned} \tag{4.2.115}$$

This example gives a nice taste of the general case. By the Miura transformation one can write down for any W algebra a hybrid field realization, i.e. a realization partly in terms of free fields and partly in affine currents. When required one can then insert for the affine currents the Wakimoto free field realization giving you a realization of the W algebra completely in terms of free fields.

Summary and conclusions

In this chapter we have shown that a large class of W algebras can be obtained by generalizing and quantizing the Drinfeld-Sokolov reduction scheme. The inequivalent reductions are related to the inequivalent embeddings of sl_2. The formalism yields completely algorithmic methods for calculating the W algebra generators and relations in both the classical and quantum case. We also generalized and quantized the Miura map. This lead to the statement that any W algebra can be embedded into a semisimple affine Lie algebra. Using this result it was possible to give a general and explicit method for constructing free field realizations for arbitrary W algebras.

The question naturally arises whether all W algebras can be described as BRST cohomologies of affine Lie algebras. The answer to this question is not known at this time but there are some indications that it is true. No matter how it turns out, the description of W algebras as reductions of Lie algebras is extremely powerful. It is in fact the only approach that, up to date, has lead to any results w.r.t. the physical models behind W algebras or their geometry.

Chapter 5

W symmetry in integrable systems

The intersection between topics related to string theory like conformal field theory, W gravity, matrix models, topological field theory etc. and the theory of integrable systems is substantial. Consider for example the facts that the second Hamiltonian stucture of the KdV hierarchy is equal to the Virasoro algebra and also that the partition function of the one matrix model is (after taking the double scaling limit) a KdV tau function. Also remember that the theory describing 2 dimensional induced gravity in the chiral gauge is nothing but Toda field theory which is a well known integrable system. It appears that string theory is connected on a fundamental level with the theory of integrable systems. Certainly one reason for this is conformal invariance which, as is well known, is infinite dimensional in 2 dimensions.

In this chapter we shall consider some applications to integrable systems of the theory developed in the previous chapters. In the first section show that finite dimensional Toda systems are reductions of systems describing free particles moving on group manifolds. Using this picture of finite Toda systems we shall also be able to generalize them by again considering nonprincipal sl_2 embeddings. This is of course a close analogue of the reduction of WZW models to Toda field theories which is also briefly reviewed in the second section. The (finite and infinite) Toda systems thus obtained have as their symmetry algebras the W algebras constructed in the previous chapters.

In the last section we construct new hierarchies of integrable evolution equations that generalize the KdV type hierarchies introduced by Gelfand and Dickii. The difference between the Gelfand-Dickii hierarchies and the hierarchies we shall be considering is that the GD hierarchies contain evolution equations for scalar fields while our hierarchies will in general be for matrix fields. Also the spatial derivatives in the equations will be replaced by covariant derivatives w.r.t. some static gauge field.

5.1 Generalized finite Toda systems

Let G be a simple Lie group of rank l and let g be its Lie algebra. Consider the following coupled set of equations of motion for the coordinates $\{q_i\}_{i=1}^{l}$

$$\frac{d^2q_i}{dt^2} + \exp(\sum_{j=1}^{l} K_{ij}q_j) = 0 \tag{5.1.1}$$

where K_{ij} is the Cartan matrix of g. Systems of this form are called finite dimensional Toda systems and are well known in the theory of integrable systems (for a review see [49]). In this section we will show, using the formalism developed in the previous chapters, that the Toda system (5.1.1) is a reduction of a system describing a free particle moving on the group G. As we will show this reduction is again associated to a principal sl_2 embedding. By also considering the other sl_2 embeddings we will also be able to generalize the class of finite dimensional Toda systems substantially.

The action of a 'free' particle moving on a Riemann manifold M with local coordinates $\{x^\mu\}$ and metric (in these coordinates) $h_{\mu\nu}(x)$ is

$$S = \frac{1}{2}\int dt\; h_{\mu\nu}(x)\frac{dx^\mu}{dt}\frac{dx^\nu}{dt} \tag{5.1.2}$$

An example of such a manifold is a simple Lie group. The Lie algebra g of a simple Lie group G is simple which means that the Cartan-Killing form on it is nondegenerate. Since the Lie algebra of G is nothing but the tangent space in the unit element we can take the Cartan-Killing form to be the 'metric in the unit element'. Because of the group structure however we can extend this metric to the whole group manifold by left translation. What we thus obtain is the unique left and right invariant metric on G. This allows us to write down the following coordinate free formula for the action of a free particle moving on the group G:

$$S[g] = \frac{1}{2}\int dt \mathrm{Tr}\left(g^{-1}\frac{dg}{dt}g^{-1}\frac{dg}{dt}\right). \tag{5.1.3}$$

where $g = g(t)$ is the world line of the particle. Indeed let $\{x^\mu\}$ be a set of local coordinates on G then we can write

$$g^{-1}\frac{dg}{dt} = R^a_\mu(x)\frac{dx^\mu}{dt}t_a \tag{5.1.4}$$

where as before $\{t_a\}$ is a basis of the Lie algebra g. Inserting this into eq.(5.1.3) we find that it reduces to an action of the form (5.1.2) with

$$h_{\mu\nu}(x) = R^a_\mu(x) R^b_\nu(x) \mathrm{Tr}(t_a t_b) \tag{5.1.5}$$

The group manifold G together with the metric h is a Riemann manifold of constant curvature. This system will be our starting point.

The action (5.1.3) satisfies the following identity which is a finite version of the Polyakov-Wiegman identity

$$S[gh] = S[g] + S[h] + \int dt \mathrm{Tr}\left(g^{-1} \frac{dg}{dt} \frac{dh}{dt} h^{-1} \right), \tag{5.1.6}$$

from which one immediately deduces the equations of motion,

$$\frac{d}{dt}\left(g^{-1} \frac{dg}{dt} \right) = \frac{d}{dt}\left(\frac{dg}{dt} g^{-1} \right) = 0. \tag{5.1.7}$$

The action (5.1.3) is invariant under the group $G \times G$ where if $(h_1, h_2) \in G \times G$ then

$$g(t) \mapsto h_1 g(t) h_2 \tag{5.1.8}$$

Also note that the phase space of a particle moving on G is the cotangent bundle T^*G of G which means that the true phase space of the free particle is

$$T^*G/(G \times G) \tag{5.1.9}$$

The invariance of the system under $G \times G$ leads to the conserved quantities $J^a, \bar{J}^a$ defined by

$$J \equiv \frac{dg}{dt} g^{-1} \equiv J^a t_a \quad \text{and} \quad \bar{J} \equiv g^{-1} \frac{dg}{dt} \equiv \bar{J}^a t_a. \tag{5.1.10}$$

The equations that express the conservation of these currents in time coincide with the equations of motion of the system, so fixing the values of these conserved quantities completely fixes the orbit of the particle once its position on $t = 0$ is specified. The conserved quantities form a Poisson algebra [83]

$$\begin{aligned} \{J^a, J^b\} &= f^{ab}_{\ c} J^c \\ \{\bar{J}^a, \bar{J}^b\} &= f^{ab}_{\ c} \bar{J}^c \\ \{J^a, \bar{J}^b\} &= 0 \end{aligned} \tag{5.1.11}$$

This is precisely the Kirillov Poisson algebra we used as a starting point for the construction of finite W algebras in chapter 3. These were obtained by imposing constraints on the Poisson algebra (5.1.11), and we want to do the same here to get systems with finite W symmetry. In terms of the decomposition $g = g_- \oplus g_0 \oplus g_+$, these constraints were $\pi_+(J) = t_+$, where $\pi_\pm$ are the projections on $g_\pm$. Here we want to impose the same constraints, together with similar constraints on $\bar{J}$,

$$\pi_+(J) = t_+ \quad \text{and} \quad \pi_-(\bar{J}) = t_-. \tag{5.1.12}$$

Let $G_\pm$ denote the subgroups of G with Lie algebra $g_\pm$, and G_0 the subgroup with Lie algebra g_0, then almost every element g of G can be decomposed as $g_- g_0 g_+$, where $g_{\pm,0}$ are elements of the corresponding subgroups, because G admits a generalized Gauss decomposition $G = G_- G_0 G_+$*. Inserting $g = g_- g_0 g_+$ into (5.1.12) we find

$$\begin{aligned} g_0^{-1} t_+ g_0 &= \frac{dg_+}{dt} g_+^{-1} \\ g_0 t_- g_0^{-1} &= g_-^{-1} \frac{dg_-}{dt}. \end{aligned} \tag{5.1.13}$$

In the derivation of these equations one uses that $\pi_+(g_- t_+ g_-^{-1}) = t_+$, and a similar equation with π_- and t_-, which follow from the fact that $t_\pm$ have degree ± 1. The constrained currents look like

$$\begin{aligned} J &= g_- \left(t_+ + \frac{dg_0}{dt} g_0^{-1} + g_0 t_- g_0^{-1} \right) g_-^{-1}, \\ \bar{J} &= g_+^{-1} \left(t_- + g_0^{-1} \frac{dg_0}{dt} + g_0^{-1} t_- g_0 \right) g_+. \end{aligned} \tag{5.1.14}$$

The equations of motion now become

$$\begin{aligned} 0 = g_-^{-1} \frac{dJ}{dt} g_- &= \frac{d}{dt} \left(\frac{dg_0}{dt} g_0^{-1} \right) + [g_0 t_- g_0^{-1}, t_+], \\ 0 = g_+ \frac{d\bar{J}}{dt} g_+^{-1} &= \frac{d}{dt} \left(g_0^{-1} \frac{dg_0}{dt} \right) + [t_-, g_0^{-1} t_+ g_0]. \end{aligned} \tag{5.1.15}$$

Note that these equations describe a particle moving on G_0 in some background potential, i.e. the system is no longer free.

If we consider again the principal sl_2 embedding then g_0 is the Cartan subalgebra. Let $\{H_i\}$ be a basis of the CSA then any element of g_0 can be written as

$$\exp(q^i H_i) \tag{5.1.16}$$

*Strictly speaking $G_- G_0 G_+$ is only dense in G but we will ignore this subtlety in the remainder

Inserting this into (5.1.15) we find that they reduce to

$$\frac{d^2q_i}{dt^2} + \exp(K_{ij}q_j) = 0 \tag{5.1.17}$$

which are exactly the Toda equations (5.1.1). For nonprincipal embeddings the equations (5.1.15) are more complicated matrix equations. For obvious reasons we call these 'generalized finite Toda systems'.

It is easy to find an action that has (5.1.15) as its equations of motion. It reads

$$S[g_0] = \frac{1}{2}\int dt \mathrm{Tr}\left(g_0^{-1}\frac{dg_0}{dt}g_0^{-1}\frac{dg_0}{dt}\right) - \int dt \mathrm{Tr}\left(g_0 t_- g_0^{-1} t_+\right). \tag{5.1.18}$$

This generalized Toda action describes a particle moving on G_0 in some background potential. Two commuting copies of the finite W algebra leave the action (5.1.18) invariant and act on the space of solutions of the equations of motion (5.1.15).

The general solution of the the equations of motion of the free particle is given by

$$g(t) = h_1 e^{tX} h_2 \tag{5.1.19}$$

where $h_1, h_2 \in G$ and $X \in g$. From this follows that the general solution of the equations (5.1.15) can be constructed as follows: Let $h_0^{(1)}, h_0^{(2)}$ be elements of G_0. Let X_0 be an arbitrary element of g_0. If $g_0(t)$ is defined by the Gauss decomposition

$$g_-(t)g_0(t)g_+(t) = h_0^{(1)} \exp t(X_0 + (h_0^{(1)})^{-1}t_+ h_0^{(1)} + h_0^{(2)} t_-(h_0^{(2)})^{-1})\ h_0^{(2)}, \tag{5.1.20}$$

then $g_0(t)$ is the most general solution of (5.1.15).

This concludes our presentation of generalized finite Toda systems.

5.2 Generalized Toda field theory

It is also possible to obtain Toda field theories by the same kind of arguments that we used in the previous section. For completeness we shall briefly describe this reduction in this section.

Consider the Wess-Zumino-Witten action

$$S[g] = -\frac{1}{4\lambda}\int_\Sigma d^2\xi\ \mathrm{Tr}(g^{-1}\partial_\mu g g^{-1}\partial^\mu g) + k\Gamma(g) \tag{5.2.21}$$

where g is a G valued matter field, λ, k are dimensionless coupling constants and $\sigma = S^1 \times \mathbf{R}$. The first term of the action (5.2.21) is a nonlinear sigma model and the second term is called the Wess-Zumino term

$$\Gamma[g] = \frac{1}{24\pi} \int_B \ \mathrm{Tr}((g^{-1}dg)^3) \tag{5.2.22}$$

where B is a three dimensional manifold such that $\partial B = \Sigma$. The Wess-Zumino term Γ is only well defined if k is an integer. The action $S[g]$ is conformally invariant for k positive and

$$\lambda^2 = \frac{4\pi}{k} \tag{5.2.23}$$

In fact the action (5.2.21) is only called the WZW action for this value of λ.

The WZW action possesses a left and right affine symmetry. As in the case of the free particle moving on G the field equations coincide with the conservation laws of the Noether currents associated to these symmetries. They read

$$\partial_- J = 0 \ \text{ and } \ \partial_+ \bar{J} = 0 \tag{5.2.24}$$

where the conserved currents J and $\bar{J}$ are given by

$$J = \partial_+ g g^{-1} \ \text{ and } \ \bar{J} = g^{-1}\partial_- g \tag{5.2.25}$$

In order to describe the reduction [68, 82] to Toda theory we again introduce (as in the previous chapter) the constraints

$$\pi_-(J) = t_- \ \text{ and } \ \pi_+(\bar{J}) = t_+ \tag{5.2.26}$$

Here again $\pi_{\pm,0}$ are the projection operators projecting onto $g_{\pm,0}$ respectively. Again using the generalized Gauss decomposition

$$g(x^+, x^-) = g_+(x^+, x^-) g_0(x^+, x^-) g_-(x^+, x^-) \tag{5.2.27}$$

we find that the constraints are equivalent to

$$\begin{aligned} g_+^{-1}\partial_- g_+ &= g_0 t_+ g_0^{-1} \\ \partial_+ g \, g_-^{-1} &= g_0^{-1} t_- g_0 \end{aligned}$$

and the constrained currents can be written as

$$\begin{aligned} J &= g_+(\partial_+ g_0\, g_0^{-1} + t_-)g_+^{-1} + \partial_+ g_+\, g_+^{-1} \\ \bar{J} &= g_-^{-1}(g_0^{-1}\partial_- g_0 + t_+)g_- + g_-^{-1}\partial_- g_- \end{aligned}$$

Now writing the WZW equation of motion as

$$[\partial_+ - J, \partial_- - 0] = 0 \tag{5.2.28}$$

and conjugating this equation by $g_+^{-1}(x^+, x^-)$ we see that (5.2.28) is equivalent to

$$\partial_-(\partial_+ g_0\, g_0^{-1}) = [t_-, g_0 t_+ g_0^{-1}] \tag{5.2.29}$$

which is the equation of motion of a generalized Toda field theory [82, 3, 68].

A particular case is again when the sl_2 embedding is principal. We then have

$$g_0(x^+, x^-) = \exp(\sum_i \phi_i(x^+, x^-)H_i) \tag{5.2.30}$$

where H_i is a basis of the Cartan subalgebra of g. Substituting (5.2.30) into (5.2.28) we find that it reduces to

$$\partial_+\partial_-\phi_i + \exp(\sum_j K_{ij}\phi_j) = 0 \tag{5.2.31}$$

which is the standard Toda field equation.

5.3 Covariant KdV type hierarchies

The nonlinear Poisson algebras $W(\bar{g}; t_0)$ constructed in the previous chapter can be interpreted in two ways: as current algebras of certain classical field theories and as the second Hamiltonian structures of certain integrable hierarchies of evolution equations. The former picture we developed in the previous section and it is the purpose of this section to construct the integrable hierarchies associated to an interesting subclass of W algebras. For this we shall closely follow the approach of Drinfeld and Sokolov (see chapter 2).

Consider the case $g = sl_{2N}$ where $N = 1, 2, \ldots$ and take the defining vector of the sl_2 embedding to be

$$t_0 = \frac{1}{2}\mathrm{diag}(1, \ldots, 1, -1, \ldots, -1) \tag{5.3.32}$$

where there are N entries 1 and N entries -1. With respect to the sl_2 subalgebra determined by this defining vector the fundamental representation of sl_{2N} branches into a direct sum of N two dimensional representations of sl_2 (doublets) while the adjoint representation branches as follows (in the notation of chapter 3)

$$\underline{ad}_{2N} \to (N^2-1)\underline{1}_2 \oplus N^2\underline{3}_2 \tag{5.3.33}$$

From this follows that the centralizer C of the sl_2 embedding is isomorphic to sl_N and that the resulting W algebra has an affine sl_N subalgebra. Furthermore, in (5.3.33) we see that there are N^2 triplets (i.e. $j=1$ representations of sl_2) from which follows that the W algebra will contain N^2 field of conformal weight 2. of course one of those will be the stress energy tensor.

The 'step up' operator t_+ of the sl_2 embedding determined by (5.3.32) is given by

$$t_+ = \begin{pmatrix} 0 & \mathbf{1} \\ 0 & 0 \end{pmatrix} \tag{5.3.34}$$

where $\mathbf{1}$ is the $N \times N$ unit matrix. According to the general prescription given in the previous chapter an general element $y \in \bar{g}_{fix}$ will be of the form

$$y(z) = \begin{pmatrix} A(z) & \mathbf{1} \\ T(z) & A(z) \end{pmatrix} \tag{5.3.35}$$

where A and T are $N \times N$ matrices. of course A is traceless, i.e. $A \in Map(S^1, sl_N)$ while $T \in Map(S^1, gl_N)$.

Let us now, in analogy with the standard DS description, construct the Lax operator. For this consider again the differential equation

$$(\partial - y)\Psi = 0 \tag{5.3.36}$$

where Ψ is a 2-vector whose entries ψ_i are $N \times N$ matrices. Here A acts on ψ_i in the adjoint representation, i.e. $A.\psi_i = [A, \psi_i] \equiv -ad_A\psi$. Eliminating the component ψ_2 equation (5.3.36) reduces to

$$(D^2 - T)\psi_1 = 0 \tag{5.3.37}$$

where $D = \partial + ad_A$. From this we find that the Lax operator is given by

$$L = D^2 - T \tag{5.3.38}$$

The covariant derivative D is a derivation, i.e.

$$D(PQ) = (DP)Q + PDQ \tag{5.3.39}$$

and has a formal inverse D^{-1} within the algebra of pseudo differential operators with matrix coefficients. The first few terms of D^{-1} are

$$D^{-1} = \partial^{-1} - ad_A\partial^{-2} + (ad_{\partial A} + ad_A^2)\partial^{-3} + \dots \tag{5.3.40}$$

It is easily checked that $DD^1 = D^{-1}D = 1$. In particular this means that $D^{-1}D(PQ) = PQ$. Using the derivation property of D and on defining $R = DQ$ we find

$$D^{-1}(PR) = PD^{-1}R - D^{-1}((DP)D^{-1}R) \tag{5.3.41}$$

Iterating this equation we find the following permutation rule

$$D^{-1}Q = \sum_{i=0}^{\infty}(-1)^i(D^iQ)D^{-i-1} \tag{5.3.42}$$

which is identical to the one for ∂^{-1}. From these considerations we find that we might as well work with the algebra of covariant pseudodifferential operators instead of with ordinary pseudodifferential operators.

Having found the Lax operator L we can now easily determine its formal root. It reads

$$L^{\frac{1}{2}} = D - \frac{1}{2}TD^{-1} + \frac{1}{4}(DT)D^{-2} + \dots \tag{5.3.43}$$

Using (5.3.40) one can easily check that in terms of ordinary pseudodifferential operators $L^{\frac{1}{2}}$ is given by

$$L^{\frac{1}{2}} = \partial + ad_A - \frac{1}{2}T\partial^{-1} + \frac{1}{4}(\partial T + \{ad_A, T\})\partial^{-2} + \dots \tag{5.3.44}$$

where $\{.,.\}$ here denotes the anticommutator.

The hierarchy of evolution equations is given by (see last section of chapter 2)

$$\frac{dL}{dt_{2k+1}} = [(L^{\frac{2k+1}{2}})_+, L] \qquad (k = 0, 1, 2, \dots) \tag{5.3.45}$$

where the + means that we are to consider only the positive power part (w.r.t. the covariant derivative). For the hierarchy (5.3.45) to be anywhere near completely integrable the different flows must commute, i.e they must correspond to commuting vector fields on the (infinite dimensional) phase space of the system. This is proved in the following

Lemma 7 *The different flows in the hierarchy (5.3.45) commute, i.e.*

$$\frac{d^2}{dt_i dt_j} L = \frac{d^2}{dt_j dt_i} L \tag{5.3.46}$$

Proof: It is not difficult to show [66] that if equation (5.3.45) holds then also

$$\frac{dL^{\frac{i}{2}}}{dt_{2k+1}} = [(L^{\frac{2k+1}{2}})_+, L^{\frac{i}{2}}] \tag{5.3.47}$$

Using this equation we find

$$\frac{d^2}{dt_i dt_j} = [[(L^{\frac{i}{2}})_+, L^{\frac{j}{2}}]_+, L] + [(L^{\frac{j}{2}})_+, [(L^{\frac{i}{2}})_+, L]] \tag{5.3.48}$$

Therefore

$$(\frac{d^2}{dt_i dt_j} - \frac{d^2}{dt_j dt_i})L = [[(L^{\frac{i}{2}})_+, L^{\frac{j}{2}}]_+ - [(L^{\frac{j}{2}})_+, L^{\frac{i}{2}}]_+ + [(L^{\frac{j}{2}})_+, (L^{\frac{i}{2}})_+], L] \tag{5.3.49}$$

where we used the Jacobi identity. From

$$[L^{\frac{i}{2}}, L^{\frac{j}{2}}] = 0 \tag{5.3.50}$$

it follows that

$$[(L^{\frac{i}{k}})_+, L^{\frac{j}{k}}]_+ = [(L^{\frac{j}{k}})_+, (L^i k)_-]_+ \tag{5.3.51}$$

Inserting this into eq. (5.3.49) we find the desired result. This proves the lemma .

The first two equations of the hierarchy (5.3.45) are

$$\frac{dT}{dt_1} = DT \tag{5.3.52}$$

$$\frac{dT}{dt_3} = \frac{1}{4}(D^3 T) - \frac{3}{4}\{DT, T\} \tag{5.3.53}$$

while the 'connection' components are constant w.r.t. all time variables, i.e.

$$\frac{dA}{dt_{2k+1}} = 0 \tag{5.3.54}$$

This can be seen as follows. The Lax equation was so constructed [66] that its right hand side is a differential operator of order 0. However, the left hand side has in the covariant case order 1 which means that the coefficient of the order 1 term must be zero. This coefficient is exactly

$$2\frac{d}{dt_{2k+1}} ad_A \tag{5.3.55}$$

Obviously the hierarchy constructed above is a covariant generalization of the KdV hierarchy. If we write $T = u + T^a I_a$ (where $\{I_a\}$ is a basis of sl_N) then equation (5.3.52) becomes

$$\begin{aligned} \frac{du}{dt_1} &= \partial u \\ \frac{dT^a}{dt_1} &= \partial T^a + f^a_{bc} T^b A^c \end{aligned} \tag{5.3.56}$$

while eqn.(5.3.53) reads

$$\begin{aligned} \frac{du}{dt_3} &= \frac{1}{4}\partial^3 u - \frac{3}{2} u \partial u - \frac{3}{4N}\partial(g_{ab} T^a T^b) \\ \frac{dT^a}{dt_3} &= \frac{1}{4}\partial T^a - \frac{3}{2}\partial(T^0 T^a) \end{aligned} \tag{5.3.57}$$

Obviously the first of these equations is the ordinary KdV equation plus a term that couples the field u to the nondiagonal fields T^a. Using eq.(5.3.45) and the explicit form of the Lax operator it is easy to construct the other elements of the covariant KdV hierarchy.

Integrable evolution equations like the KdV equation are characterized by the fact that they have infinitely many conserved quantities. This property, which signals soliton like solutions, is of fundamental importance. We will now show that the covariant KdV hierarchy constructed above also admits infinitely many conserved quantities.

Theorem 12 *Let $Res(\sum_i A_i D^i) = A_{-1}$, then the quantities*

$$H_{2k+1} = \int Tr\, Res(L^{\frac{2k+1}{2}})\, dz \tag{5.3.58}$$

are conserved w.r.t. all flows in the hierarchy.

Proof: Consider

$$Res[AD^k, BD^l] \tag{5.3.59}$$

This quantity is 0 if $l, k > 0$ or $l, k < 0$. The only interesting case is $k > 0, l < 0$. Take $l = -p$ where $p > 0$ then $Res[AD^k, BD^{-p}] = 0$ if $p > k + 1$ and

$$(\frac{k}{p+1})(A(D^{k-p+1}B) + (-1)^{k-p}(D^{k-p+1}A)B) \tag{5.3.60}$$

Also note that if

$$g = (\frac{k}{p-1})\sum_{i=0}^{k-p}(-1)^i(D^iA)D^{k-p-i}B \tag{5.3.61}$$

then

$$Dg = (\frac{k}{p-1}(AD^{k-p+1}B + (-1)^{k-p}(D^{k-p+1}A)B) \tag{5.3.62}$$

which means that

$$\mathrm{Tr}(Dg) = \mathrm{Tr}Res[ADk, BD^{-p}] \tag{5.3.63}$$

However, $\mathrm{Tr}(Dg) = \partial\mathrm{Tr}(g)$ from which it follows that $\mathrm{Tr}Res[AD^k, BD^{-p}]$ is a total derivative. Using this and eq.(5.3.47) the theorem follows vspace5mm.

The first two Hamiltonians are given by

$$\begin{aligned} H_1 &= \frac{1}{2}\int u(z)\,dz \\ H_2 &= \frac{1}{2}\int (u^2 + \frac{1}{N}g_{ab}T^aT^b)\,dz \end{aligned}$$

Note that for $N = 1$ the hierarchy discussed above reduces to the ordinary KdV hierarchy. We have therefore obtained a generalization of this hierarchy and the W algebra $W(\bar{g}; t_0)$ is its second Hamiltonian structure.

It is possible to repeat the whole procedure oulined above for the the Boussinesq hierarchy. For this consider the sl_2 embedding into sl_{3N} determined by the following defining vector

$$t_0 = \mathrm{diag}(1, \ldots, 1, 0, \ldots, 0, -1, \ldots, -1) \tag{5.3.64}$$

where there are N entries 1,0 and -1. With respect to the sl_2 subalgebra determined by this defining vector the fundamental representation of sl_{3N} banches into a direct sum of N three dimensional sl_2 representations (triplets) while the adjoint representation branches as follows

$$\underline{ad}_{3N} \to (N^2 - 1)\underline{1}_2 \oplus N^2\underline{3}_2 \oplus N^2\underline{5}_2 \tag{5.3.65}$$

Again from this we see immediately that the W algebra $W(\bar{g};t_0)$ will contain an affine sl_N subalgebra, N^2 fields of conformal dimension 2 and N^2 fields of conformal dimension 3. According to the general prescription given in the previous chapter a general element $y \in \bar{y}_{fix}$ will be of the form

$$y(z) = \begin{pmatrix} A & T & W \\ \mathbf{1} & A & T \\ 0 & \mathbf{1} & A \end{pmatrix} \tag{5.3.66}$$

where A, T and W are again $N \times N$ matrices.

Again we can easily construct the Lax operator. It reads

$$L = D^3 - 2TD - (DT) - W \tag{5.3.67}$$

where as before $D = \partial + ad_A$. The formal cube root of this operator is given by

$$L^{1/3} = D - \frac{2}{3}TD^{-1} + \frac{1}{3}(\partial T - W)D^{-2} + \dots \tag{5.3.68}$$

As before the hierarchy is then given by

$$\frac{dL}{dt_{2k+1}} = [(L^{2k+1/3})_+, L] \tag{5.3.69}$$

Again let us give the first few elements of this hierachy. The first one reads

$$\begin{aligned} \frac{dT}{dt_1} &= DT \\ \frac{dW}{dt_1} &= DW \end{aligned}$$

and the second one is given by

$$\begin{aligned} \frac{dT}{dt_3} &= DW \\ \frac{dW}{dt_3} &= -\frac{1}{3}D^3T + \frac{4}{3}\{T, DT\} - \frac{4}{3}[T, W] \end{aligned} \tag{5.3.70}$$

which is a covariant version of the Boussinesq equation. As in the case of the covariant KdV hierarchy the connection components A are constant w.r.t. all times.

Again it is possible, in precisely the same way, to show that all the flows commute and also that the quantities

$$H_{2k+1} = \int \mathrm{Tr} Res(L^{2k+1}/3) \tag{5.3.71}$$

are conserved.

It is easy to construct covariant versions of all the elements of the Gelfand-Dickii hierarchy of hierarchies of which the KdV and Boussinesq hierarchies are the first two. For this one has to consider the sl_2 embedding into sl_{MN} with respect to which the fundamental representation of sl_{MN} branches into N M-dimensional sl_2 representations.

5.4 Discussion

In this chapter we have considered some applications of the theory developed in the previous chapters to the theory of integrable systems. This lead to generalized Toda theories in finite and infinite dimensions and also to new integrable hierarchies. The hierarchies we presented above are however related to a very special kind of sl_2 embedding and one might wonder as to the general structure. A possible answer to this question has been given in [99] where it was shown that to any conjugation class of the Weyl group associated a given simple Lie algebra one can associate an integrable hierarchy. This approach seems quite different from our approach in which sl_2 embeddings are the fundamental objects, but this is only superficial since sl_2 embeddings and conjugation classes of the Weyl group are in 1-1 correspondence for almost all simple Lie algebras. We therefore conjecture that the W algebras constructed in this thesis are the second Hamiltonian structures of the hierarchies of [99]. It should be noted here that in the approach of [99] the covariant derivative structure (which is of course related to affine subalgebras of the W algebra) is not clear.

Bibliography

[1] A. Kato *Classification of modular invariant partition functions in 2 dimensions*, Mod. Phys. Lett. **A2** (1987) 585; J. L. Cardy, *Conformal invariance*, in 'Phase transitions and critical phenomena', Vol.11, C. Domb and J. Lebowitz (Acedemic Press, 1986)

[2] L. Alvarez-Gaume, J. B. Bost, G. Moore, P. Nelson and C. Vafa, Phys. Lett. **B178** (1986) 41; CMP **112** (1987) 503

[3] W. P. G. van Veldhoven, F. A. Bais, *Generalizations of the Toda molecule*, Physica **139A** (1986) 326

[4] R. J. Baxter, *Exactly solved models in statistical mechanics*, Acedemic press London, 1982

[5] M. Henneaux, *Hamiltonian form of the path integral for theories with a gauge freedom*, Phys. Rep. **126** (1985) 1; L. Baulieu, *Perturbative gauge theories*, Phys. Rep. **129** (1985) 1

[6] M. Bershadski, H. Ooguri, CMP **126** (1989) 49

[7] J. Balog, L. Feher, P. Forgac, L. O'Raifeartaigh, A. Wipf, *Toda theory and W algebra from a gauged WZNW point of view*, Ann. Phys. **203** (1990) 76

[8] A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov, *Infinite conformal symmetry in two dimensional quantum field theory*, Nucl. Phys. **B241** (1984) 333

[9] J. de Boer and J. Goeree, *Covariant W gravity and its moduli space from gauge theory*, THU-92/14; Phys. Lett. **B274** (1992) 289; Nucl. Phys. **B381** (1992) 329; *The effective Action of W_3 gravity to all orders*, THU-92/93

[10] R. Bott, *Homogeneous vector bundles*, Ann. Math. **66** (1957) 203

[11] A. Cappelli, C. Trugenberger, G. R. Zemba, *Infinite symmetry in the quantum Hall effect*, CERN-TH 6516/92; A. Cappelli, G. Dunne, C. A. Trugenberger, G.

R. Zemba, *Conformal symmetry and universal properties of quantum Hall states* CERN-TH 6702/92.

[12] P. Ginsparg, *Applied conformal field theory*, in Les Houches, session XLIX, Fields, strings and critical phenomena, E. Brezin and J. Zinn-Justin eds. (Elsevier 1989); J. Bagger, *Basic conformal field theory*, Harvard Preprint.

[13] P. Goddard, A. Kent, D. Olive, *Virasoro algebras and coset space models*, Phys. Lett. **152** (1985) 88; CMP **103** (1985) 105

[14] K. Gawedzki, A. Kupiainen, Phys. Lett. **215B** (1988) 119; Nucl. Phys. **320** (1989) 625; P.Bowcock, Nucl. Phys. **B316** (1989) 80

[15] A. Deckmyn, Phys. Lett. B **298** (1993) 318

[16] J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509

[17] D. Olive, N. Turok, Nucl. Phys. **B215** (1983) 470; N. Ganoulis, P.Goddard, D. Olive, Nucl. Phys. **B205** (1982) 601; E. Witten, Phys. rev. Lett. **38** (1977) 121; F. A. Bais, H. A. Weldon, Phys. Rev. D **18** (1978) 591

[18] P. A. M. Dirac, *Lectures on quantum mechanics*, Belfer graduate school of science, Yeshiva Univ. of New York, 1964

[19] V. A. Fateev, S. L. Lukyanov, *The models of 2 dim CFT with Z_n symmetry*, Int. Journ. Mod. Phys. **A3** (1988) 507

[20] B. Feigin, E. Frenkel, Phys. Lett. **B246** (1990) 75; E. Frenkel, *W algebras and Langlands correspondence*, Harvard preprint 1991; *Affine Kac-Moody algebras at the critical level and quantum Drinfeld-Sokolov reduction*, Harvard thesis.

[21] G. Felder, J. Frohlich, G. Keller, CMP **124** (1989) 417

[22] P. Forgac, A. Wipf, J. Balog, L. Feher, L. O'Raifeartaigh, Phys. Lett. **227B** (1989) 214

[23] J. Frohlich, T. Keller, *Universality in quantum Hall systems*, Zurich preprint ETH-TH/90-26

[24] N. Woodhouse, *Geometric quantization*, Oxford University Press, Oxford 1980; J. Sniatycki, *Geometric quantization and quantum mechanics*, Springer-Verlag, New York, 1980; V. Guillemin, S. Sternberg, *Geometric quantization and multiplicities of group representations*, Invent. math. **67** (1982) 515

[25] D. Gepner, E. Witten, *String theory on group manifolds*, Nucl. Phys. **B278** (1986) 493; E. Witten, *Nonabelian bosonization in 2 dimensions*, CMP **92** (1984) 255

[26] I. A. Gelfand and L. A. Dickii, *Fractional powers of operators and Hamiltonian systems*, Finct. Anal. Appl. **10** (1976) 4; and *The resolvent and Hamiltonian systems*, Funct. Anal, Appl. *11* (1977) 2

[27] P. Goddard and D. Olive, *Kac-Moody and Virasoro algebras in relation to quantum physics*, Int. Journ. Mof. Phys. **A1** (1986) 303

[28] D. Gross, V. Periwal, *String perturbation theory diverges*, Phys. Rev. Lett. **60** (1988) 2105

[29] M. Green, J. Schwartz, E. Witten, *Superstring theory*, Cambridge University Press, Cambridge 1987

[30] B. Hatfield, Quantum field theory fo point particles and strings, Frontiers in physics, Addison-Wesley publishing company, Redwood City, 1992

[31] C. M. Hull, Nucl. Phys. **367** (1991) 731; Nucl. Phys. **B353** (1991) 707; Phys. Lett. **B269** (1991) 257

[32] M. Kaku, Introduction to superstrings, Graduate texts in contemporary physics, Springer Verlag, New York, 1988; Strings, conformal fields and Topology: An introduction, Graduate texts in contemporary physics, Springer Verlag, New York, 1991.

[33] V. G. Kniznik, A. B. Zamolodchikov, *Current algebra and Wess-Zumino model in two dimensions*, Nucl. Phys. **B247** (1984) 83

[34] B. Kostant, S. Sternberg, Ann. Phys. **176** (1987) 49

[35] A. Lopez and E. Fradkin, *The fractional quantum Hall effect and Chern-Simons gauge theories*, University of Illinois preprint PACS 73.20.Dx; A. Cappelli, C. A. Trugenberger, G. R. Zemba, *Large N limit in the quantum Hall effect*, preprint CERN-TH 6810/93; G. Nagao, *Toward a conformal field theory for the quantum Hall effect*, Preprint DOE-287-CPP-25; *A conformal field theory for the quantum Hall effect*, DOE-288-CPP-26; $c = 1$ conformal field theory and the fractional quantum Hall effect, NUS/HEP/920501; G. Moore, N. Read, *Nonabelions in the fractional quantum Hall effect*, Nucl. Phys. **B360** (1991) 362

[36] D. Luest and S. Theisen, *Lectures on string theory*, Lecture notes on physics 364, Springer Verlag Berlin 1989

[37] W. Nahm, *Conformal quantum field theory in 2 dimensions*, World scientific, to appear.

[38] N. P. Landsman, *Rieffel induction as generalized quantum Marsden-Weinstein reduction*, Preprint DAMTP-93-22; Preprint DAMTP to appear; Private communication.

[39] M. A. Olshanetsky and A. M. Perelomov, *Classical integrable finite dimensional systems related to Lie algebras*, Physics reports **71** (1981)313

[40] H. Ooguri, A. Sevrin, P. van Nieuwenhuizen, CMP **145** (1992) 515; K. Schoutens, A. Sevrin, P. van Nieuwenhuizen, Nucl. Phys. **B349** (1991) 791; Phys. Lett. **B243** (1991) 248; Nucl. Phys. **B364** (1991) 584; Nucl. Phys. **B371** (1992) 315

[41] E. Witten, *Two dimensional gravity and intersection numbers on moduli space*, Surveys in Differential geometry **1** (1991) 243; J. Horne, *Intersection theory and two dimensional gravity at genus 3 and 4*, Mod. Phys. Lett. **A5** (1990) 2127

[42] V. G. Kac, *Infinite dimensional Lie algebras*, Cambridge University Press, 1985

[43] V. G. Kniznik, A. M. Polyakov, A. B. Zamolodchikov, *Fractal structure of 2d quantum gravity*, Mod. Phys. Lett. **A3** (1988) 819

[44] T. Kimura, *Generalized classical BRST cohomology and reduction of Poisson manifolds*, To appear in CMP.

[45] M. Kontsevic, *Intersection theory on the moduli space of curves*, Funk. Anal. i Pril. **25** (1991) 50; *Intersection theory on the moduli space of curves and the matrix Airy function*, 30 Arbeitstagung Bonn, Max-Planck-Institut preprint MPI/91-47 and MPI/91-77

[46] J. L. Gervais, A. Neveu, *Novel triangle relation and absence of tachyons in Liouville string field theory*, Nucl. Phys. **B238** (1984) 125 and references therein.

[47] K. Osterwalder, R. Schrader, CMP **31** (1973) 83; CMP **42** (1975) 281; V. Glaser, CMP **37** (1974) 257

[48] W. Oevel, Mastersymmetries: weak action angle structure for Hamiltonian and non-Hamiltonian systems, Paderborn preprint

[49] M. A. Olshanetski and A. M. Perelomov, *Classical integrable finite dimensional systems related to Lie algebras*, Phys. Rep. **71** (1981) 313

[50] A. M. Polyakov, Mod. Phys. Lett. **A2** (1987) 893; Int. Journ. Mod. Phys. **A5** (1990) 833

[51] M. Dubois-Violette, Ann. Inst. Fourier **37** (1987) 45; D. Wilbour, J. M. Arms, *Reduction procedures for Poisson manifolds*, Washington Preprint.

[52] R. E. Prange, Chapter 1 in *The quantum Hall effect*, Graduate texts in contemporary physics, R. E. Prange, S. M. Girvin (eds.), Springer Verlag 1987

[53] *The quantum Hall effect*, Graduate texts in contemporary physics, R. E. Prange, S. M. Girvin (eds.), Springer Verlag 1987

[54] A. Reichl, A modern course in statistical physics, Edward Arnold publishing ltd, Texas, 1980

[55] B. Zwiebach, *Closed string field theory: Quantum action and the Batalin Vilkvisky equation*, Nucl. Phys. **B390** (1993) 33; W. Siegel, *Introduction to string field theory*, World scientific Singapore, 1988; A. Giveon, *On string field theory and effective actions*, Nucl. Phys. **B391** (1993) 229

[56] E. Witten, *Topological quantum field theory*, CMP **117** (1988) 353; *On the topological phase of two dimensional gravity*, Nucl. Phys. **B340** (1990) 281; J. Labastida, M. Pernici, E. Witten, *Topological gravity in two dimensions*, Nucl. Phys. **B310** (1988) 611; E. Verlinde, H. Verlinde, *A solution of two dimensional topological quantum gravity*, Nucl. Phys. **B348** (1991) 457

[57] K. Thielemans, Int. Journ. Mod. Phys. **C2** (1991) 787

[58] D. Friedan, Z. Qui, S. Shenker, Phys. Rev. Lett. **52** (1984) 1575.

[59] J. Goeree, Nucl. Phys. **B358** (1991) 737

[60] E. Witten, *Quantum field theory and the Jones polynomial*; CMP **121** (1989) 351

[61] E. Witten, *Coadjoint orbits of the Virasoro group*, CMP **114**(1988)1

[62] A. B. Zamolodchikov, *infinite additional symmetries is two dimensional CFT*, Theor. Math. Phys. **65** (1985) 347

[63] P. Bouwknegt, K. J. Schoutens, 'W symmetry in CFT', CERN-TH.6583/92, ITP-SB-92-23, To be published in Physics reports

[64] I. Bakas, Comm. Math. Phys. **123** (1989) 627
P. Mathieu, Phys. Lett. **B208** (1988) 101

[65] P. Ginsparg, *Applied conformal field theory*, in Les Houches, Session XLIX, 1988, 'Fields, strings and critical phenomena', E. Brezin, J. Zinn-Justin, eds. (Elsevier, 1989); E. Brezin and V. Kazakov, Phys. Lett. **B236** (1990) 144; M. Douglas and S. Schenker, Nucl. Phys. **335** (1990) 635; D. J. Gross and A. Migdal, Phys. Rev. Lett. **64** (1990) 127; Nucl. Phys. **B340** (1990) 333; A. Bilal, '2d gravity from matrix models', John Hopkins Lectures, CERN-TH.5867/90

[66] V. Drinfeld, V. Sokolov, Journ. of Soviet Math. **30** (1984) 1975

[67] F. A. Bais, T. Tjin, P. van Driel, Nucl.Phys. **B357** (1991) 632

[68] L. O'Raifeartaigh, A. Wipf, Phys. Lett. **B251** (1990) 361
L. Feher, L. O'Raifeartaigh, P. Ruelle, I. Tsutsui, 'Generalized Toda theories and W algebras associated to integral gradings', ETH-TH/91-16, DIAS-STP-91-17; Phys. Lett. **B283** (1992) 243
J. Balog, L. Feher, P. Forgac, L. O'Raifeartaigh, A. Wipf, Phys. Lett. **B227** (1989) 214; Phys. Lett. **B244** (1990) 435
L. O'Raifeartaigh, P. Ruelle, I. Tsutsui, A. Wipf, 'W algebras for generalized Toda theories', dublin Preprint; L. Feher, L. O'Raifeartaigh, P. Ruelle, I. Tsutsui, 'On the completeness of the set of classical W algebras obtained from DS reductions', Bonn-HE-93-14.

[69] J. de Boer, J. Goeree, 'Covariant W gravity and its moduli space from gauge theory', THU-92/14; Phys. Lett. **B274** (1992) 289; Nucl. Phys. **B381** (1992) 329; 'The Effective Action of W_3 Gravity to All Orders', THU-92/33

[70] H. Ooguri, K. Schoutens A, Sevrin, P. van Nieuwenhuizen, Comm. Math. Phys. **145** (1992) 515; K. Schoutens, A. Sevrin, P. van Nieuwenhuizen, Nucl. Phys. **B349** (1991) 791; Phys. Lett. **B243** (1991) 248; Nucl. Phys. **B364** (1991) 584; Nucl. Phys. **B371** (1992) 315; CMP **124** (1989) 87; Int. Journ. Mod. Phys. **A6** (1991) 2891; 'Example of a QFT based on a nonlinear Lie algebra', in Festschrift on the occasion of M. Veltman's birthday, Eds. R. Akhoury et al.; A. Sevrin, P. van Nieuwenhuizen, 'Quantum gauging from classical gauging of nonlinear algebras', ITP-SB-92-67, LBL-33339; A. Sevrin, K. Thielemans, W. Troost, Preprint LBL-33778, To appear in Phys. Rev. **D**; A. Sevrin, K. Thielemans, W. Troost, Preprint LBL-33738.

[71] E. Frenkel, V. Kac, M. Wakimoto, Comm. Math. Phys. **147** (1992) 295;

[72] M. Rocek, Phys. Lett. **B255** (1991) 554

[73] K. J. Schoutens, A. Sevrin, P. van Nieuwenhuizen, Phys. Lett. **B255** (1991) 549

[74] T. Tjin, Phys. Lett. **B292** (1992) 60

[75] P. Bouwknegt, J. McCarthy, K. Pilch, Comm. Math. Phys. **131** (1990) 125

[76] R. Abraham, J. E. Marsden, 'Foundations of classical mechanics', second edition, The Benjamin/Cummings Publishing Company inc.

[77] M. Dubois-Violette, Ann. Inst. Fourier **37** (1987) 45
D. Wilbour, J. M. Arms, 'Reduction procedures for Poisson manifolds', Washington Preprint.

[78] K. Sundermeyer, Lecture Notes in Physics 169 (Springer-verlag, Berlin, 1982)

[79] E. B. Dynkin, Amer. Math. Soc. Transl. **6** (1967) 111

[80] L. Feher, L. O'Raifeartaigh, Ruelle, I. Tsutsui, A. Wipf, 'On the general structure of Hamiltonian reductions of WZW theory', DIAS -STP -91 -29, UdeM -LPN -TH -71/91

[81] A. Bilal, G. Gervais, Phys. Lett. **B206** (1988) 412; Nucl. Phys. **B314** (1989) 646; Nucl. Phys. **B318** (1989) 579
P. Mansfield, B. Spence, Nucl. Phys. **B362** (1991) 294

[82] A. N. Leznov, M. Saveliev, Lett. Math. Phys. **6** (1982) 505; Comm. Math. Phys. **89** (1983) 59

[83] G. Papadopoulos, 'The global phase space structure of the WZW model', Imperial TH/91-92/28

[84] Z. Bajnok, L. Palla, G. Takács, 'A_2 Toda theory in reduced WZNW framework and the representations of the W algebra', Budapest preprint

[85] V. Pasquier, *Two dimensional critical systems labelled by Dynkin diagrams*, Nucl. Phys. **B285** (1987) 162; J. Phys. A: Math. gen. *20* (1987) L1229

[86] B. Kostant, S. Sternberg, Ann. Phys. **176** (1987) 49

[87] A. W. Knapp, *Lie groups, Lie algebras and cohomology*, mathematical notes Princeton university press, Princeton, 1988.

[88] B. Kostant, Inv. Math. **48** (1978) 101

[89] F. Bais, P. Bouwknegt, M. Surridge, K. Schoutens, Nucl. Phys. **B304** (1988) 348

[90] R. Bott and L. W. Tu, 'Differential Forms in Algebraic Topology', Graduate Texts in Mathematics 82, Springer Verlag, New York, 1982

[91] J. McCleary, 'User's Guide To Spectral Sequences', Mathematics Lecture Series 12, Publish or Perish, Washington, 1985

[92] B. Kostant, Lect. Notes Math. **466** (1974) 101

[93] M. Bershadsky, Comm. Math. Phys. **139** (1991) 71

[94] W. P. G. van Veldhoven, F. A. Bais, Physica **139 A** (1986) 326

[95] B. A. Khesin, B. Z. Shapiro, Commun. Math. Phys. **145** (1992) 357.

[96] T. Tjin, P. van Driel, 'Coupled WZW-Toda models and covariant KdV hierarchies', preprint ITFA-91-04

[97] J. de Boer and T. Tjin, 'Quantization and representation theory of finite W algebras', To appear in Comm. Math. Phys.

[98] J. de Boer and T. Tjin, *The relation between quantum W algebras and Lie algebras*, To appear in CMP.

[99] M. F. de Groot, T. J. Hollowood, J. L. Miramontes, *Generalized Drinfeld-Sokolov hierarchies*, IASSNS-HEP-91/19,PUPT-1251; N. J. Burroughs, M. F. de Groot, T. J. Hollowood, J. L. Miramontes, *Generalized Drinfeld-Sokolov hierarchies 2*, PUPT-1263, IASSNS-HEP-91/42